

Monograph of *Coccinia* (Cucurbitaceae)

Norbert Holstein¹

¹ *Nees-Institute for Biodiversity of Plants, Meckenheimer Allee 170, 53115 Bonn, Germany*

Corresponding author: *Norbert Holstein* (holstein@lrz.uni-muenchen.de)

Academic editor: *Sandra Knapp* | Received 25 April 2012 | Accepted 23 September 2014 | Published 3 August 2015

Citation: Holstein N (2015) Monograph of *Coccinia* (Cucurbitaceae). *PhytoKeys* 54: 1–166. doi: 10.3897/phytokeys.54.3285

Abstract

This monograph deals with all 95 names described in the Cucurbitaceae genus *Coccinia* and recognizes 25 species. Taxonomic novelties are *Coccinia adoensis* var. *aurantiaca* (C.Jeffrey) Holstein, **stat. nov.**, *C. sessilifolia* var. *variifolia* (A.Meeuse) Holstein, **stat. nov.**, and *C. adoensis* var. *jeffreyana* Holstein, **var. nov.** For the 25 species 3157 collections were examined, of which 2024 were georeferenced to produce distribution maps. All species are distributed in sub-Saharan Africa with one species, *C. grandis*, extending from Senegal in West Africa east to Indonesia and being naturalized on Pacific Islands, in Australia, the Caribbean, and South America. *Coccinia* species are dioecious creepers or climbers with simple or bifid tendrils that occupy a range of habitats from arid scrubland, woodlands to lowland rainforest and mist forest. The corolla of *Coccinia* species is sympetalous, usually pale yellow to orange, and 1 to 4.5 cm long. Pollination is by bees foraging for pollen or nectar. After pollination, the developing ovary often exhibits longitudinal mottling, which usually disappears during maturation. All species produce berries with a pericarp in reddish colors (orange-red through to scarlet red), hence the generic name. The globose to cylindrical fruits contain numerous grayish-beige flat to lenticular seeds. Chromosome numbers are $2n = 20, 24$, and $22 + XX/XY$. Many *Coccinia* species are used for food, either as roasted tubers, greens as spinach, or the fruits as vegetables. Medicinal value is established in *C. grandis*, of which leaves and sap are used against diabetes.

Keywords

Cucurbitaceae, *Coccinia*, molecular phylogeny, biogeography, taxonomy, morphology, sex expression, useful plants

Table of contents

| | |
|------------------------------------------------------------------------------------|----|
| Introduction..... | 4 |
| Materials and methods | 4 |
| General morphology | 4 |
| Phylogenies | 5 |
| Distribution maps | 5 |
| Morphology and anatomy | 6 |
| Habit | 6 |
| Roots | 6 |
| Hypocotyl and shoots | 7 |
| Cotyledons..... | 9 |
| Leaves | 9 |
| Probracts and bracts | 13 |
| Extranuptial glands | 13 |
| Peduncles and pedicels | 14 |
| Perianth | 15 |
| Androecium | 17 |
| Pollen..... | 18 |
| Gynoecium | 19 |
| Female gametophyte development and embryology..... | 19 |
| Fruits | 19 |
| Seeds | 21 |
| Germination | 23 |
| Genome, chromosomes, and hybridization..... | 23 |
| Chromosomes and sex determination..... | 23 |
| Genome of <i>C. grandis</i> | 26 |
| Hybridization and crossing experiments | 26 |
| Plant – biotic environment interactions..... | 29 |
| Pollination | 29 |
| Seed dispersal | 30 |
| Interaction with ants | 31 |
| Diseases and parasites..... | 31 |
| Use, economic potential, and phytochemistry | 34 |
| Evolution and phylogeny..... | 36 |
| Identification of <i>Coccinia</i> species | 41 |
| Possible confusion with other genera..... | 41 |
| Characters for species discrimination..... | 41 |
| Key to <i>Coccinia</i> species..... | 41 |
| Taxonomic treatment | 54 |
| 1. <i>Coccinia abyssinica</i> (Lam.) Cogn. | 56 |
| 2a. <i>Coccinia adoensis</i> (Hochst. ex A.Rich.) Cogn. var. <i>adoensis</i> | 59 |
| 2b. <i>Coccinia adoensis</i> var. <i>aurantiaca</i> (C.Jeffrey) Holstein | 67 |

| | |
|-----------------------------------------------------------------------------------|-----|
| 2c. <i>Coccinia adoensis</i> var. <i>jeffreyana</i> Holstein | 70 |
| 3. <i>Coccinia barteri</i> (Hook.f.) Keay | 72 |
| 4. <i>Coccinia grandiflora</i> Cogn. | 75 |
| 5. <i>Coccinia grandis</i> (L.) Voigt | 78 |
| 6. <i>Coccinia heterophylla</i> (Hook.f.) Holstein | 89 |
| 7. <i>Coccinia hirtella</i> Cogn. | 92 |
| 8. <i>Coccinia intermedia</i> Holstein..... | 94 |
| 9. <i>Coccinia keayana</i> R.Fern. | 95 |
| 10. <i>Coccinia longicarpa</i> Jongkind | 97 |
| 11. <i>Coccinia mackenii</i> Naudin ex C.Huber | 99 |
| 12. <i>Coccinia megarrhiza</i> C.Jeffrey | 102 |
| 13. <i>Coccinia microphylla</i> Gilg | 104 |
| 14. <i>Coccinia mildbraedii</i> Gilg | 107 |
| 15. <i>Coccinia ogadensis</i> Thulin | 109 |
| 16. <i>Coccinia pwaniensis</i> Holstein | 111 |
| 17. <i>Coccinia quinqueloba</i> (Thunb.) Cogn. | 113 |
| 18. <i>Coccinia racemiflora</i> Keraudren | 114 |
| 19. <i>Coccinia rehmannii</i> Cogn. | 115 |
| 20. <i>Coccinia samburuensis</i> Holstein | 122 |
| 21. <i>Coccinia schliebenii</i> Harms | 123 |
| 22. <i>Coccinia senensis</i> (Klotzsch) Cogn..... | 124 |
| 23a. <i>Coccinia sessilifolia</i> (Sond.) Cogn. var. <i>sessilifolia</i> | 129 |
| 23b. <i>Coccinia sessilifolia</i> var. <i>variifolia</i> (A.Meeuse) Holstein..... | 132 |
| 24. <i>Coccinia subsessiliflora</i> Cogn. | 133 |
| 25. <i>Coccinia trilobata</i> (Cogn.) C.Jeffrey..... | 135 |
| Insufficiently known taxa | 137 |
| Dubious names | 138 |
| Invalid names..... | 140 |
| Excluded taxa | 145 |
| Acknowledgements..... | 150 |
| References | 151 |
| Supplementary material 1..... | 165 |
| Supplementary material 2..... | 165 |

Introduction

Coccinia Wight & Arn. comprises 25 species and is the 11th largest of the 97 genera of the Cucurbitaceae (Schaefer and Renner 2011a). Especially in the 19th century, it drew gardeners' attention probably because of its striking fruits (André 1900; Edwards 1815; Huber 1864; 1865; Koch 1865; Sims 1816). All species are dioecious, and one species, *C. grandis* (L.) Voigt, has heteromorphic sex chromosomes and therefore has been studied cytologically (Agarwal and Roy 1975; 1983; 1984; Bhaduri and Bose 1947; Chakravorti 1948; Datta 1988; Kumar and Deodikar 1940; Kumar and Vishveshwaraiah 1952; Roy 1974; Roy and Roy 1971b). The last complete taxonomic treatment of *Coccinia* is by Cogniaux (1881), more than 130 years ago. Since then, 16 new species have been described, and the genus has only been revised regionally (Hutchinson et al. 1954; Jeffrey 1967; 1978; 1995; Jeffrey and Fernandes 1986; Kéraudren-Aymonin 1975a; Kéraudren 1967; Meeuse 1962). The position of *Coccinia* in the Benincaseae has been confirmed by molecular data (Kocyan et al. 2007; Schaefer and Renner 2011b), and the monophyly has been tested with almost complete species sampling by Holstein and Renner (2011b).

The delimitation of *Coccinia* from other genera is difficult. The scarlet-red fruits to which the genus name – *Coccinia* from Latin *coccineus* – refers are also found in other African genera, such as *Eureiandra* Hook.f. Therefore, it is not surprising that early botanists described several species now considered to belong to *Coccinia* in other genera (*Cephalandra* Schrad. ex Eckl. & Zeyh., *Physedra* Hook.f., and *Staphylosyce* Hook.f.), or species described in *Coccinia* now belong to different genera. In all, 113 names at various ranks have been proposed for what are here considered 25 species. The species concepts in the present revision are based on 3157 herbarium collections and fieldwork in Tanzania, geo-referencing of 2024 collections and cultivation of 10 species in the greenhouse. In combination, plastid and nuclear data obtained for multiple accessions representing most species and ecological information coming from the mapping effort provide a modern understanding of the evolution and species relationships in *Coccinia*.

Materials and methods

General morphology

During this study, the present author examined 3157 herbarium collections from 39 herbaria from (physical or digital) loans or *in situ* (B, BM, BR, BRI, C, CANB, CBG, COI, DSM, EA, FR, FT, G, GAT, GOET, H, HBG, HEID, JE, K, L, LISC, LISU, M, MO, MSB, NHT, P, PERTH, PR, PRC, S, U, UBT, UPS, W, WAG, Z, ZT). Additional collections were obtained via personal communication (BO, DNA, TUB, US), from web pages such as like JStor Plant Science (JPS, <http://plants.jstor.org/>), Chinese Virtual Herbarium (CVH, <http://www.cvh.org.cn/>), and homepages of the following herbaria: A, AAU, BAR, CAY, FLAS, FSU, FTG, GH, HAST, NTUF, NY, PDA, TAIF, and USE.

New collections were added if the photograph allowed identification or if misidentification appeared to be unlikely (esp. *C. grandis* collections from the Pacific area), while duplicates were added without visual inspection of the specimen photo. Online availability of specimen images is mentioned in the list of exsiccatae (Suppl. material 1). Ten species were cultivated in greenhouses of Munich Botanical Garden: *C. abyssinica* (Lam.) Cogn. (1 origin, 4 individuals), *C. adoensis* var. *jeffreyana* (1 origin, 4 individuals), *C. grandiflora* Cogn. (1 origin, 1 individual), *C. grandis* (3 origins with 2, 61, 3 individuals, respectively), *C. hirtella* Cogn. (1 origin, 12 individuals), *C. megarrhiza* C.Jeffrey (1 origin, 8 individuals), *C. microphylla* Gilg (2 origins, 2 and 1 individual, respectively), *C. rehmannii* Cogn. with two varieties (2 origins, each 3 individuals), *C. sessilifolia* (Sond.) Cogn. var. *sessilifolia* (1 origin, 8 individuals), and *C. trilobata* (Cogn.) C.Jeffrey (2 origins, each 3 individuals). The present author performed crossing experiments among eight of these species. Morphological features were documented photographically and in the form of vouchers, with 47 collections deposited in M. Field data were obtained on a trip to NE Tanzania in 2009 resulting in 28 *Coccinia* collections.

Phylogenies

For this monograph the phylogenetic data of Holstein and Renner (2011a; 2011b) were augmented with 20 new sequences from 8 accessions (GenBank accession numbers are given in Suppl. material 2), and new phylogenies were calculated using RAxML v. 7.2.6 (Stamatakis 2006) and MrBayes v. 3.2 (Ronquist et al. 2012). The substitution model was GTR+ Γ as used before, and 1000 ML replicates were used to infer statistical support for the nodes via bootstrapping. For Bayesian analysis, four chains were run with 2,000,000 generations, with a sampling frequency of 1000. The first 25% of the trees were discarded as burn-in, and the rest were plotted as a 50% majority rule consensus tree using FigTree 1.3.1 (<http://tree.bio.ed.ac.uk/software/figtree/>). Gaps in the plastid matrix occurring in more than one accession were coded as “0”, “1”, or “?”, with “?” when data were missing or when shorter gaps were coded in the same place, but in different accessions.

Distribution maps

Of the examined collections, 2024 were geo-referenced and mapped in Google Earth (Google Inc., Mountain View, CA, USA). Cultivated plants were geo-referenced according to the original collection site. If collecting sites were given as distances from locations, a path along major roads was used, beginning from the center of the starting location. Collecting sites were geo-referenced according to the description even if coordinates were given on the label, except for cases in which the coordinates were clearly derived from GPS or if the description did not allow further improvement. Location names were cross-validated from printed maps and then imported into DIVA-GIS 7.1.6.2 (<http://www.diva-gis.org>). If collecting sites of specimens appeared to be too

useful to ignore in the distribution maps despite the large uncertainty of the position (radius > 5 km) or if the collecting site was only given as “nearby” a distinct locality, then geo-referenced coordinates are given in brackets. Political administrative borders were taken from GADM v.1 (Jan 2009) or v.2 (Jan 2012) (<http://www.gadm.org/>) and elevation data (1 km resolution) from CGIAR Consortium for Spatial Information (Jarvis et al. 2008). The geodetic datum for the maps is WGS84; the projection in each case is equirectangular. Coordinates are given in decimal degrees in Suppl. material 1 with WGS84 as geodetic datum.

Morphology and anatomy

Habit

Coccinia species are perennial climbers or creepers. The lignification of the mature shoots differs among the species from unlignified to completely lignified. Climbing is enabled by tendrils, which are either simple or bifid. Tendril development in young plants is delayed and emerges in *C. abyssinica* after the 6th node (Getahun 1974a). The tendril arms are only rarely equally sized, as one is usually much smaller; true dichotomy of tendrils is unknown from *Coccinia*. Whether a species has simple or bifid tendrils is often not fixed, but there is a strong predominance of one kind. Bifid tendrils regularly occur or are predominant in *C. grandiflora* Cogn., *C. heterophylla* (Hook.f.) Holstein, *C. hirtella* Cogn., *C. intermedia* Holstein, *C. mackenii* Naudin ex C. Huber, *C. mildbraedii* Gilg, *C. racemiflora* Keraudren, *C. schliebenii* Harms, and in some forms of *C. barteri* (Holstein and Renner 2011b). Strikingly, *Coccinia* species with bifid tendrils occur in rather humid habitats. This suggests an adaptive advantage, because more tendril arms increase stability, as the leaves of rainforest species are larger, coriaceous, and thus heavier than leaves of species from drier habitats. Some species are regularly described as having simple tendrils in floristic treatments, but they may bear bifid tendrils such as *C. sessilifolia* (N. Holstein 13) and *C. senensis* (H.J.E. Schlieben 5745 in B, K, and MO). *Coccinia adoensis* has bifid tendrils even in some type specimens (e.g., G.H.W. Schimper 166 in BR8886781 and on the sheet with a drawing in K) and is still listed as simple-tendrilled. All three species with this polymorphism, however, have predominantly simple tendrils. Interestingly, these species are also closely related to species with predominantly bifid tendrils: *C. sessilifolia* with *C. hirtella* and *C. mackenii*, and *C. adoensis* with *C. grandiflora* and *C. schliebenii*.

Roots

Coccinia species have perennial roots. Most (if not all) species are woody at the base, and most of them produce hypocotyl tubers (Fig. 1a). Some species, such as *Coccinia adoensis* and *C. grandiflora* (and most likely also *C. senensis* (Klotzsch) Cogn.




Figure 1. a Young plant of *C. grandis*. The hypocotyl is already thickened and lignifies later-on to a tuber. The cotyledons are glabrous, have an entire margin and an obtuse apex **b** Adventitious root on *C. grandis*; the beige structure to the left is a dry tendril.

and *C. schliebenii*), however, produce globular subterranean root tubers, much like potatoes, but smaller in size (Holstein and Renner 2011b; Zimmermann 1922b; pers. observ.). Root tubers in *Coccinia adoensis* are likely to be an adaptation to fire, as this species predominantly occurs in woodlands. In contrast to rather mild fires in semi-arid savannas with less inflammable biomass, woodland fires produce temperature rises of 60 °C in 0–3 cm depth (Bradstock and Auld 1995; Gignoux et al. 1997), so vegetative buds near the ground (hemicryptophytes) might be damaged, whereas root tubers (geophytes) have a higher chance of survival.

Coccinia grandis and *C. barteri* produce adventitious roots if stems touch the soil (Fig. 1b). *Coccinia hirtella*, *C. sessilifolia* and their F1 hybrids with *C. grandis* appear to lack this ability (pers. observ.). Adventitious roots also occur along the hypocotyl of *C. abyssinica* seedlings (Getahun 1974a). There is barely any research on root anatomy, solely Getahun (1974a) reports tetrarch vascular bundles in the primary roots of *C. abyssinica* seedlings and di- to triarch bundles in secondary roots.

Hypocotyl and shoots

Many species, such as *C. abyssinica*, *C. grandis*, *C. hirtella*, *C. megarrhiza*, *C. microphylla*, *C. rehmannii*, *C. sessilifolia*, and *C. trilobata*, produce a lignified tuber that is derived from the hypocotyl (Zimmermann 1922b; pers. observ., Fig. 1a). The tuber, at least of some species, contains starch as a storage nutrient (Getahun 1974b). It develops during the first season, and lignification may begin as soon as the appearance of the first tendrils, such as in *C. abyssinica* (N. Holstein 132). Some species, such as *C. adoensis* and *C. grandiflora* (and most likely also *C. senensis* and *C. schliebenii*) do not produce hypocotyl tubers but root tubers. In *C. adoensis* var. *jeffreyana* the hypocotyl is



Figure 2. **a** Female flower bud of *C. microphylla* (N. Holstein et al. 90); picture taken during the dry season. The stem is completely lignified, and only green short shoots are produced **b** Male plant of *C. sessilifolia*. The stem is glaucous and does not lignify. Unusually, the bract is 3-lobate leaf-like.

minute (N. Holstein 130), which prevents the development of a tuber. Whether West and Central African forest species produce tubers is unknown.

Each plant produces one to several shoots, which can persist or die back completely during the dry season or due to fire or grazing. *Coccinia microphylla* shoots can lignify completely and produce short green branches with flowers and small leaves during the dry season (Fig. 2a), whereas shoots of *C. sessilifolia* do not lignify at all (pers. observ. from greenhouse cultivation over 4 years). The shoots of *C. grandis* can become slightly succulent. The length of the shoots varies from 70 cm in *C. microphylla* to 20 m in *C. grandiflora* and *C. mildbraedii*. Zimmermann (1922a) reports a stem of *C. grandiflora* being 6 cm in diameter. Usually, the bark of the hypocotyl tuber and the shoots is grayish in color. Fresh shoots and twigs are usually deep green to brownish green, sometimes speckled with pale to whitish pustules. In *C. abyssinica* and *C. megarrhiza* the shoots and tendrils can turn purple during maturity. *Coccinia sessilifolia* produces glaucous shoots that bear a waxy bloom (Fig. 2b). The indumentum of *Coccinia* species, if present, is composed of simple, oligo- to multicellular eglandular trichomes up to 2 mm in length. The long trichomes consist of oblong cells that may appear articulate when dried (Fig. 3a). Shorter trichomes can be lineal to conical (Fig. 3b). Sometimes, trichomes have a thickened base that appear warty when the trichomes break off. The density of the trichomes is often increased on the nodes. Trichome type and length on shoots are like those of the abaxial surface of the petioles, but usually less dense. Young shoots often exhibit short (< 0.5 mm), weak trichomes, even in species that are later glabrous, e.g., in *C. grandis* or *C. sessilifolia*. Glandular trichomes are rare, few-celled, not visible with the naked eye and have been found, e.g., in *C. grandiflora* and *C. grandis* (pers. observ.; Thanki 1989; Fig. 3c). Glandular trichomes are also observed in young stems of *C. abyssinica* (Getahun 1974a), which are usually covered with long multicellular eglandular trichomes.




Figure 3. Trichomes on *Coccinia* species. **a** Node of *C. adoensis* var. *jeffreyana* (J.C. Lovett 1197). The black bar equals 1 mm. **b** Part of a leaf *C. adoensis* var. *jeffreyana* (A.F. Stolz 504). The white bar equals 1 mm. **c** Glandular trichome from *C. grandis*.

Cotyledons

Zimmermann (1922b) reports epigeous cotyledons for *C. grandiflora* and *C. grandis*, of which the latter is confirmed by personal observations (Fig. 1a). Epigeous cotyledons also occur in *C. abyssinica*, *C. adoensis* var. *jeffreyana*, *C. microphylla*, *C. rehmannii* aff. var. *littoralis*, and *C. sessilifolia*. The hypocotyl and cotyledons of all observed taxa are glabrous. The cotyledons are elliptical to obovate and have an entire margin. The cotyledons are slightly fleshy and green, which is also observed in those of *C. abyssinica* (Getahun 1974a), and the cotyledonous apex has a pale marking and is obtuse to retuse. Getahun reports that the prominent veins and the margins on the lower cotyledon surface in *C. abyssinica* are covered with multicellular trichomes. However, prominent veins in *C. abyssinica* cotyledons cannot be confirmed, and if multicellular trichomes occur, then they are not visible to the naked eye. The first normal leaf in this species, however, emerges in the axilla of the cotyledons (N. Holstein 132, Fig. 4a), and thus might have been confused.

Leaves

The leaves of *Coccinia* species are simple, alternate, and paired with a tendril on each node, except for the first nodes (Figs 1a, 4a, b). Leaves of all species are petiolate, except for *C. sessilifolia* var. *sessilifolia*, which only develops petioles when young (N. Holstein 131, Fig. 4b) or rarely sessile leaves when older; full petioles in this species are only realized in *C. sessilifolia* var. *variifolia* (A.Meeuse) Holstein. Sessile leaves are common in *C. quinqueloba* and *C. senensis*, while the other species' leaves are usually distinctly petiolate. The petioles' surface can be glabrous, at maturity speckled with hyaline to white cell clusters (*C. grandis*, *C. heterophylla*, *C. intermedia*, *C. quinqueloba*, *C. rehmannii*, *C. samburuensis*, *C. senensis*, *C. subsessiliflora*), or have an indumentum. The petiole contains several vascular bundles arranged in a U-shape (Fig. 5a). However,




Figure 4. **a** Young plant of *C. abyssinica* (N. Holstein 132). The first node is in the axilla where the glabrous cotyledons split off. The first nodes lack probracts and tendrils **b** Young plant of *C. sessilifolia* (N. Holstein 131). The first leaves in this species are petiolate, sessile leaves are produced later-on. The glabrous cotyledons are already dried (plant had the same age as the one in Fig. 4a).




Figure 5. **a** Cross-section of a petiole of *C. grandiflora*, stained with astra blue and safranin (3:2). The bicollateral vascular bundles are arranged in a U-shape **b** Trichome on the adaxial ridge of a cross-section of a petiole of *C. grandiflora*. Although not visible by naked-eye, the petiole is also covered with few-celled glandular trichomes **c** Young plant of *C. adoensis* var. *jeffreyana*. The trichomes are mainly occurring on the prominent veins. The adaxial side of the petiole bears smaller trichomes on the ridges, which fade into the leaf margin.

Hussain et al. (2011) report a ring of vascular bundles in *C. grandis*. The adaxial side of the petiole often bears two ridges above the “lateral” vascular bundles (Fig. 5a). These ridges merge into the leaf margin and usually bear trichomes (Figs 3b, 5b, c). The abaxial side of the petiole shares its indumentum with the lower leaf lamina, at least at the base of the veins (Fig. 5c).

The venation in *C. grandis* is reticulate, and the mid rib is reported to contain three bicollateral vascular bundles with xylem and phloem arranged in a ring (Hussain et al. 2011). Reticulate venation can be confirmed for all *Coccinia* species except *C. ogaden-sis*, in which only the central vein in each lobe is visible.




Figure 6. Leaf shape variability in *Coccinia senensis*. **a** H.J. Schlieben 5259 (HBG) **b** E.M.C. Groenendijk *et al.* 1031 (WAG) **c** A.R. Torre *et al.* 18788 (MO). Black bars equal 1 cm.

Young leaf buds often bear a dense indumentum, even in species that are glabrous at maturity, e.g., in *C. grandis*. The leaf lobes are linear, elliptic, (ob-)ovate to triangular. The incision depth of the lobes can be consistent (*C. ogadensis*, *C. subsessiliflora*) or highly variable (*C. adoensis*, *C. grandis*, *C. senensis* (Fig. 6)), even within a single individual. In taxa with a variable degree of lobation, young leaves tend to be not or shallowly lobed (e.g., *C. grandis*, *C. megarrhiza*, *C. rehmannii* aff. var. *littoralis*, *C. sessilifolia*), a differentiation according to light exposure might also be possible. The leaf margin usually is beset conspicuously with small teeth and often bears trichomes (Figs 3b, 5b, c), even in otherwise glabrous species (e.g., *C. grandis*, *C. sessilifolia*). The teeth are at the apex of lobes, lobules and smaller orders of serration or situated along the entire margin. The term “dentate” (toothed) is therefore ambiguous in literature describing *Coccinia*, as it might also refer to the margin morphology (Stearn 2004). The teeth are often pale, but can also be colored, esp. when dry, such as in *C. abyssinica*, *C. grandis*, *C. intermedia*, *C. longicarpa*, *C. megarrhiza*, and *C. samburuensis* (Fig. 7a). The coloration of teeth is inconspicuous in young plants and develops during maturation (as observed in *C. abyssinica*, *C. grandis*, and *C. megarrhiza*). The teeth are interpreted as hydathodes by Zimmermann (1922a), because he observed water drops in *C. grandis* and *C. trilobata* on the teeth of the 2nd order (except those of the tip of



Figure 7. a Male flower of *C. grandis* (N. Holstein 37). Apices of the petals and calyx lobes, as well as major teeth on the leaf margin are colored in red. Minor margin teeth are inconspicuously colored. The calyx lobes in *C. grandis* are spreading in flower buds and reflexed in mature flowers **b** Male plant of *C. sessilifolia*. Darkish glands (extranuptial nectaries) are commonly found at the base of a lower leaf lamina. The calyx lobes are unusually large in this specimen (cp. Fig. 2b).

the lobes) in the morning. A white deposit at the teeth on the upper side of the leaf of a *C. adoensis* plant (P. Quarre 75; PR) seems to support Zimmermann's interpretation.

The upper leaf lamina is often covered with transparent to white pustules, that contain cystoliths (Avetta 1894; Solereder 1899; Zimmermann 1922a; pers. observ.). The pustules consist of up to 25 cells in *C. mackenii* (Avetta 1894) but are larger and denser in glabrous species from dry habitats (esp. *C. ogadensis*). As they develop over time (they are smaller and less well visible in forest species), it can be assumed that the pustules are an adaptation towards protection against high solar radiation. When acetic acid is applied to microscopic sections of the leaves, heavy gas development suggests that the cystoliths consist of CaCO_3 (pers. observ.). This can be observed in *C. grandis*, *C. hirtella*, and *C. sessilifolia*, hence also when the pustules are not conspicuous as in the latter two species. The pustules may form the base of small trichomes, such as in *C. adoensis* var. *jeffreyana* (Fig. 3b) or *C. microphylla*. In some species, the upper surface is usually covered with an indumentum (*C. hirtella*, *C. schliebenii*, *C. senensis*, and *C. trilobata*), but it may also be reduced, and other species rarely exhibit a trichome-bearing upper surface, e.g., *C. adoensis*. In each case, the trichomes are simple, < 1 mm, and whitish. The veins on the upper surface are either glabrous to the naked eye or are covered with small < 0.5 mm long simple trichomes. Zimmermann (1922a) observed in *C. grandis* that the glabrous surface of the lamina is only slightly wettable, whereas a drop of eosine disperses along the veins rapidly. Zimmermann argues that these "capillary drainage lines" might serve to transport water to the hydathodes during the dry season.

The lower leaf lamina is paler than the upper side (Fig. 3b) and can be glabrous or bear an indumentum. The highest density of the indumentum can be found on the

prominent veins (Fig. 5c). The indumentum on the lower leaf surface and the abaxial surface of the petiole can consist of eglandular oligo- to multicellular trichomes. The trichomes are appressed or upright (Fig. 5c), usually filiform, sometimes also narrowly conical (e.g., *C. abyssinica*). Filiform trichomes are straight, curved, or sinuate. Long filiform trichomes often appear articulate when dry due to sunken lateral cell walls (Fig. 3a). Dry trichomes are hyaline, whitish, beige, or yellowish. The lower lamina often displays deeply colored to dark green to blackish extranuptial glands (Fig. 7b). The glands usually occur at the base of the leaf between the veins, sometimes also between secondary ramifications (*C. grandis*) or along the main veins (*C. grandiflora*). The epidermis of the lower leaf lamina in *C. grandis* consists of cells with undulating anticlinal cell walls and anomocytic stomata (pers. observ.).

Probracts and bracts

In addition to the foliose leaves, most *Coccinia* species have bracteose prophylls on sterile nodes, which are called “probracts” (Zimmermann 1922b). The probracts can be up to 5 mm long, but also rather small (< 1 mm) or caducous. The first nodes of the seedling lack probracts, and they are developed on later nodes. The shape of the probracts is ovate and entire with a round to acute apex. They are often spoon-like presenting the lower surface (e.g., *C. adoensis*, *C. barteri*, *C. grandis*, *C. megarrhiza*, *C. sessilifolia*; Figs 3a, 8b), or they are folded in the middle with a prominent keel (*C. grandiflora*; Fig. 8a). Probracts can be glabrous or bear short (< 1 mm) trichomes, and bear extranuptial glands on the lower surface (Fig. 8a; Okoli and Onofeghara 1984).

Bracts (leaves subtending inflorescences or flowers), if present, look like the probracts. Bracts below inflorescences are as large as probracts, while bracts below flowers tend to be smaller. Bracts can be present or absent, the latter being an indicative character for some species.

In rare cases, probracts and bracts can be leaf-like (e.g., *N. Holstein* 126, Fig. 2b; *P.C.M. Jansen* 2065; *H. Wanntorp* & *H.E. Wanntorp* 1159) indicating that the (pro-) bracts are likely derived from usual leaves.

Extranuptial glands

The conspicuous glands on the lower leaf surface, probracts, and bracts (Figs 7b, 8a) are of the *Benincasa*-type (sensu Zimmermann 1932), meaning that they are flat and consist of several layers of secretory cells, which are surrounded by a single-layered sheath (Muhammad Ilyas 1992; Okoli and Onofeghara 1984; pers. observ.). This sheath is lignified in *C. microphylla* and *C. trilobata* (Zimmermann 1922b). However, Zimmermann (1932) cites Nieuwenhuis von Üxküll-Güldenbandt as saying that the sheath in *C. grandis* is suberitized, but the present author did not find such a statement in the citations given in that paper (eventually, this was a personal communication).




Figure 8. **a** The probract of *C. grandiflora* is fleshy and has a keel. The abaxial side bears many extranuptial nectaries. The structure pointing to the lower left of the picture is the tendril **b** The probract of *C. trilobata* (sampled as *N. Holstein* & *P. Sebastian* ♀) is spoon-shaped, papery, and without a keel.

Schrödter (1926), however, finds that young sheaths in *Luffa aegyptiaca* are lignified but become suberized with age, so the difference might be explained by different stages. Chakravarty (1948) interprets the sheath as filter tissue that is surrounded by an “external osmotic tissue”. Also Muhammad Ilyas (1992) interprets these radially elongated cells as secretory and notes that they have a connection to the vascular strand. However, Okoli and Onofeghara (1984) find that the glands in *C. barteri* are too distant to be interpreted as vascularized. Zimmermann (1922b) observes intermediate forms between few-celled, stalked glandular hairs and the *Benincasa*-type glands in *C. microphylla* and *C. trilobata*, including the sheath that forms the base of the protruding glandular tissue. The glands secrete a clear, rarely slightly colored, sweet-tasting exudate (pers. observ. in *C. grandiflora* and *C. grandis*). In *C. grandis* the exudates contain sucrose, glucose, fructose, alanine, tryptophane, threonine, and an unidentified amino acid (Muhammad Ilyas 1992).

Peduncles and pedicels

Male flowers mostly occur in racemes that are usually accompanied by 1–2 solitary flowers on the same node (Fig. 2b). The first flowers in male plants of *C. hirtella*, *C. rehmannii*, and *C. sessilifolia* are solitary. Racemes appear later in the course of the flowering season, although racemes are generally rare in the first species (pers. observ.). If solitary flowers and racemes are produced on the same node, then the solitary flower(s) precede(s) those of the racemes in time of maturity (Fig. 2b). The trigger to produce racemes instead of or additionally to solitary flowers is not known. The racemes bear up to 35 flowers (e.g., in *C. pwaniensis*, *C. racemiflora*). Within the racemes, flowering starts at the basalmost branches. If the peduncle is reduced, flowers

appear clustered on the node. In *C. grandis*, which usually produces single flowers only, flower clusters (short-peduncled racemes) rarely occur. This can be seen in plants from Ethiopia, Saudi Arabia but also from India and Sri Lanka. The pedicels of solitary male flowers of *C. hirtella*, *C. megarrhiza*, and *C. rehmannii* exhibit a negative gravitropism. In creeping plants, pedicels that grow downwards in the beginning make a sharp bent upwards to present the flower upright.

Female flowers are mostly solitary. Only in some species, female flowers are usually in racemes, such as in *C. heterophylla*, *C. keayana*, and *C. racemiflora*. Few-flowered female racemes or clustered flowers might also occur in *C. grandiflora*, *C. intermedia*, and *C. subsessiliflora*. In *C. barteri*, female flowers can be solitary or in few- or many-flowered racemes. Two female flowers per node have also been observed in *C. microphylla*. The pedicels of solitary female flowers are negatively gravitropic during flower development. After pollination, the pedicels of solitary female flowers of *C. grandis*, *C. hirtella*, *C. megarrhiza*, *C. microphylla*, *C. rehmannii*, and *C. sessilifolia* exhibit positive gravitropism. The downturn is not due to slackness caused by the weight of the developing fruit but an active process, as the pedicels thicken and remain firm. However, only fertilized flowers turn downwards completely, as aborted flowers from mis-pollination never reach this state (pers. observ. in cultivated plants).

Perianth

The perianth of all *Coccinia* species is synsepalous and sympetalous. At the base, calyx tube and corolla tube are connected with each other and form a perianth tube or funnel. Depending on the exertion point of the staminodes in female flowers, parts of the tube form a hypanthium (e.g., *C. grandiflora*).

The calyx differentiates as a bulge (Fig. 7a, b) with usually five lobes, or only the lobes emerge from the perianth tube (Fig. 9). If the calyx emerges as a bulge, then it and the perianth tube are rather conspicuously differentiated from the corolla in terms of color. If only the calyx lobes emerge, then the color of perianth tube fades to green color towards the receptacle, with the veins of the corolla remaining more intensely colored. Whether calyx and corolla are non-differentiated (congenital fusion) or postgenitally fused, is not known for *Coccinia*, but in the distantly related *Echinopepon wrightii* (A.Gray) S.Watson the perianth tube is non-differentiated (Leins and Galle 1971). The outside of the perianth tube can bear long trichomes of the type on as the lower leaf surface or the petioles (Fig. 10a). The calyx lobes are acute triangular to subulate or linear, rarely slightly lanceolate. The orientation of the calyx lobes is erect, spreading, or reflexed, although they can be curved inwards (e.g., *C. rehmannii* aff. var. *littoralis* (Fig. 10a) or outwards (e.g., *C. intermedia*). The color of the calyx lobes can be more intense (green) than the perianth tube or the pedicel (Fig. 7b). In *C. grandis*, the tip of the calyx lobes is brownish to reddish just as the teeth on the leaves and the corolla (Fig. 7a).

The petals of *Coccinia* species are fused at the base, usually for at least one third of the total length. Rarely, the petals are free down to the height of the calyx lobes (pers.



Figure 9. Leaky dioecy in a plant of *C. megarrhiza*. The plant was flowering male (the bud on the left) through the season, but a single female flower developed (the second flower on this node was male). The flower was receptive and produced a normal-sized fruit and normal-shaped seeds. The probracts (left node) are spoon-shaped, the tendrils are purplish.




Figure 10. **a** Male plant of *C. rehmannii* [aff. var. *littoralis*] (N. Holstein 126). The plant was raised from seeds of a female plant with ovoid fruits (B. Jarret – pers. comm.) **b** Male flower of *C. adoensis* var. *aurantiaca* (N. Holstein et al. 85). The halictid bee (H. Schaefer – pers. comm.) circled around the globose anther head harvesting pollen. The scent of the flower was strong and honey melon-like.

observ. in *C. megarrhiza*, *C. rehmannii* var. *rehmannii*, and *C. sessilifolia*). Perianth tube and corolla tube are often campanulate, rarely funnel-shaped or tubular. The perianth tube can be urceolate in *C. longicarpa*, *C. racemiflora* and sometimes in *C. barteri*. The tips of the (4–)5(–7) corolla lobes are rounded to acute with an apical tooth. The



Figure 11. **a** Longitudinal section of a female *C. hirtella* flower (size of the small squares is 1 mm²). Perianth and style were detached. The pollen sacs on the staminodes are highly reduced. The introrse side of the staminodes bears long trichomes that in intact flowers touched the style **b** Female flower of *C. rehmannii* var. *rehmannii* with bilobate stigma arms **c** Female flower of *C. megarrhiza* with bulging stigma arms.

apical tooth can be inconspicuous or colored claret red or brown, such as in *C. adoensis* var. *aurantica* or *C. grandis*. Outside, the perianth tube and the corolla is glabrous or covered with short (< 10 globose cells in *C. grandis*) trichomes. Inside, the corolla is covered with long trichomes (up to 20 cells in *C. grandis*), sometimes with a glandular apical cell (Fig. 10b). The trichomes become shorter towards the receptacle. The inner side of the hypanthium of female flowers is glabrous and smooth, which suggests nectary tissue, in *C. grandiflora*, *C. grandis*, and *C. hirtella* (Fig. 11a). The size of the corolla does not differ conspicuously between staminate and pistillate flowers; pistillate flowers might be a bit smaller.

Androecium

In staminate flowers, the three stamens originate from the base of the perianth tube, and the filaments are fused to a central column (Fig. 10a). The fusion point sometimes leaves a small gap to the hollow receptacle center. This gap, however, can be filled with long multi-cellular trichomes (e.g., in *C. abyssinica* and *C. megarrhiza*). Rarely, the filaments can also be separate. The filaments are glabrous and white, greenish, yellowish, or orange. The number of vascular bundles in the stamens is disputed. Chakravarty (1954) reports five vascular bundles in *C. grandis*: two stamens have two bundles each, and the third stamen has a single bundle. Later research shows three vascular bundles for the same species with one per stamen (Bhuskute et al. 1986; Deshpande et al. 1986). In *C. hirtella* each stamen contains a single vascular bundle (pers. observ.).

The anthers together form a globose head (Fig. 10b). Each anther is bithecate; sometimes one can be monothecate (Bhuskute et al. 1986; Chakravarty 1954). Each theca is sinuate. Deshpande et al. (1986) report a bi-layered fibrous endothecium and a secretory tapetum, which they found differs from distantly related *Momordica charantia* L.

In pistillate flowers, the three, now free, stamens are reduced to staminodes that originate from the interior perianth wall, forming a hypanthium. Introrsely, the staminodes of *C. grandiflora*, *C. grandis* and *C. hirtella* bear long, multicellular trichomes, except for the apex, extrorsely the staminodes are glabrous (Fig. 11a). The anthers of the staminodes are strongly reduced to a slightly yellowish spot at the apex in *C. grandis* and *C. hirtella*. The staminodes of *C. megarrhiza* bear long multicellular trichomes introrsely and laterally but are glabrous extrorsely.

Pollen

Pollen in *Coccinia* species shows little diversity. The pollen is oblate-spheroidal to prolate with a reticulate exine (Table 1). Additionally, the pollen of *C. pwaniensis* (Holstein and Renner 2010), *C. hirtella*, and *C. trilobata* is prolate, the exine texture is unknown. The sampling of the examined species covers all clades and suggests uniformity in shape and exine texture, which negates systematic value of pollen in *Coccinia*. The color is yellow in *C. abyssinica*, *C. grandiflora* (Zimmermann 1922b; and pers. observ.), *C. grandis*, *C. hirtella*, *C. megarrhiza*, *C. microphylla*, *C. rehmannii*, *C. sessilifolia*, and *C. trilobata*, and orange in *C. adoensis* var. *aurantiaca*. Zimmermann (1922b) reports that pollenkitt of *C. grandiflora* contains a yellow colorant that is soluble in peanut oil, but not in water and only slightly in heated chloral hydrate solution. It changes its color in concentrated sulfuric acid to blue, in Lugol’s iodine (I₂KI) to green, and in osmic acid to brown. As in several other cucurbit species, *in vitro* germination of *C. grandis* pollen increases from pH = 7 towards alkalinity and is maximal at pH = 8.5 (Zaman 2009).

Table 1. Pollen characters in *Coccinia* species.

| | shape | Size (P × E) [μm] | Exine texture | source |
|-------------------------------------------------|---------------------------------|-----------------------------|---------------------|-------------------------|
| <i>C. abyssinica</i> | Prolate-spheroidal | 60–70 × 56–65 | Reticulate | Martcorena 1963 |
| | Oblate-spheroidal to spheroidal | 76 × 81 | Reticulate | Khunwasi 1998 |
| <i>C. adoensis</i> | Prolate | 66–73 × 45–50 | Reticulate | Martcorena 1963 |
| | Prolate-spheroidal | 72 × 61 | Reticulate | Khunwasi 1998 |
| <i>C. barteri</i> | Prolate | 70–80 × 50–60 | Reticulate | Martcorena 1963 |
| | Prolate-spheroidal | 71 × 58 | Reticulate | Khunwasi 1998 |
| <i>C. grandiflora</i> | Prolate | | | Zimmermann 1922b |
| <i>C. grandis</i> | Prolate | 60–63 × 34–40 | Reticulate | Martcorena 1963 |
| | Prolate-spheroidal | 58 × 52 | Reticulate | Khunwasi 1998 |
| | Prolate | 47.61–64.62 × 35.91–44.80 | Coarsely reticulate | Perveen and Qaiser 2008 |
| | Subprolate to prolate | 34–52 × 28–35 | Reticulate | Awasthi 1962 |
| | Prolate | 41.20 ± 0.61 × 34.00 ± 0.45 | Reticulate | Datta 1988 |
| <i>C. megarrhiza</i> | Oblate-spheroidal to spheroidal | 92 × 92 | Reticulate | Khunwasi 1998 |
| <i>C. mildbraedii</i> | Prolate | 55–60 × 35–41 | Reticulate | Martcorena 1963 |
| <i>C. sessilifolia</i> var. <i>sessilifolia</i> | Prolate-spheroidal to prolate | 70 × 58 | Reticulate | Khunwasi 1998 |

Gynoecium

Pistillate flowers are epigynous and have three (rarely two or four) carpels. The ovary is narrowly spindle-shaped, oblong to globose. The surface is smooth or warty; it is glabrous or has the indumentum of the pedicel. The style is often greenish-white or pale-yellowish; the stigmas are frequently in yellowish colors and covered with long trichomes (Fig. 11c). Each stigma in *C. grandiflora*, *C. grandis*, *C. hirtella*, *C. megarrhiza*, *C. rehmannii*, and *C. sessilifolia* is U-shaped with the ends of the lower sides of the arms touching each other. The stigmatic branches can be long and free, such as in *C. grandiflora*, *C. grandis* and *C. rehmannii* var. *rehmannii* (Fig. 11b), or short and bulbous, such as in *C. hirtella*, *C. megarrhiza* (Fig. 11c), *C. microphylla*, and *C. sessilifolia*.

The placentation of the ovules in *Coccinia* is involute, which is also discussed for other Cucurbitaceae by Leins and Galle (1971). The funicle appears to be attached to the outer wall, but actually attaches to a septum coming from the axis (Fig. 11a, b, c), which itself is connected to the outer wall. The axis-wall septum, however, might be reduced during ripening, but this needs further study.

The anatomy of the ovules is barely surveyed in *Coccinia*. In *C. abyssinica* and *C. grandis*, the ovules are reported to be anatropous, bitegmic, and crassinucellate (Getahun 1973; Zahur 1962). Anatropous ovules also occur in *C. hirtella* (Fig. 12a). The position of the ovules is horizontal in *C. grandis* (Kirkwood 1904), *C. hirtella* (Fig. 12a), and *C. megarrhiza* (Fig. 12b) with the micropyle facing outwards. Horizontal ovules are regularly reported in the Benincaseae and the Cucurbitaceae (Schaefer and Renner 2011a).

In staminate flowers, a pistil is not developed because the stamens fuse to a central column. The pedicel is narrow and reaches the perianth, and there is no indication of even a thin (sterile) inferior ovary in the flower.

Female gametophyte development and embryology

The development of the embryo has only been investigated in *C. grandis*. Chakravorti (1947) and Zahur (1962) report that the female gametophyte development is according to the *Polygonum*-type. Both observe that the synergides possess hooks instead of the filiform apparatus. Chakravorti (1947) describes the developing endosperm as a nuclear type, which is confirmed by Chopra (1955). By formation of a large central vacuole, the nuclei become displaced to the periphery. After the endosperm becomes cellular, the often lateral chalazal haustorium remains coenocytic. Then, the haustorium becomes cellular with multinuclear cells except for the apex (Chopra 1955).

Fruits

The fruits are many-seeded berries, which vary in size and shape between species (and within *C. grandis*, *C. rehmannii*, and *C. subsessiliflora*). The smallest fruits occur in



Figure 12. **a** Cross-section through an ovary of *C. hirtella*. The ovules are anatropous with the micropyle facing outwards **b** Cross- and longitudinal section of a *C. megarrhiza* fruit. The seeds are enclosed in a hyaline hull (aril) and seemingly attached to the periphery **c** Cross-section through a fruit of *C. sessilifolia*. Note that the vascular bundles in the lower left of the picture bend in the periphery, so the placentation is not parietal but involute.




Figure 13. **a** Ripening fruit of *C. hirtella*. Note the typical lobulate leaves of this species in the lower right **b** Ripening fruit of *C. sessilifolia*. The fruit, like the plant, bears a waxy bloom **c** Ripening fruits of *C. megarrhiza* have a dark green halo around the white longitudinal mottling. The left fruit is derived from pollination with *C. megarrhiza* pollen, whereas the smaller fruit on the right is derived from cross-pollination with *C. trilobata* (both pollinations were conducted on the same day).

C. rehmannii var. *rehmannii* and *C. microphylla* with globose berries as small as 1 cm in diameter at maturity. However, in both species larger globose fruits (up to 2.5 cm in diam.) and in *C. rehmannii* ovoid fruits may occur additionally, in the latter case especially in more humid habitats. The largest fruits occur in *C. samburuensis* and the rainforest species *C. grandiflora*, *C. longicarpa*, *C. mildbraedii*, and *C. schliebenii*, which have long elliptical to cylindrical (sausage-shaped) fruits up to 20 cm long and up to 5 cm in diameter.

Immature fruits often have a white or pale-green (*C. hirtella*, *C. sessilifolia*) or dark green (*C. adoensis*) longitudinal mottling or lines, even when the ovary and the ripe fruit is single-colored (Figs 2a, 13a, b, c). In the *C. rehmannii* clade, the white spots or lines become surrounded by a dark green halo during ripening (Fig. 13c). Rarely, if no white mottling develops, e.g., in some *C. microphylla*, dark green spots develop nevertheless. In any case, the mature fruit in species of the *C. rehmannii* clade is usually uniformly colored red (Zimmermann 1922b; and pers. observ.). Ripening usually oc-

curs from green with or without mottling via yellow to orange to the final coloration. The color changes from the apex of the fruit downwards (Fig. 13a, b, c), independent of the position (hanging vs. horizontal) in *C. sessilifolia*. In *C. megarrhiza*, pendulous fruits ripen from the apex to the base, which sometimes remains green even when the apex already turns soft. In lying fruits from creeping *C. megarrhiza* plants, ripening does not proceed from the apex, but starts from point that is closest to the source of either warmth or light (pers. observ. from greenhouse cultivation). The degree of the yellow to orange ripening zone varies. In *C. sessilifolia*, fruits directly turn red, whereas in *C. grandis* the color change includes a well visible yellow zone. Unripe fruits collected of *C. grandis* tend to turn yellow outside and pink to red inside (Imbumi 2004). Mature fruits are in deep red colors (hence the genus name) or orange-red. Rarely, a white longitudinal mottling is described in ripe fruits (e.g., *C. mackenii*).

Immature fruits are glabrous or have the same indumentum as the ovary. By ripening, the indumentum is usually reduced. The exocarp of *Coccinia* fruits is papery thin and has a waxy bloom when ripe. The endo- and mesocarp are red, fleshy and soft (Fig. 12b, c). The pulp is nerved with a dense network of tubular tissue. Shah et al. (1983) report that such a network consists of sieve tubes in *C. grandis*. The sieve tubes are not connected to the main vascular strands and are filled with a proteinaceous material. The authors suggest that the sieve tube network aids nutrient transport during the rapid growth of the fruit.

Seeds

The seeds (Fig. 14) in *Coccinia* species are beige to grayish with a small margin, which often has a darker coloration. The shape is more (esp. in the *C. rehmannii* clade) or less asymmetrically ovate (especially *C. adoensis*, *C. pwaniensis*, and *C. senensis*). The surface is flat, esp. in the *C. rehmannii* clade, to lenticular (esp. *C. adoensis*, *C. pwaniensis*, and *C. senensis*). The size varies from $4.5\text{--}7 \times 3\text{--}5 \times 1\text{--}2$ mm (L/W/H). Seed numbers per fruit vary drastically from about 10 (*C. microphylla*) to c. 100 (+ c. 20 infertile) in *C. sessilifolia* (N. Holstein 119). Species with larger fruits might contain more seeds.

Detailed observations of the seed anatomy have been made by Getahun (1973) for *C. abyssinica* and by Chakravorti (1947) for *C. grandis*. Getahun describes the mature seed as consisting of the embryo, a membrane-like structure (pellicle) closely adhering the embryo but separated from the hard testa. However, Chakravorti does not recognize a pellicle in *C. grandis*. Both authors agree that the inner integument disappears and the testa develops solely from the outer integument. The testa of *C. abyssinica* is described as comprising four layers (from center outwards): (1) a thin-walled parenchyma, (2) a sclerenchyma of macrosclereids, (3) a thick-walled parenchyma, and (4) an epidermal layer. The outermost layer, the epidermis, is disintegrated, leaving the cell walls as slender rods of 500 μm length. This has also been noticed in other species, and the surface has been described as a fibrillose testa (Jeffrey 1967; Kéraudren 1967). De Wilde et al. (2011) also interpret the seed surface of *C. grandis* as having a disintegrated, pulpy, radiately striate exotesta.




Figure 14. Seeds of *Coccinia*. The lack of fibers in e and f are preparation artifacts due to mechanical removal of the hyaline aril. Maceration (coarse crushing of the fruit and soaking of the mass in water for 2–3 weeks; R. Brüggemann – pers. comm.) retains the surface fibers. Length of white bars equals 1 mm. **a** Seeds of *C. adoensis* var. *jeffreyana* (plant derived from seed of the same fruit: *N. Holstein* 130). Note the lenticular face and symmetrical shape of the seed **b** seeds of *C. abyssinica* (plant derived from seed of the same fruit: *N. Holstein* 120 and 132) **c** Seeds of *C. trilobata* **d** Seeds of *C. sessilifolia* (harvested by maceration) **e** Seeds of *C. sessilifolia* (harvested by mechanical extraction; taken from *N. Holstein* 119) **f** Seeds of *C. grandis* (harvested by mechanical extraction).

Getahun contrasts his observations with those of Chakravorti (1947) in *C. grandis*, but the seeds are in fact similar, just incompletely described by Chakravorti. Chakravorti draws a four-layered testa, but does not name the innermost layer that has the same hatching as the third layer, which he describes as “cells with thickened walls”. This is what Getahun calls parenchyma. Chakravorti’s outermost layer, the epidermis, consists of radially elongated cells with thin walls. These cells have likely just not yet disintegrated as observed by Getahun. The only difference between both observations is the second layer, which consists of macrosclereids in *C. abyssinica* and of radially slightly elongated cells with thin walls in *C. grandis*. These different observations are explainable by two possibilities: 1) different developmental stages of the seeds, since Chakravorti surveys the seed development, so the layers are immature, while Getahun surveys mature seeds and germination, or 2) different staining. Chakravorti uses haematoxylin alone, which does not stain lignified cell walls, whereas Getahun uses haematoxylin with safranin as counter stain, which makes lignin, and thus sclerenchyma, well visible (von Aufseß 1973).

The seeds in *Coccinia*, at least in *C. abyssinica* (Hora 1995), *C. grandis*, *C. hirtella*, *C. megarrhiza*, *C. microphylla*, *C. mildbraedii*, *C. sessilifolia*, and *C. subsessiliflora* (pers.

observ.) are surrounded by a hyaline red juicy envelope (Fig. 12b). As the ovule is bitegmic (see above), one might assume the hyaline envelope is the testa. Chakravorti (1947), however, observes that the juicy envelope is derived from carpellary tissue. However, Getahun (1973) does not recognize the hyaline hull in *C. abyssinica*, which is surprising as it also occurs in the closely related species *C. megarrhiza* and *C. microphylla*. Similar structures to the hyaline hull are also found in other Cucurbitaceae, esp. in *Momordica*. Van der Pijl (1982) interprets these as “endocarp-pulpa” taking over the function of an aril for seed dispersal as the fruits of *Momordica* species dehisce at maturity. However, *Coccinia* fruits disintegrate and do not dehisce, e.g., into valves.

Germination

The seeds of *C. abyssinica* maintain a high germination rate (100%) after four years of storage at room temperature (Getahun 1973). However, time from watering until germinating increases from 4 days (after one year of storage) to 16 days (after four years of storage). Seeds of *C. grandis* are also able to germinate after four years of storage, while seeds of *C. ogadensis* Thulin (3 seeds tested) did not germinate after five years (pers. observ.). Getahun (1973) reports that *C. abyssinica* seeds do not germinate below 10 °C and above 35 °C. In the latter case, he observes thermal damage to hypocotyls and primary roots. The optimum for germination in *C. abyssinica* is between 20 and 30 °C and that of *C. grandis* is 35 °C, whereas temperatures < 23.5 °C and > 40 °C inhibit germination (Li et al. 2001). The germination rate of *C. abyssinica* seeds in the light is decreased by 35% compared to germination in darkness (Getahun 1973). In *C. sessilifolia*, seed viability declines after 9 months, and germination is at a maximum after 10–20 min smoke exposure or red:far red light treatment, followed by burying and a long-day cycle (Weiersbye and Witkowski 2003). Rotting of a crushed ripe fruit in water (for seed extraction) resulted in germination of two seeds in an artificial hybrid (*C. megarrhiza* ♀ × *C. rehmannii* aff. var. *littoralis* ♂) after 3 weeks of soaking (pers. observ.). *Coccinia grandis* seeds do not exhibit dormancy (Motooka et al. 2003); for the other species there is no information available.

Genome, chromosomes, and hybridization

Chromosomes and sex determination

Coccinia is one of the few examples in the plant kingdom, in which at least one species has heteromorphic sex chromosomes (Ming et al. 2011). *Coccinia grandis* contains 22 autosomes plus 2 gonosomes. Female individuals have homomorphic XX, whereas male individuals have heteromorphic XY chromosomes (Fig. 15a). Although Kumar and Deodikar (1940) report males to have two large “X” and females a large X and a smaller Y chromosome, later studies (Chakravorti 1948; Kumar and Vishveshwaraiah 1952) reveal that males are heteromorphic and the Y is 2.5 (Bhaduri and Bose 1947)




Figure 15. Chromosome preparations fixed in 3:1 EtOH – acetic acid and stained with orcein, objective lens: 100×. **a** Mitotic plate of a male *C. grandis* ($2n = 22 + XY$) **b** Mitotic plate of a male plant of *C. hirtella* ($2n = 24$) **c** Phase contrast image of a mitotic plate of a male plant of *C. sessilifolia* ($2n = 24$).

Table 2. Chromosome numbers of *Coccinia* species and sexes of surveyed individuals.

| Species | Sex of individual | Chromosomes ($2n$) | Voucher |
|----------------------------------------------------|-------------------|----------------------------|-----------------------------------------------------------|
| <i>C. grandiflora</i> | female | 24 | <i>N. Holstein</i> 114 (EA, M) |
| <i>C. grandis</i> | male | 22 + XY (Fig. 15a) | <i>N. Holstein</i> 32 (M) |
| <i>C. grandis</i> | female | 22 + XX | <i>N. Holstein</i> 25 (M) |
| <i>C. hirtella</i> | male | 24 (Fig. 15b) | <i>N. Holstein</i> 29 (M) |
| <i>C. rehmannii</i> aff. var. <i>littoralis</i> | male | 20 | <i>N. Holstein</i> 126 (M) |
| <i>C. sessilifolia</i> | male | 24 (Fig. 15c) | <i>N. Holstein</i> 13 (M), <i>N. Holstein</i> 109 (M (3)) |
| <i>C. sessilifolia</i> | female | 24 | <i>N. Holstein</i> 119 (B, M) |
| <i>C. trilobata</i> | male | 20 (A. Sousa, pers. comm.) | <i>N. Holstein</i> & <i>P. Sebastian</i> 9 (M) |

to 3–4 times longer (Guha et al. 2004) than the other chromosomes. Some years before Kumar and Deodikar, Sutaria (1936) reported $n = 12$ from pollen mother cells of *C. grandis*, without finding the large Y chromosome. Although scientists from India conducted some research for *C. grandis*, chromosome work in other *Coccinia* species is almost none-existent. McKay (1930) reports $n = 12$ for *C. hirtella*, without mentioning whether he studied a male or a female individual. The author's own chromosome counts (Table 2; Fig. 15b) support McKay's report and reject the existence of heteromorphic sex chromosome in *C. hirtella* males. This is also the case for *C. sessilifolia* (Fig. 15c). Two counts in the *C. rehmannii* clade reveal a reduction of chromosome number and the non-existence of a heteromorphic Y chromosome there.

Due to the sex chromosomes, sex expression in *C. grandis* plants is pre-determined and sex ratios in the offspring basically follow Mendelian inheritance of a single allele. However, Agarwal and Roy (1983) report that of 500 planted seeds only 181 (36.2%) were male, and their interpretation is that there might be a genetic mechanism to reduce the number of male plants. As they do not report XY females, their finding might be explained rather by an increased lethality of XY embryos due to deleterious mutations on the single X or on the Y in Y-containing pollen. The Y chromosome in *C. grandis* is dominant, as the presence of a single Y results in male phenotypes, regardless of the number of X chromosomes (Agarwal and Roy 1975; Kumar and

Vishveshwaraiah 1952; Roy and Roy 1971b). Triploids of *C. grandis* with a $3n = XYY$ constitution are also male but bear flowers in clusters instead of the usually solitary ones and exhibit leaf deformations (Roy and Roy 1971b).

Evidence for Mendelian inheritance of sex in *C. hirtella* is not so clear, as the same plant can produce flowers of the opposite sex in succeeding seasons. Two plants marked as female and one as male from observation of flowers produced flowers of the opposite sex in the following year (pers. observ.), making *C. hirtella* functionally dioecious, but genetically hermaphroditic. On the other hand, there are several observations of flowers of the opposite sex in otherwise unisexual plants. Kumar and Vishveshwaraiah (1952) report a gynodioecious form of *C. grandis* that has homomorphic chromosomes (XX). Although bisexual flowers are reported to develop, pollen grain development is arrested, and the male function remains suppressed. Roy and Saran (1990), however, report fully fertile hermaphroditic flowers in an otherwise female individual. Holstein and Renner (2011a) report a collection of *C. intermedia* that bears male floral buds and female flowers and young fruits on the same nodes. This observation can be interpreted as 'leaky dioecy' (Baker and Cox 1984). Among the author's own cultivated plants, a male individual of *C. megarrhiza* produced a single female flower towards the end of the season (Fig. 9). Although two male flowers on the same individual were open at the same time when the female flower was mature, the pollen sacs did not open. It is not known whether this was a coincidence or functionally significant, e.g., to prevent selfing. Selfing is often discussed as being advantageous in small population sizes, e.g., when new islands are colonized. Prevention of simultaneous flowering of both sexes on the same plant implies that leaky dioecy would not immediately aid the establishment of new distant populations per se. It might require the establishment of several plants or clonal separation. In any case, the single female flower was receptive and was fertilized by another male *C. megarrhiza* plant derived from the same fruit as the "female" plant. The resulting fruit and seeds developed normally.

The production of hermaphroditic flowers in X-radiation studies (Agarwal and Roy 1983; Roy 1974) shows that dicliny in *Coccinia* is kept up actively. Agarwal and Roy found hermaphroditic flowers on two plants with otherwise female flowers and an XX configuration. They also report the development of a normal fruit without mentioning the fertility of pollen from hermaphroditic flowers, but interpret their finding as cleistogamy. However, fruit development without previous pollination (parthenogenesis) or from pseudogamy with pollen from different genera as fructification stimulus is described by Lal (1973). True selfing from own pollen (in irradiated XX individuals) would mean that the Y chromosome is not important for fertile pollen development. Furthermore, this means that it only carries at least one gene for suppression of the development of the female organs, and the occurrence of a second X suppresses pollen development in "normal" plants. Agarwal and Roy (1983) also report that X-ray dosages of 5 to 50 R [1.29×10^{-3} to 12.9×10^{-3} C/kg] result in a drastic reduction of the sex ratio (11 males, 2 hermaphrodites, and 127 females out of 500 irradiated seeds). This might indicate that the single X bears many functionally important genes in contrast to the Y, as mutations in the single X lead to an increased mortality compared to females with a balancing second X chromosome.

Genome of *C. grandis*

Aside from research on the sex chromosomes, a few studies on the genome of *C. grandis* have been undertaken. Guha et al. (2004) report the 4C nuclear DNA content of female *C. grandis* as 8.37 ± 0.14 pg, whereas that of male *C. grandis* is 10.17 ± 0.24 pg. This means that the difference between X and Y chromosome adds about 20% to the complete DNA content. Patankar et al. (1985) report a DNA content of $1C = 2.75$ pg for *C. grandis*, however, they do not report the sex of the analyzed individual. Interphase nuclear structure in *C. grandis* is chromocentric with 14 ± 0.25 chromocenters (Patankar et al. 1985).

Surveys on the reassociation kinetics in Cucurbitaceae suggest that *C. grandis* has the lowest amount of repetitive DNA among the six species studied in the Cucurbitaceae (Bhave et al. 1984). Fragments of 550 bp length have 25% of repetitive elements (Bhave et al. 1984), whereas 7400 bp long fragments consist of 49% repetitive DNA (Bhave et al. 1986). However, Pasha and Sen (1995) report different results as they find 400–600 bp long fragments to comprise 38% highly repetitive DNA (52% total repetitive DNA), which appears to be average in the Cucurbitaceae. Pasha and Sen do not discuss this difference, which cannot be explained by difference in sex, as the large amounts of plant material used (1 kg seeds or 100 seeds respectively) suggest that it must have comprised elements of both sexes. A sex discriminating study of *C. grandis* reassociation kinetics has not been undertaken yet.

Hybridization and crossing experiments

Charles Naudin's famous work on the effects of hybridization included crosses between *Coccinia* plants. He reports successful crosses between Asian *C. indica* (nom. illeg. for *C. grandis*) and the NE African *C. schimperi* (P06809214, P06809215, P06809216; Naudin 1862), which is now seen as a synonym of *C. grandis*. Naudin's *Coccinia schimperi*, however, has buff petals, whereas the Asian *C. grandis* has snow-white petals. Both supposed species hybridized without problems. During the following two years, Naudin could not intercross within the F1 generation because plants of different sexes did not flower at the same time, so he crossed F1 individuals with a female *C. grandis* from Asia, which again produced offspring. As Naudin erroneously supposed that he dealt with two species, he deduced that hybrids between species could be fully fertile, have a reduced fertility or be sterile, and that there was no clear boundary between species and varieties. However, he proved rather that the buff-petaled, African *C. schimperi* and the white-petaled Asian *C. grandis* are a single species obeying the biological species concept.

Naudin also crossed other *Coccinia* species that he had in cultivation. *Coccinia quinqueloba* and *C. mackenii*, although sometimes not easily distinguishable, were not amenable to crossing (Naudin 1866). Only 1 out of 20 crossing trials resulted in a fruit that developed poorly. Naudin did not publish whether the hybrid seeds were fertile or even viable, but his observations are valuable as each one accession of *C. quinque-*

loba and *C. mackenii* were not distinguishable using more than 3500 bp of plastid sequences, and hence might share the same plastid haplotype (Holstein and Renner 2011b). There are collections that share characters of *C. mackenii* and *C. quinqueloba*, but these are not intermediates. In these collections, long petioles (typical for *C. mackenii*) are coupled with simple tendrils (typical for *C. quinqueloba*), and thus cannot be unambiguously allocated to either species. However, if both typical forms are indeed reproductively isolated, then they are species sensu Mayr (1942), and the crossing behavior of these species needs to be tested reciprocally to define the morphological scope of the two species.

Naudin also crossed male *C. diversifolia* (*C. abyssinica*) with a female *C. mackenii*, which are rather distantly related and do not co-occur in nature. However, the cross resulted in onset of mediocre fruits with only few, but well-developed and viable seeds (Naudin 1866). Naudin did not report further results for this cross either.

As reproductive isolation between species is often assumed but rarely tested, crossing experiments among species that are cultivated in Munich Botanical Garden have been performed. Positive results are given in Table 3.

Interspecific fertilization succeeded or failed without correlation of relatedness or co-occurrence (Table 4). Female flowers of *C. sessilifolia* could not be fertilized with pollen of *C. megarrhiza* (4 trials), *C. trilobata* (4 trials), *C. rehmannii* (2 trials), *C. hirtella* (3 trials) or *Diplocyclos palmatus* (2 trials). Pollinated female flowers were discarded like non-fertilized flowers. Hence, hybridization seems to be prevented prezygotically in female *C. sessilifolia* with members of the *C. rehmannii* clade and *C. hirtella* as pollen donor. As *C. sessilifolia* and *C. rehmannii* co-occur widely in their range and share flowering time and floral syndrome, the production of hybrids would reduce fitness drastically. *Coccinia sessilifolia* and *C. hirtella* do not co-occur, but belong to the same clade (see chapter Evolution and phylogeny). Although a female *C. sessilifolia* could not be fertilized by pollen from *C. hirtella*, pollination of a female *C. hirtella* with pollen from *C. sessilifolia* resulted in fruit onset.

In contrast to *C. sessilifolia*, *C. grandis* is fertilized easily by *C. hirtella* and *C. sessilifolia*, although the species neither co-occur, nor are closely related. The cross resulted in offspring, which was growing vigorously but sterile, as pollen sacs did not open (Table 3). Hybrid pollen that was extracted from the pollen sacs of fully open flowers was also not able to fertilize *C. sessilifolia* (1 trial). The occurrence of sex chromosomes in *C. grandis* might result in gene dosage imbalance, which interferes with the floral development, leading to sterile offspring. The inability of female *C. grandis* to be fertilized by *C. rehmannii* aff. var. *littoralis* and *C. trilobata* (Table 4) might be explained by the fact that the chromosome numbers differ (see Table 2) and translocations lead to gene loss in hybrid genomes and thus inviability of the offspring. On the other hand, the cross between female *C. hirtella* and male *C. trilobata* (not sympatric) produced a purely intermediate F1 generation, which flowers vigorously despite the difference in chromosome numbers (see Table 2 and 3). Although the anthers open like in fertile flowers, unlike in *C. grandis* hybrids, the pollen of this hybrid was not able to fertilize female flowers of *C. hirtella* (1 trial), *C. grandis* (2 trials), or *C. sessilifolia* (2 trials).

Table 3. Description of the F1 from crosses and a natural hybrid between *Coccinia* species. Species used for the artificial crosses are not sympatric.

| Parent species | Offspring | Occurrence |
|------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|
| <i>C. grandis</i> ♀ × <i>C. hirtella</i> ♂ | F1 vegetatively morphologically intermediate; flowers are either aborted or sterile (pollen sacs remain closed); pollen globose; corolla is smaller than in each parent species (Fig. 16a) | Artificially in Munich Botanical Garden; voucher: <i>N. Holstein</i> 108 (M) |
| <i>C. hirtella</i> ♀ × <i>C. grandis</i> ♂ | F1 vegetatively morphologically intermediate; flowers smaller (Fig. 16b), sterile | Artificially in Munich Botanical Garden; voucher: <i>N. Holstein</i> 116 (M) |
| <i>C. grandis</i> ♀ × <i>C. pwaniensis</i> ♂ | F1 morphologically intermediate; flowers sterile (pollen sacs remain closed) (Fig. 16c) | Naturally in Pugu Hills, Dar es Salaam, Tanzania; vouchers: <i>N. Holstein et al.</i> 102 (DSM, M), 103 (M), 104 (M), 105 (DSM, M) |
| <i>C. grandis</i> ♀ × <i>C. sessilifolia</i> ♂ | F1 vegetatively morphologically intermediate; flowers are either aborted or sterile (pollen sacs remain closed); corolla is smaller than in each parent species (Fig. 16d) | Artificially in Munich Botanical Garden; voucher: <i>N. Holstein</i> 113 (B, M); <i>N. Holstein</i> 115 (M) |
| <i>C. hirtella</i> ♀ × <i>C. trilobata</i> ♂ | F1 morphologically intermediate; males flowering vigorously with intermediate flowers, pollen sacs open, but pollen is sterile | Artificially in Munich Botanical Garden; <i>N. Holstein</i> 121 (M) |

Table 4. Observations on fruit development of crosses between *Coccinia* species (except for crosses mentioned in Table 3). Viability and morphology of the F1 is not known so far. ^a = sympatrically distributed species, ^b = close relatives/sister species, ^c = species occur in the same area but in different habitats.

| Parent species | crossability |
|-------------------------------------------------------------------------------------------------|-----------------------------------------------|
| <i>C. grandis</i> ♀ × <i>C. abyssinica</i> ♂ | Onset of fruit (1 trial) ^c |
| <i>C. grandis</i> ♀ × <i>C. megarrhiza</i> ♂ | Onset of fruit (1 trial) ^a |
| <i>C. grandis</i> ♀ × <i>C. rehmannii</i> aff. var. <i>littoralis</i> ♂ | Abortion of flower (1 trial) |
| <i>C. grandis</i> ♀ × <i>C. trilobata</i> ♂ | Abortion of flower (2 trials) ^c |
| <i>C. hirtella</i> ♀ × <i>C. rehmannii</i> aff. var. <i>littoralis</i> ♂ | Onset of fruit (1 trial) ^c |
| <i>C. hirtella</i> ♀ × <i>C. sessilifolia</i> ♂ | Onset of fruit (1 trial) |
| <i>C. megarrhiza</i> ♀ × <i>C. abyssinica</i> ♂ | Onset of fruit (1 trial) ^{b & c} |
| <i>C. megarrhiza</i> ♀ × <i>C. hirtella</i> ♂ | Abortion of flower (1 trial) |
| <i>C. megarrhiza</i> ♀ × <i>C. rehmannii</i> aff. var. <i>littoralis</i> ♂ | Onset of fruit (1 trial) |
| <i>C. megarrhiza</i> ♀ × <i>C. sessilifolia</i> ♂ | Abortion of flower (2 trials) |
| <i>C. megarrhiza</i> ♀ × <i>C. trilobata</i> ♂ | Onset of fruit (Fig. 13c; 1 trial) |
| <i>C. microphylla</i> ♀ × <i>C. megarrhiza</i> ♂ | Abortion of flower (1 trial) ^a |
| <i>C. microphylla</i> ♀ × <i>C. trilobata</i> ♂ | Onset of fruit (1 trial) ^{b & c} |
| <i>C. rehmannii</i> var. <i>rehmannii</i> ♀ × <i>C. rehmannii</i> aff. var. <i>littoralis</i> ♂ | Onset of fruit (1 trial) ^b |
| <i>C. rehmannii</i> var. <i>rehmannii</i> ♀ × <i>C. trilobata</i> ♂ | Abortion of flower (2 trials) ^b |

Roy and Roy (1971a) report an intergeneric cross between a female *C. grandis* and a male individual of the monoecious *Diplocyclos palmatus*, resulting in a morphologically intermediate F1 offspring in 5% of the trials. All F1 individuals are female, indicating that the X chromosome bears at least one gene for maleness suppression, which is dominant over the maleness genes of *D. palmatus*. Whether the F1 is fertile, is not clear, as the authors report successful back-crossing only with a female [*sic*, male?]




Figure 16. Hybrids of *Coccinia* species. **a** *Coccinia grandis* ♀ × *C. hirtella* ♂ **b** *Coccinia hirtella* ♀ × *C. grandis* ♂ **c** *Coccinia grandis* ♀ × *C. pwaniensis* ♂; note that the pollen sacs are not open although the flower is in full bloom indicating sterility **d** Left flower: male *C. sessilifolia*, right flower: male flower of *C. grandis* ♀ × *C. sessilifolia* ♂ (*C. grandis* flowers are about equally sized as *C. sessilifolia*).

C. grandis but not with *D. palmatus*. To the author's knowledge, there are no reports of the F₂ generation. However, if the parental sexes were the other way around (male *C. grandis* × female *D. palmatus*), fertilization was not possible.

Plant – biotic environment interactions

Pollination

Although bee pollination is observed for only a few species, most *Coccinia* species exhibit characters that support a general attraction to bees. The petal color is commonly pale yellow but can also range to white, pale pink or bright orange, with green, yellow, orange to purple venation. Anthesis is during the day in *C. abyssinica*, *C. adoensis* var. *aurantiaca*, *C. grandiflora*, *C. grandis*, *C. hirtella*, *C. megarrhiza*, *C. microphylla*, *C. rehmannii*, *C. sessilifolia*, and *C. trilobata* (pers. observ.) but often only for a few hours (in, e.g., *C. megarrhiza*, *C. rehmannii*). Zimmermann (1922b) reports pollen release in *C. grandiflora* at 6.30 a.m. before blooming of the flower, opening of the flower

between 7 and 8 a.m. and wilting after noon. Ash (*J. W. Ash* 898) reports flower opening a.m. in *C. schliebenii*. Anthesis time in the other *Coccinia* species is not reported, but also likely to happen during the day. The scent is rather weak and dull sweetish, resembling that of honey melon, in *C. abyssinica*, *C. grandiflora*, *C. hirtella*, *C. rehmannii*, *C. sessilifolia*, and *C. trilobata*, weak but fresh in *C. grandis*, and intense, sweet, and fruit-like (like honey melon) in *C. adoensis* var. *aurantiaca* and *C. megarrhiza*. The only evidence of an alternative to bee pollination is *C. ogadensis*, which is reported to smell of rotten meat (*P. Ellis* 163 and 383). However, it is unclear whether the flowers emit a fetid scent or whether the smell comes from crushed vegetative parts, as it is known from *Momordica foetida* or *Kedrostis foetidissima* (Jacq.) Cogn. (Jeffrey 1967).

Bee pollination is confirmed for *C. rehmannii* (*C. J. Ward* 12250), *C. adoensis* var. *aurantiaca* (Fig. 10b), *C. grandiflora*, and *C. grandis*. Observed pollinators of *C. grandis* are *Trigona apicalis* Smith, 1857 and *Trigona collina* Smith, 1857 in Thailand (Jongjitvimon and Wattanachaiyingcharoen 2006), and *Megachile* sp. in Cambodia (H. Schaefer, pers. comm.). The present author observed a halictid bee (Fig. 10b; identification by H. Schaefer, pers. comm.) in a male *C. adoensis* var. *aurantiaca* walking on the globose anther head and collecting pollen in the corbicula. Stigmas in *Coccinia* are lobed (Fig. 11b) or bulging (Fig. 11c), and nectaries are located presumably in the hypanthium, so one can assume stripping of the pollen from the venter when crawling into the flower. Zimmermann (1922b) also observed the circling around the anther head in the large-flowered *C. grandiflora*. He identified the visiting small bee as *Trigona* sp. He also noted that a bee just having visited a *Momordica* flower walked on the inner side of the corolla loading dorsally located pollen on the anthers of a male *C. grandiflora* flower.

Seed dispersal

There are no observations of actual seed dispersal but mammals and birds appear to be attracted by the fruits and likely act as seed dispersers. Fruit bats such as *Cynopterus sphinx* (Vahl, 1797) feed on *C. grandis* fruits in Thailand (Elangovan et al. 2001; Ruby et al. 2000). Fruits of *C. grandis* are also taken up by birds (Bhatt and Kumar 2001) and eaten by humans (Voigt 1845). Elephants also feed on *C. grandis* (Mubalama 2000) and are possibly also seed dispersers. From its introduction to Pacific Islands, dispersal of *C. grandis* by humans is well-known (Muniappan et al. 2009). Human dispersal, in many cases likely to be intentional (Starr et al. 2010), also explains the occurrence of this species in the Neotropics and even in Missouri, USA. Some occurrences in Australia can also be explained by escape from gardens. Zimmermann reports feeding on *C. grandiflora* fruits by birds, small mammals but also snails and beetles (Zimmermann 1922a), the latter two unlikely being seed dispersers. The forest weaver *Ploceus bicolor* Vieillot, 1819 was observed to feed on fruits of *C. mackenii* (Bleher et al. 2003). Stanford and Nkurunungi (2003) report differing preference of *Coccinia* plant parts by gorillas. Whereas the gorillas feed on the leaves and fruit pulp of *C. mildbraedii* but not the seeds, they take only the leaves of *C. barteri*.

Successful seed germination in Munich Botanical Garden indicates that passage through a digestive tract is not necessary at least for *C. abyssinica*, *C. adoensis* var. *jeffreyana*, *C. grandiflora*, *C. grandis*, *C. hirtella*, *C. megarrhiza*, *C. microphylla*, *C. rehmannii*, *C. sessilifolia*, and *C. trilobata*. However, whether seeds would survive intestine passage and the role of endozoochoric dispersal is also not known.

Interaction with ants

Many species of *Coccinia* bear extranuptial glands (nectar producing glands outside of the flower) on the lower lamina of the leaves and/or on the bracts and probracts (Figs 7b, 8a). The glands are sunken into the surface and are surrounded by cells with a thicker cell wall (Muhammad Ilyas 1992; pers. observ.). Ants take up the sweet-tasting sap in *C. grandiflora* (Zimmermann 1922b) and in *C. grandis* (pers. observ.). Whereas Ilyas (1992) reports aggressive behavior of the ants on herbivores for Indian *C. grandis*, the present author could not observe this in Tanzanian *C. grandis*. Nieuwenhuis von Üxküll-Güldenbandt (1907) found a weak attraction of ants and heavy damage by herbivores in *C. grandis* in Bogor Botanical Garden (Java, Indonesia). In addition, Zimmermann (1922b) does not find aggressive behavior in *C. grandiflora* either but reports that the ants attacked a caterpillar he had placed onto the plant. Agarwal and Rastogi (2008), on the other hand, report a significant reduction in residence time of herbivores on the cucurbit *Luffa aegyptiaca* when ants are patrolling on the plant. Most likely, there is no close relationship to certain ant species as guardians, and plant-defense is carried out only by a few ant species. How *Coccinia* species without or few probracts, bracts or sublaminal extrafloral nectaries (e.g., *C. microphylla*) react when damage by herbivores occurs, is unknown. Agarwal and Rastogi (2008) found an increase of total numbers of extrafloral nectaries over time but did not discuss changes of nectary density as reaction to grazing.

Diseases and parasites

Some research has been undertaken on parasites and diseases for *C. grandis* for its status as crop but also as weed. As *C. grandis* is naturalized on several Pacific islands, in Australia, and the Neotropics, the plants can either overgrow other plants or represent a non-specific host for diseases of cucurbitaceous crops (Bamba et al. 2009; Muniappan et al. 2009). Its rapid growth can become problematic, as Pangelinan (2002) reports that *C. grandis* covered 35% of the vegetation of the island of Saipan only eleven years after its introduction.

Many different organisms are reported to live in, on, or to feed from *Coccinia* species. Beetle and fly larvae are either a disease for *Coccinia*, or in some cases, they are used to eradicate *C. grandis*. Fruits of *C. grandis* are a host for the larvae of the melon fly *Bactrocera* (= *Dacus*) *cucurbitae* (Coquillett, 1899), a tephritid fruit fly (Uchida et al. 1990). *Bactrocera cucurbitae* larvae usually populate the fruits but are also reported to hatch from

galls (Murthy 1959). The galls are not produced by these flies, however, but by the gall midge *Lasioptera* (= *Bimba*) *toombii* (Grover, 1962) (Bhatia and Mahto 1968). The gall infestation is interpreted as non-specific, as the female fly might not be able to differentiate between the gall and an unripe fruit, which would be the usual target. In addition, also the tephritid fruit fly *Dacus ciliatus* Loew, 1862 infests the galls, sometimes even together with *Bactrocera cucurbitae* (Bhatia and Mahto 1968). The galls in *C. grandis* do not only result from *Lasioptera toombii* but can also be produced by the Itonidid gall midge *Neolasioptera cephalandrae* Mani, 1934 (Dharmamaraju 1968), which is reported to be the major disease in *C. grandis* in India (Unni et al. 1976). The galls induced by *Neolasioptera cephalandrae* also appear to be gateway for a fungal infection with a mold, which is identified tentatively as *Cladosporium* sp. (Krishnamurthy 1984).

Other major cucurbit pests can also use *C. grandis* as a host such as *Diaphania* (= *Palpita*) *indica* (Saunders, 1851) (Lepidoptera: Pyralidae), *Aulacophora foveicollis* (Lucas, 1849) (Coleoptera: Chrysomelidae), *Leptoglossus australis* (Fabricius, 1775) (Hemiptera: Coreidae), *Aphis gossypii* Glover, 1877 (Hemiptera: Aphididae), *Liriomyza* spp. leafminers (Diptera: Agromyzidae), *Bemesia* spp. white flies (Hemiptera: Aleyrodidae) (Bamba et al. 2009), and *Epilachna vigintioctopunctata* (Fabricius, 1775) (Coleoptera: Coccinellidae) (Maurice and Ramteke 2012).

As a result of the damage that can be done to cucurbitaceous crops and of its weedy behavior on Pacific islands, larvae of the clearwing moth (Sesiidae) *Melittia oedipus* Oberthür, 1878 and the weevil (Curculionidae) species *Acythopeus burkhartorum* O'Brian, 1998 and *Acythopeus cocciniae* O'Brian, 1998 were introduced to Hawaii for biological pest control against *C. grandis* (Muniappan et al. 2002). Immediately after hatching from the eggs, *Melittia oedipus* larvae bore into the stems, where they live and pupate after two to four months (Chun 2002). This moth, originating from Zanzibar (Oberthür 1878), appears to be quite specific as larvae only rarely develop on *Cucumis sativus* L. (Chun 2001). Also *Zehneria guamensis* (Merrill) Fosberg, a Guam endemic, is not attacked by *M. oedipus* (Bamba et al. 2009; Reddy et al. 2009). As *C. grandis* is a noxious weed in Hawaii (Hawaiian Department of Agriculture 1992) active search for pests for biological control was undertaken, which led to the discovery of two new beetle species from Kenya: *Acythopeus burkhartorum* whose larvae produce galls in young shoots, and *A. cocciniae* whose larvae mine the leaves (Chun 2002; O'Brian and Pakaluk 1998). O'Brian and Pakaluk report a close morphological similarity of both *Acythopeus* species to *A. cucurbitae* (Marshall), which is a major pest on various cucurbitaceous crops in Africa, the Middle East, and South India.

Many crop plants are attacked by root parasites or diseases, but there is little known from *Coccinia*. Only root lesion nematodes *Pratylenchus dasi* Fortuner, 1985 (nom. nov. for *P. capitatus* Das & Sultana, 1979) and *P. crassi* Das & Sultana, 1979 were described from the soil around the roots of *C. grandis* (Das and Sultana 1979; Siddiqi 2000), but it is not known if they harm the plants.

The only known plant parasite growing on *Coccinia* is the hemiparasitic vine *Cuscuta chinensis* Lam., which is reported to grow on *C. grandis* in Gujarat, India (Patel and Patel 2010).

Table 5. List of fungi reported from *Coccinia* species (sorted by phylum of the fungus).

| Fungus | Host | Symptom | Citation |
|----------------------------------------------------------------------------------------------------|-----------------------|------------------------|--------------------------------------|
| <i>Plasmopara cubensis</i> (Berk. & M.A.Curtis) C.J.Humphrey (Peronosporales, Oomycota) | <i>C. grandis</i> | Downy mildew | Selby (1899) |
| <i>Alternaria pluriseptata</i> (P.Karst. & Har. ex Peck) Jørst. (Saccharomycetales, Ascomycota) | <i>C. grandis</i> | Fruit rot | Chagale and Bhale (2010) |
| <i>Cercospora coccinae</i> Munjal, Lall & Chona (Dothideales, Ascomycota) | <i>C. grandis</i> | Leaf spot disease | Rangaswami and Chandrasekaran (1961) |
| <i>Cercospora elaterii</i> Pass. | <i>C. grandis</i> | Leaf spot disease | Rangaswami and Mahadevan (2004) |
| <i>Colletotrichum gloeosporioides</i> (Penz.) Sacc. (Glomerellales, Ascomycota) | <i>C. grandis</i> | Fruit rot | Bhagavan Reddy and Reddy (1987) |
| <i>Colletotrichum orbiculare</i> (Berk. & Mont.) Arx | <i>C. grandis</i> | Anthrachnose fruit rot | Imbumi (2004) |
| <i>Corynespora cassiicola</i> (Berk. & M.A.Curtis) C.T.Wei (Pleosporales, Ascomycota) | <i>C. grandis</i> | Leaf blight | Philip et al. (1972) |
| <i>Curvularia pallescens</i> Boedijn (Pleosporales) | <i>C. grandis</i> | Black rot | Imbumi (2004) |
| <i>Erysiphe cichoracearum</i> DC. ex Merat (Erysiphales, Ascomycota) | <i>C. grandis</i> | Powdery mildew | Imbumi (2004) |
| <i>Fusarium moniliforme</i> J.Sheld. (Hypocreales, Ascomycota) | <i>C. grandis</i> | Fruit rot | Kapoor et al. (1981) |
| <i>Geotrichum candidum</i> Link (Pleosporales) | <i>C. grandis</i> | Fruit rot | Chagale and Bhale (2010) |
| <i>Sphaerotheca fuliginea</i> (Schltdl.) Pollacci (Erysiphales) | <i>C. grandis</i> | Powdery mildew | Imbumi (2004) |
| <i>Puccinia cephalandrae</i> Thümen (Uredinales, Basidiomycota) | <i>C. quinqueloba</i> | Rust | Berndt (2007) |
| <i>Puccinia cephalandrae-indicae</i> Syd. & P.Syd. | <i>C. grandis</i> | Rust | Berndt (2007) |
| <i>Puccinia physedrae</i> Syd. | <i>C. barteri</i> | Rust | Berndt (2007) |
| <i>Puccinia windhoekensis</i> Mennicken, Maier & Oberw. | <i>C. rehmannii?</i> | Rust | Mennicken et al. (2005) |
| <i>Rhizoctonia solani</i> Khun (Cantharellales, Basidiomycota) | <i>C. grandis</i> | Fruit rot | Bhagavan Reddy and Reddy (1988) |

Several fungi have been reported from *Coccinia* (Table 5). The rust fungus *Puccinia windhoekensis* Mennicken, Maier & Oberw. was described on *Coccinia rehmannii* (Mennicken et al. 2005), although Berndt (2007) noticed a great similarity of this rust to *P. ctenolepidis* Ramachar & Bagyanar. Berndt could not confirm the identity of the host specimen, so it seems to be likely that it was misidentified, since *Ctenolepis cerasiformis* looks quite similar to *C. rehmannii*.

There are several reports of plant viruses from *Coccinia* species. Purcifull and colleagues (1988) tested the infectability of several Cucurbitaceae to different plant viruses. They found that *C. grandis* can be infected by the papaya ringspot virus type W (PRSV-W) and the Trichosanthes virus but not by the cucumber mosaic virus, squash mosaic virus, watermelon mosaic virus-2, and the zucchini yellow mosaic virus. PRCV-W infections of *C. grandis* are also reported from several Pacific islands (Davis and Ruabete 2010).

Verma et al. (1983) suggest a yet undescribed mosaic virus, which is expressed in the occurrence of deformed leaves and a mosaic pattern in *C. grandis* leaves. A strain

of the Moroccan watermelon mosaic virus, a Potyvirus, can infest *Coccinia barteri* (Owolabi et al. 2012), whereas the infection of *C. sessilifolia* with this virus, maybe a different strain, failed (van der Meer and Garnett 1987).

Use, economic potential, and phytochemistry

Several *Coccinia* species are used by tribal communities, mainly as a food source but also for cultural applications (for details see species descriptions). *Coccinia grandis* is notable for its economic value (although often cited erroneously as *Coccinia cordifolia* or *C. indica*), whereas the importance of *C. abyssinica* is only regional. Other species are used by local tribes only.

Coccinia grandis is used in a wide variety of applications. The plant is well-known in India, where its fruits had an impact even in classical Sanskrit literature. The red fruits are regularly used to describe lips, such as those of a beloved wife, who is described by her husband in Kālidāsa's poem Meghadūta (Wilson 1867) or those of the goddess Sita and the god Rama in the epic Ramayana (Dutt 1891–1894). However, the fruits are also edible (raw when ripe and cooked when unripe) and are valued for their high content of carotenoids, esp. lycopene (Barua and Goswami 1979). Also young shoots and leaves are eaten as spinach and contain high amounts of lutein and other carotenoids (Addis et al. 2009; Wasantwisut and Viriyapanich 2003). The high carotenoid value is of special importance in developing countries, as vitamin A deficiency is widespread among young children and pregnant women (WHO 2009). Social marketing has proven to be valuable in promoting the use of *C. grandis* to prevent vitamin A deficiency (Chittchang et al. 1999). Domestication of *C. grandis* is in an early stage but promising cultivars are developed in South and SE Asia (Bharathi 2007; Engle et al. 1998; Ramachandran and Subramaniam 1983). Additionally, the leaves seem to be a good source of selenium and potassium, as well as vegetable protein (Xu et al. 2003; Xu et al. 2004). In Africa, *C. grandis* is mostly used from wild collections (Addis et al. 2009; Imbumi 2004). Contraindications to the use of *C. grandis* are also reported (Adanson 1757; Orech et al. 2005), but these might also be the result from either misidentification or regional chemo-varieties with differing amounts of secondary metabolites.

Coccinia grandis has been used in Indian traditional medicine for several hundred years (Nadkarni and Nadkarni 1976; Ramachandran and Subramaniam 1983). There are some studies that suggest a high potential for the use of *C. grandis* leaf extracts in diabetes treatment (Azad Khan et al. 1980; Kuriyan et al. 2008; Munasinghe et al. 2011). Parts of the observed effects are explained by inhibition of gluconeogenesis in the liver due to repression of glucose-6-phosphatase (Hossain et al. 1992) and fructose-1,6-bisphosphatase (Shibib et al. 1993). Also an activating effect on the promotor of the glucose transporter gene GLUT1 from rats is reported (Graidist and Purintrabipan 2009). Eshrat (2003) observes a positive effect of *C. grandis* in rats with hyperlipidemia, which is often connected to diabetes. However, its effectivity in

diabetes treatment and the overall experimental design is in dispute (Ramachandran and Subramaniam 1983; Sadikot 2009), and more research to test the medical value is necessary. Since 2005, more than 15 studies researched chemical compounds in *C. grandis* and tested their validity in folk medicine. Some applications by tribal people could be reproduced *ex situ* but research is still in its infancy. Suggested effects are, e.g., anti-anthelmintic (Dewanjee et al. 2007b), anti-tussive (Pattanayak and Sunita 2009), hepatoprotective (Moideen et al. 2011; Vadivu et al. 2008), antioxidative (Umamaheswari and Chatterjee 2008), antipyretic, analgesic, and anti-inflammatory (Niazi et al. 2009), anti-ulcerogenic (Mazumder et al. 2008), and antimicrobial (Bulbul et al. 2011; Dewanjee et al. 2007a; Farrukh et al. 2008; Shaheen et al. 2009). Antimicrobial activity is explained by the occurrence of a protease inhibitor (Satheesh and Murugan 2011). Observed xanthine oxidase inhibition and antiuricaemical activity (Umamaheswari et al. 2007) suggests a use for gout treatment. Female rats with hyperprolactinemia-caused infertility regain fertility when treated with an aqueous extract of *C. grandis* stems and leaves (Jha et al. 2010).

Coccinia abyssinica is mainly an Ethiopian tuber crop. Under the name *anchote*, its starch containing (c. 20%) tubers are an important staple food in the SW semi-humid highland regions (Aga and Badada 1997; Asfaw 1997; Hora 1995). Additionally, the tubers contain a relatively high amount of calcium, which might explain the local belief that the plant helps with repairing bone fractures and displaced joints (Hora 1995). Locally (around Dembi Dolo, Oromia), young shoots and leaves are also eaten (Hora 1995). Although the fruits of the cultivated landraces are not eaten (Getahun 1973), the use might be beneficial due to the carotenoid content of the fruits, which are likely to be comparable to those of *C. grandis*. However, fruits of wild races of *C. abyssinica* are already used (Asfaw and Tadesse 2001). In Wollega (W Ethiopia), *C. abyssinica* is also used to treat gonorrhea, tuberculosis, and cancer, as well as in traditional ceremonies and celebrations and for animal fattening (Gelmesa 2010). Currently, much effort is put into the development of *anchote* to increase the yield by selection of cultivars with larger tubers and by improving crop growing with better suited fertilizers (Abera and Guteta 2007; Bekele et al. 2014; Mengesha et al. 2012).

Also other species of *Coccinia* are used as food sources but if so, then only locally. In these species, such as *C. sessilifolia*, some wild landraces lack bitter substances (Bosch 2004; Dinter 1912). Bitterness in Cucurbitaceae is mainly caused by triterpenoids called cucurbitacins, although not all cucurbitacins are bitter. Cucurbitacins are often cytotoxic and often exist as β -glucosides (Miró 1995). All *Coccinia* species screened so far contain cucurbitacins, although the cucurbitacin type, organ, and time of expression differ greatly. Whereas fruits of *C. hirtella* and *C. quinqueloba* contain glycosidic cucurbitacin B, *C. adoensis* from South Africa contains aglycosidic cucurbitacin B only in unripe fruits and traces of cucurbitacin D but not in ripe fruits (Rehm et al. 1957). Unripe fruits of *C. rehmannii* and *C. sessilifolia* are not bitter, and therefore lack bitter cucurbitacins (Enslin et al. 1956). Njoroge and Newton (1994) tested the type and distribution of cucurbitacins within the plant in different Cucurbitaceae and found in Kenyan *C. adoensis* plants cucurbitacins H, I, and R in the stem but no cucurbitacins

in the roots, leaves, fruits, or seeds. *Coccinia trilobata* was found to contain the cucurbitacins B, D, and G in the stems, cucurbitacin D, H, I, and R in the leaves, and cucurbitacin G in the fruits, with no cucurbitacins in the roots and seeds. However, there seems to be much variability, as there are reports of edible (non-bitter) *C. trilobata* leaves (Coilly? 24, F. Msajiri 19). *Coccinia grandis* is also reported to contain cucurbitacin B (Bhakuni et al. 1962), and bitter and sweet fruited varieties are known. Guha and Sen (1973) find that cucurbitacin B has an antigibberelic effect, and its occurrence in seeds of *C. grandis* might enable or increase dormancy of the seeds.

Cucurbitaceae are also known for the occurrence of non-coded amino acids, such as citrulline in *Citrullus lanatus* (Thunb.) Matsum. & Nakai (Wada 1930). In a survey of such amino acids in Cucurbitaceae *C. grandis* and *C. hirtella* seeds were found to contain citrulline in low amounts, β -(pyrazol-1-yl)-L-alanine in very high amounts and the peptide Γ -L-glutamyl- β -(pyrazol-1-yl)-L-alanine in intermediate amounts (Dunnill and Fowden 1965). This pattern is similar to those that were found in *Diplocyclos tenuis* (Klotzsch) C. Jeffrey, *Acanthosicyos horridus* Welw. ex Hook.f., *Peponium hirtellum* Keraudren, *Ruthalicia eglandulosa* (Hook.f.) C. Jeffrey, *Dactyliandra welwitschii* Hook.f., and *Ctenolepis cerasiformis* (Stocks) C.B. Clarke (all in the same tribe as *Coccinia*).

Evolution and phylogeny

Recent phylogenetic analyses (Kocyan et al. 2007; Schaefer and Renner 2011b) show that *Coccinia* belongs to the tribe Benincaseae with a moderately supported sister group relationship to the genus *Diplocyclos*. However, the backbone of the tribe is not resolved and the relationship of the *Coccinia-Diplocyclos* clade to the other genera is unknown. *Citrullus*, *Cucumis*, or *Scopellaria* cluster with this clade but each without support, and morphological characters also do not seem to suggest any closer relatives.

Both phylogenies, plastid (Fig. 17) and the nuclear *LEAFY*-like 2nd intron (Fig. 18), suggest four major clades, although the backbone lacks bootstrap or posterior probability support (Holstein and Renner 2011b). The *Coccinia rehmannii* clade (IV) is well-supported in all phylogenies. The *Coccinia quinqueloba* group (II) is well-supported in the plastid DNA analysis, and consists of *C. hirtella*, *C. mackenii*, and *C. quinqueloba*. Additionally, *C. sessilifolia* belongs to this group, but it is only supported here by the nrDNA data. According to the nuclear data, the *C. barteri* clade (III) is nested within the *C. adoensis* clade (I). The plastid analysis tree separates these two clades but without support.

The *C. rehmannii* clade (IV) consists of five species. *Coccinia abyssinica* and *C. megarrhiza* are sister species from Ethiopia and semi-arid parts of N Kenya and Somalia (Fig. 20). They differ ecologically with the former species occurring in the semi-humid highlands and the latter one in the semi-arid lowlands. Both species differ weakly in morphology, and hybridization cannot be ruled out. The plastid haplotypes of both species do not form clades in the tree, which might be explained best by incomplete lineage sorting. The other three species of clade IV contain several plastid haplotypes and




Figure 17. Phylogenetic relationships in *Coccinia* based on five plastid DNA loci (*matK*, *ndhF-rpl32* intergenic spacer (IS), *rpl20-rps12* IS, *trnL* intron, *trnL-trnF* IS, *trnS-trnG* IS) obtained for 75 accessions from 24 species. Shown is the topology of the 50% majority rule consensus tree obtained from Bayesian analysis including simple gap coding for ingroup InDels. Numbers above the branches are posterior probability values ≥ 0.98 with values “with InDel coding” first, followed by “without InDel coding.” Numbers below the branches are bootstrap support values from ML analysis. Topologies from the different analyses were not contradictory, although some clades were not resolved without gap coding. Roman numerals indicate clades as discussed in the text: I = *C. adoensis* clade, II = *C. quinqueloba* group, III = *C. barteri* clade, and IV = *C. rehmannii* clade.

nrDNA sequences that each also do not form clades. The geographical distribution of the haplotypes is not assessed. The three species, however, are distinct. *Coccinia rehmannii* occurs in southern Africa while the other two species occur in NE Africa. In *Coccinia rehmannii* four forms can be recognized, which are included in the plastid phylogeny: (1) an inland form from dry habitats with small globose fruits (type form / var. *rehmannii*),




Figure 18. Phylogenetic relationships in *Coccinia* based on 505 nucleotides of the nuclear *LEAFY*-like 2nd intron, obtained for 37 accessions from 23 species analyzed under maximum likelihood (ML) and the GTR + Γ model. Numbers below branches refer to ML bootstrap support > 80% from 1000 replicates. The dots at nodes and behind the two accessions refer to uniquely shared indels. Roman numbers indicate clades as discussed in the text: I = *C. adoensis* clade, II = *C. quinqueloba* group, III = *C. barteri* clade, and IV = *C. rehmannii* clade.

(2) a form similar to var. *rehmannii*, but with larger globose fruits (described by Dinter and Gilg as *C. ovifera*), (3) a long-petiolate and long-peduncled coastal form from the (semi-)humid Southeast (described by Meeuse as var. *littoralis*), and (4) plants with oblong fruits occurring in all semi-humid areas from the Southeast to the northern parts in the periphery of the *C. rehmannii* distribution (*C. rehmannii* 5; here referred to *Coccinia rehmannii* aff. var. *littoralis*). None of these forms cluster together. The other two species differ morphologically and ecologically from each other: *C. trilobata* has, e.g., oblong fruits and occurs in the semi-humid uplands, and *C. microphylla* has globose fruits and occurs in the semi-arid lowlands (Fig. 33). Interestingly, *C. microphylla* does not differ morphologically from the *C. rehmannii* form from the dry inland. This scenario suggests incomplete lineage sorting and a speciation event with ecological differentiation in the northeastern Africa but not in southern Africa as intermediate collections between the four forms are found regularly. The distribution of these three species and the estimated age of this clade of 3.2 Ma suggest either a long distance dispersal or vicariance. As each of the three species contains several plastid haplotypes, vicariance is more likely, which indicates that semi-arid conditions might have prevailed between today's Tanzania and Zimbabwe. This has been suggested several times for different clades under the term "arid track" (Balinsky 1962; de Winter 1971).

The *C. quinqueloba* clade (II) is only supported in the nrDNA phylogeny, as plastid sequences of *C. sessilifolia* and its distinctly petiolate variety *variifolia* lack synapomorphies that support a closer relationship to any clade in *Coccinia*. The two varieties of *C. sessilifolia* occur in the semi-arid and sub-semi-humid inland (Fig. 40; see species description), whereas the other three species prefer more humid habitats in the Southeast (Fig. 30; see species description; Holstein and Renner 2011b). One species, *C. hirtella*, occurs in the rather open habitats, especially in the Drakensberg Mountains, which receive high amounts of rainfall. *Coccinia mackenii* occurs in remnant forest sites in the humid Southeast of southern Africa, whereas *C. quinqueloba* occurs only in coastal bushlands of the Eastern Cape, where it receives less precipitation than the other two species but has a more evenly distributed water availability all over the year (Holstein and Renner 2011b). As *C. mackenii* and *C. quinqueloba* do not co-occur but have similar ecologies, and as they only slightly differ morphologically but hardly produce hybrids (see chapter Hybridization and crossing experiments), a recent allopatric speciation event is probable. The lack of differentiation in the plastid sequences over 3500 bp between two accessions might support this hypothesis. In contrast to the *C. rehmannii* clade, all species of this clade occur exclusively in southern Africa, although the clade is older (c. 5.0 Ma vs. 3.2 Ma).

The *C. adoensis* clade (I) contains several morphologically and ecologically well differentiated species (Holstein and Renner 2011b). There are three subclades in the plastid tree with accessions having the name *Coccinia adoensis*. The type form (Fig. 21; see species description) from Ethiopia (no DNA sequences available) is morphologically inseparable from South African forms (*C. adoensis* 1 and 6). Geographically between those two populations, however, there are many populations that mostly differ gradually in length and density of trichomes. Two forms (with especially dense and long trichomes, respectively) could be assessed geographically and are accepted by the present author as varieties. The *Coccinia adoensis* var. *aurantiaca* accessions are neither in the plastid nor in the nuclear tree monophyletic but share a dense indumentum. These forms cluster in the nuclear tree with collections that have a less dense indumentum and thus are rather referred to as *C. adoensis* var. *adoensis* (Fig. 18). In the plastid tree, these collections cluster together with a Kenyan specimen of *C. adoensis* var. *adoensis* and var. *jeffreyana* (Fig. 17). *Coccinia adoensis* var. *jeffreyana*, however, shares the longer trichomes (Figs 3a, 5c) of some *C. senensis*, but it differs from these by lacking subulate calyx lobes and a 569 bp deletion in the *trnS*^{GCU}–*trnG*^{UUC} intergenic spacer region. However, one collection that does not differ morphologically from the variety *jeffreyana* (R.E. Gereau and C.J. Kayombo 3582) clusters within the East African forms of *C. adoensis*, which indicates either homoplasy of the trichome length or gene flow. Additionally, gene flow among the *C. adoensis* clades might also occur. Holstein and Renner (2011b) found a collection from Namibia (*C. adoensis* 5) that contained ITS sequences that are otherwise found exclusively in the South African and East African plastid haplotypes. Thus it can be suggested that all these forms belong to one widespread species, *C. adoensis*, which contains different plastid haplotypes. From this widespread species, several populations might have undergone ecological and morphological divergence. *Coccinia grandiflora* and *C. schliebenii* are nested within one *C. adoensis* subclade, and they occupy rather humid




Figure 19. Scenario of evolution in the *C. adoensis* clade. The green line surrounds today's distribution of *C. adoensis*. Blue lines surround today's distributions of *C. senensis* and *C. pwaniensis*. Blue arrows indicate peripatric speciation without shift in precipitation preference. Yellow arrows indicate speciation with shifts towards more arid habitats. Black arrows indicate speciation with shift towards more humid habitats.

habitats while *C. grandis* and *C. ogadensis* occupy more arid habitats. Some populations probably evolved parapatrically in former times with morphological shifts (*C. senensis*, *C. pwaniensis*, *C. samburuensis*) or evolved in allopatry (*C. intermedia*) (Holstein and Renner 2011b; Fig. 19). Some populations, however, did not diverge sufficiently to be taxonomically well-separated as a species, such as *C. adoensis* var. *aurantiaca*.

The *C. barteri* clade (III) mostly consists of rainforest species from West and Central Africa, except for the recently described *C. intermedia* (Holstein and Renner 2011a). *Coccinia mildbraedii* (including *C. ulugurensis*) also differs ecologically, as it occurs in mountain forest communities not in typical lowland rainforests as does the rest of the species. The phylogenetic position of *C. intermedia* is unclear as the resolution within this clade is generally poor. *Coccinia intermedia* shares morphological characters with *C. adoensis*, especially the open campanulate flowers. Both species occur in the same habitat type with the former occurring in West Africa and the latter north and east of the Central African rainforests. If the *C. barteri* clade is indeed nested in the *C. adoensis* clade, as suggested by the nuclear phylogeny, then it is possible that *C. intermedia* might have split allopatrically from a proto-*C. adoensis* species

and is sister to the rest of the *C. barteri* clade (Fig. 19). Then, the common ancestor of the other species of the *barteri* clade might have shifted the habitat preference towards perhumidity once and evolved allopatrically in refugia during arid periods of the Pliocene and Pleistocene. Alternatively, the habitat of *C. intermedia* would be explained as a reversal from a rainforest distributed common ancestor of the *C. barteri* clade. As the frequency of the Pleistocene climatic oscillations increased, reproductive isolation did not always occur, leading to weak morphological differentiation of interbreeding populations, such as in the polymorphic *C. barteri* (Holstein and Renner 2011b).

Identification of *Coccinia* species

Possible confusion with other genera

Some *Coccinia* species are easily confused with collections of other Cucurbitaceae genera (Table 6). The similarity is sometimes striking and without generative structures, one might need some experience to differentiate between the genera.

Characters for species discrimination

There is no character that is useful for all species. For example, whereas the direction of the calyx lobes can be a useful character for some species (e.g., *C. grandis*, *C. intermedia*, *C. keayana*), it is less useful in others (e.g., in the *C. quinqueloba* clade). Collections without flowers are harder to identify. In some cases it is almost impossible to discriminate between species if flowers are lacking. Identification of only vegetative material is often possible but needs experience. The indumentum can be a useful character; especially the trichomes (length, somewhat also the shape) on the abaxial side of the petiole and the lower leaf lamina can be helpful. However, the trichomes on the adaxial side of the petiole and the leaf margin do not seem to have any purpose for species identification.

Key to *Coccinia* species

The key is made from observations of herbarium material but also includes some characters from personal observations of living material and observations as given on herbarium labels. Fresh material is not needed, however, to use the key. The term ‘articulate’ refers to dried trichomes that appear wrinkled due to equatorially sunken cell walls (see Fig. 3a) but not to trichomes with ramifications, which have never been observed in *Coccinia*. In the living state, these trichomes are rather long and stiff. The term “dentate” refers to the sometimes colored structures (hydathodes?) at the leaf margin and leaf tip (Figs 6, 7a, 8a, 16a, 16b, 21, 39).

Table 6. African taxa that *Coccinia* species might be misidentified with.

| <i>Coccinia</i> species | Similar taxon | Differences |
|-----------------------------------------------------------------------|--------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>C. barteri</i> , <i>C. heterophylla</i> , <i>C. racemiflora</i> | <i>Bambekea</i> <i>racemosa</i> Cogn. | <i>B. racemosa</i> : petals free, has veins running along the leaf margin <i>Coccinia</i> : petals connate, veins not running directly along the margin |
| <i>C. barteri</i> , <i>C. heterophylla</i> , <i>C. racemiflora</i> | <i>Cogniauxia</i> spp. | <i>Cogniauxia</i> : petals free and up to 8 cm long, has veins running along the leaf margin, prophylls lanceolate, fruits up to 15 × 8 cm (L × D) large, seeds up to 2 cm long (Schaefer and Renner 2011a) <i>Coccinia</i> : petals connate and < 3.5(–4.5 cm), leaf veins not directly running along the leaf margin, probracts ovate or missing, fruit diameter < 4(–5) cm, if length > 6 cm, then not ovate, seeds < 0.8 cm long |
| W African rainforest species | <i>Ruthalicia</i> spp. | <i>Ruthalicia</i> : bracts lanceolate, petals free, seeds black or dark brown; <i>R. eglandulosa</i> with trichomes with a claret-red color (coloration is often at the ends of the long, centrally sunken-in cells) <i>Coccinia</i> : bracts ovate or missing, petals connate, seeds gray to beige; trichomes whitish, beige, yellowish or rarely light brownish |
| rainforest species | <i>Peponium</i> spp. | <i>Peponium</i> : petals free, male flowers with long-stretched hypanthium and three free stamens, which connect only with the long-stretched anthers, seeds dark colored; <i>P. vogelii</i> : sessile probracts and bracts are round and up to 3 cm long <i>Coccinia</i> : petals connate, male flowers with perianth tube of which the length does not exceed two times the diameter, three already connected filaments and a globose anther head, seeds gray to beige; short petiolate probracts and bracts ≤ 0.5 cm in C & W African species |
| <i>C. schliebenii</i> | <i>Luffa aegyptiaca</i> Mill. | <i>L. aegyptiaca</i> : mostly (2–)3–5-fid tendrils (check as many as possible), petals free, petals bright yellow (in <i>L. acutangula</i> (L.) Roxb. also dull yellowish), stamens 5 <i>C. schliebenii</i> : (1–)2-fid tendrils, petals connate, petals dull yellowish or yellow-orange, stamens 3 |
| <i>C. schliebenii</i> | <i>Lagenaria</i> spp. | <i>Lagenaria</i> : often tooth-like glands at the base of the lamina or along the petiole, trichomes > 1 mm, petals free and white, anthers serpentine <i>C. schliebenii</i> : never glands as above, trichomes < 1 mm, petals connate and dull yellowish or yellow-orange, anthers S-shaped |
| <i>C. adoensis</i> | <i>Eureiandra</i> spp. | <i>Eureiandra</i> : petals free, calyx lobes triangulate to lanceolate, stamens 5, seeds almost globose and whitish (Jeffrey 1967) <i>Coccinia</i> : petals connate, calyx lobes lineal, subulate to narrowly triangulate, stamens 3 in a central column, seeds grayish to beige, flattened |
| <i>C. microphylla</i> , <i>C. rehmannii</i> | <i>Ctenolepis</i> <i>cerasiformis</i> C.B.Clarke | <i>C. cerasiformis</i> : large roundish, sinuate-ciliate probract, petals < 5 mm (Jeffrey 1967; Meeuse 1962) <i>Coccinia</i> : probract < 3 mm, ovate or missing, petals > 1 cm |
| <i>C. microphylla</i> , <i>C. rehmannii</i> | <i>Dactyliandra</i> spp. | <i>Dactyliandra</i> : large roundish, sinuate-ciliate probract, petals < 5 mm; <i>D. stefaninii</i> (Chiov.) C.Jeffrey from N Africa lacks the probracts but the seed shape is conspicuously rounded (Jeffrey 1967; Meeuse 1962) <i>Coccinia</i> : probract < 3 mm, ovate or missing, petals > 1 cm; seeds asymmetrical (almost falcate) |

Habitats in this key (not the species descriptions) are given rather crudely and reflect the vegetation that would be found naturally. Savannas and woodlands (tree stands with not largely overlapping canopies) can also include mopane, but also dry forests (larger amounts of deciduous trees and overlapping canopies), deciduous thickets, tall grasslands, and secondary vegetation derived from these. “Rainforests” include gallery forests, semi-deciduous forests derived from rainforests, e.g., in relict areas, perhumid savanna types, and open areas, in which rainforest would be predominant if it was not for human impact, or swamps.

A local key for *Coccinia* from West Africa is provided separately by Holstein and Renner (2011a). If the plant is collected from outside of Africa, then it is *C. grandis*.

- 1 Mature leaves sessile (first leaves may be petiolate), rarely subsessile; alive usually bluish-green; glabrous; male flowers solitary or in few-flowered racemes, female flowers solitary; fruit long ovoid, elliptical to spindle-shaped; preferring dry habitats; from S Africa (Figs 2b, 7b, 13b)..... ***C. sessilifolia* var. *sessilifolia***
- 1* All leaves petiolate; plant not like in 1..... 2
- 2 Tendrils mostly bifid; usually forest species or from Drakensberg Mts or humid coastal bushland in SE Africa (in E and W Africa also in woodlands or savannas) 3
- 3 Plant with flowers 4
- 4 Corolla ≥ 4 cm long, calyx lobes > 3 mm long; ovaries and fruits long ovoid to cylindrical; E Africa or Ethiopia..... 5
- 5 Leaf surface usually glabrous, rarely with sparse weak thin trichomes on the abaxial side; leaves profoundly lobed (Fig. 8a); E African (rain) forests of C Zimbabwe and Mozambique to S Kenya ***C. grandiflora***
- 5* Leaf surface, at least below (secondary and tertiary veins) densely covered with small trichomes; leaf shallowly or rarely profoundly lobed; margin of humid forests and in forests; from N Mozambique to C–S Tanzania or Ethiopia to South Sudan ***C. schliebenii***
- 4* Corolla < 4 cm long..... 6
- 6 Calyx lobes (> 2.5 mm) subulate (Fig. 28), western C Africa... ***C. heterophylla***
- 6* Calyx lobes < 2.5 mm, if longer then from S Africa 7
- 7 Calyx lobes > 3 mm, plant from S Africa 8
- 8 Leaf lamina and stem usually densely covered with long (> 0.5 mm) trichomes; lamina profoundly lobate and lobulate; lobe tips usually rounded; pedicels covered with long (> 0.5 mm) trichomes (Fig. 13a) ***C. hirtella***
- 8* Leaf lamina and stem glabrous or rarely sparsely covered with long trichomes, with lobes often tapering into an acute tip, only side lobes with a slight lobule on outer side; pedicels glabrous..... ***C. mackenii***
- 7* Calyx lobes < 3 mm, plant not from S Africa 9
- 9 Flowers in lax many-(> 6 -)flowered racemes, western C Africa... ***C. racemiflora***

- 9* Flowers in dense racemes, few-flowered or on a long common peduncle that surpasses the length of the branched part; female flowers may also be solitary **10**
- 10 Male flowers on a long common peduncle that surpasses the length of the branched part; female flowers solitary with cylindrical ovary; mountain forests of Kivu Mts, Livingstone Mts, and Eastern Arc Mts, introduced into Kenyan high mts..... ***C. mildbraedii***
- 10* Male flowers in a raceme, in which the common peduncle is shorter than the branched part; female flowers in racemes, clustered or if solitary, then with subglobose to elliptical ovary **11**
- 11 Corolla campanulate, calyx lobes erect with recurved tips. Lower leaf surface at maturity often with white speckles and leaf margin with colored teeth. W African semi-humid savannas and woodlands ***C. intermedia***
- 11* Corolla urn-, cup-, funnel-shaped or narrow campanulate. Calyx lobes variable but not as above. Lower leaf surface rarely with white speckles, teeth on leaf margin not conspicuously colored. Rainforests of W Africa, C Africa, and in relict forests to Angola, Zambia?, W Tanzania, Uganda, and the Chimanimani Mts (Mozambique, Zimbabwe) ***C. barteri***
- 3* Plant with fruits or vegetative parts only **12**
- 12 Plant with fruits..... **13**
- 13 Fruit oblong to cylindrical (mature > 5 cm long), plant from E or NE Africa... **14**
- 14 Lower leaf surface, often also upper surface densely covered with short trichomes; N Mozambique, C and S Tanzania or W Ethiopian to SE South Sudanian mts..... ***C. schliebenii***
- 14* Upper leaf surface glabrous, lower leaf surface glabrous or rarely with some trichomes; plant from E Africa, incl. Kivu Mts and Chimanimani Mts (Mozambique, Zimbabwe); hard to differentiate in shared mountain ranges ... **15**
- 15 Probracts > 3.5 mm (Fig. 8a); corolla > 3 cm, calyx lobes > 3 mm; forests and forest relicts in S Kenya, Mt Meru to Usambara Mts, Eastern Arc Mts, SE Tanzania, N Mozambique, Chimanimani Mts (Mozambique, Zimbabwe) ***C. grandiflora***
- 15* Probracts < 3.5 mm; corolla < 3 cm, calyx lobes < 3 mm; forests of Kivu Mts, Livingstone Mts, and Eastern Arc Mts, introduced into Kenyan high mts
..... ***C. mildbraedii***
- 13* Fruits ovoid, if long elliptical, then from S Africa **16**
- 16 Fruits in lax racemes, plant from western C Africa ***C. racemiflora***
- 16* Fruits in dense racemes or solitary..... **17**
- 17 Plant from S Africa; fruits solitary..... **18**
- 18 Leaf surface and stem usually densely covered with long (> 0.5 mm) trichomes; lamina profoundly lobate and lobulate; lobe tips usually rounded (Fig. 13a) ..
..... ***C. hirtella***
- 18* Leaf surface and stem glabrous or sparsely covered with long (> 0.5 mm) trichomes, with lobes often tapering into an acute tip, only side lobes with a slight lobule on outer side..... ***C. mackenii***

- 17* Plant from W to C Africa to Chimanimani Mts (Mozambique, Zimbabwe)... **19**
- 19 Plant from western C Africa, not distinguishable with confidence without flowers *C. heterophylla* or *C. barteri*
- 19* Plant not from western C Africa **20**
- 20 Lower leaf surface at maturity often with white speckles and leaf margin with colored teeth when dry. W African semi-humid savannas and woodlands.....
..... *C. intermedia*
- 20* Lower leaf surface rarely covered with white speckles, teeth on leaf margin not conspicuously colored. Rainforests of W to C Africa to Chimanimani Mts (Mozambique, Zimbabwe) *C. barteri*
- 11* Plant vegetative only **21**
- 21 Lower leaf surface, often also upper surface densely conspicuously covered with short trichomes; N Mozambique, C and S Tanzania or W Ethiopian to SE South Sudanian mts *C. schliebenii*
- 21* Leaves glabrous, or if covered with trichomes, then they are long (> 0.7 mm) or inconspicuous..... **22**
- 22 Plant from S Africa **23**
- 23 Leaf surface and stem usually densely covered with long (> 0.5 mm) trichomes; leaves profoundly lobate and lobulate; lobe tips usually rounded (Fig. 13a) *C. hirtella*
- 23* Leaf surface and stem glabrous or sparsely covered with long (> 0.5 mm) trichomes, with lobes often tapering into an acute tip, only side lobes with a slight lobule on outer side *C. mackenii*
- 22* Plant from W to E Africa..... **24**
- 24 Plant from E African (incl. S Kenyan) rainforests or forest relicts, Mt Meru to Usambara Mts, Eastern Arc Mts to Chimanimani Mts (Mozambique/Zimbabwe); probracts > 3 mm (Fig. 8a) *C. grandiflora*
- 24* Plant from W, C, or E Africa, if from E Africa, then probracts < 3 mm.... **25**
- 25 From mountain forests of E Africa, incl. Kivu Mts *C. mildbraedii*
- 25* Rather from lowland rainforests from W Africa, C Africa, or from rainforests surrounding the Western Rift **26**
- 26 Plant from western C Africa not confidently distinguishable without flowers
..... *C. heterophylla*, *C. racemiflora*, or *C. barteri*
- 26* Plant not from western C Africa **27**
- 27 Lower leaf surface at maturity often with white speckles and leaf margin with colored teeth. W African semi-humid savannas and woodlands... *C. intermedia*
- 27* Lower leaf surface rarely with white speckles, teeth on leaf margin not conspicuously colored. Rainforests of W to C Africa to Chimanimani Mts (Mozambique, Zimbabwe) *C. barteri*
- 2* Tendrils usually simple, if not, then from semi-arid habitats or E and NE-African woodlands **28**
- 28 Leaves deeply palmately lobed with lineal lobes. If lobes lobulate, then leaf lamina at lobe base as broad as vein. E Ethiopia and C Somalia... *C. ogadensis*

- 28* Leaves profoundly, but not deeply lobed, or if deeply lobed, then lobes lanceolate or lobe base broader than vein.....**29**
- 29 Leaves deeply lobed with lanceolate lobes. Male flowers in racemes with short peduncle and pedicels. Rainforests of C Africa and around the Western Rift***C. subsessiliflora***
- 29* Plant not as above.....**30**
- 30 Leaves 7-lobate, rarely 5-lobate. Outer side of lobes serrate to lobulate with conspicuously colored tips. Calyx lobes > 4 mm, corolla > 2.5 cm, fruit ripe > 10 cm long, from Samburu area (C Kenya, E Africa).....***C. samburuensis***
- 30* Leaves cordate or 3-lobate. If 5-lobate, then plant not as above. Calyx lobes shorter than above or if longer then corolla shorter. Fruit shorter than above or if longer then from S Africa**31**
- 31 Plant glabrous; leaves usually subsessile, 5-lobate. Plant from coastal bushlands of Eastern Cape (South Africa)***C. quinqueloba***
- 31* Plant with trichomes or if glabrous, then from different region.....**32**
- 32 Plant glabrous, glaucous. Fruit long (> 6 cm) elliptical to spindle-shaped. Limpopo Province (South Africa) ***C. sessilifolia* var. *variifolia***
- 32* Plant with trichomes, or if glabrous then not glaucous and from different region..**33**
- 33 Lower leaf surface, often also upper surface densely conspicuously covered with short trichomes; bracts > 3 mm; calyx > 1 cm, corolla > 4 cm; fruit oblong to short cylindrical, > 5 cm long; N Mozambique, C and S Tanzania or W Ethiopian to SE South Sudanian mts.....***C. schliebenii***
- 33* Lower leaf surface glabrous or sparsely covered with trichomes. If densely covered with trichomes, then bracts < 3 mm and flowers smaller.....**34**
- 34 Plant with male flowers.....**35**
- 35 Plant glabrous, rarely some trichomes on adaxial petiole and leaf margin. Lower leaf lamina with pale (rarely also black when oxidized) glands towards the base, sometimes also between secondary veins. Margin of mature leaves with claret-red or brownish (black when dry) teeth, when young pale. Flowers solitary, rarely clustered, calyx lobes spreading to reflexed, corolla white or buff (Fig. 7a). NW (Senegal) to NE and E Africa, southern Arabia, S and SE Asia, naturalized or likely to become so in (sub)tropical regions worldwide ***C. grandis***
- 35* Plant with trichomes, or if glabrous then different from above.....**36**
- 36 Plant glabrous or rarely with soft multicellular trichomes. Flowers in lax ebracteate racemes, calyx lobes linear, > 2 mm long, in buds spreading, when mature reflexed (Fig. 32). Rainforests of W Africa (W of the Dahomey Gap) ***C. keayana***
- 36* Plant not as above.....**37**
- 37 Plant glabrous. Leaves cordate to subhastate, rarely 3-lobate. Flowers in ebracteate racemes. Calyx lobes erect, at base broader than 0.75 mm, corolla urceolate. Rainforests of W Africa..... ***C. longicarpa***
- 37* Plant different and not from W Africa, or if from W Africa then calyx lobes narrower or spreading to reflexed.....**38**
- 38 Plant glabrous, at maturity often with white speckles on stem, petiole, and lower leaf lamina. Flowers in racemes or 1 solitary. Calyx lobes erect with

- recurved tips. Corolla campanulate. Semi-humid savannas and woodlands of W Africa *C. intermedia*
- 38* Plant not from W Africa or if so, then rainforest species (sometimes hard to distinguish from *C. intermedia*), or lower leaf surface conspicuously covered with trichomes 39
- 39 Plant glabrous (or puberulous), leaves usually coriaceous. W or C (or western E) African rainforests 40
- 40 Male flowers in racemes with common peduncle shorter than racemose part. Lowland rainforests or rainforest relicts in higher altitudes or along rivers.....
..... *C. barteri*
- 40* Male flowers in racemes with common peduncle longer than racemose part. Mountain forests from Kivu Mts, Eastern Arc Mts, Livingstone Mts, also introduced in Kenyan high mts *C. mildbraedii*
- 39* Plant conspicuously covered with trichomes or if glabrous, then leaves papery or from NE, E, or S Africa 41
- 41 Plant from S Africa (except C and N Mozambique) 42
- 42 Plant (esp. stem oder petioles) with or without white speckles, perianth tube/hypanthium with long (> 0.7 mm) trichomes or if glabrous then calyx lobes > 2 mm (Fig. 10a) *C. rehmannii*
- 42* Plant without white speckles, perianth tube/hypanthium with short (< 0.7 mm) trichomes or if glabrous then calyx lobes < 2 mm (Fig. 21)
..... *C. adoensis* var. *adoensis*
- 41* Plant from E (incl. C and N Mozambique), NE, or NC Africa 43
- 43 Flowers clustered, common peduncle < 1 cm, if flower solitary then pedicel usually < 1 cm. NE Africa (incl. N Tanzania) 44
- 44 Upper and lower leaf surface rather densely covered with multicellular trichomes. Plant usually from higher elevations of N Tanzania and Kenya *C. trilobata*
- 44* Upper leaf surface pustulate but without trichomes, or with minute trichomes from pustules. Plant rather from dry habitats and lower elevations. May be hard to distinguish 45
- 45 Leaf margin in mature leaves with conspicuously colored teeth. Plant densely covered with long trichomes that appear articulate when dry (Fig. 9)
..... *C. megarrhiza*
- 45* Leaf margin in mature leaves without conspicuously colored teeth. Plant less densely covered with trichomes or if densely, then trichomes minute (< 0.2 mm) or if longer then not appearing articulate when dry (Fig. 2a) *C. microphylla*
- 43* Flowers in racemes with peduncle > 1 cm (if smaller then from C Tanzania), or if solitary then either pedicel > 1 cm or plant from C Tanzania 46
- 46 Calyx lobes subulate to narrowly triangulate with pointed tip, > 2.5 mm. Petiole and lower leaf surface not puberulous. Plant from E Africa (Tanzania, Mozambique, Malawi) 47
- 47 Leaves 3-lobate, distinctly petiolate, often with few short trichomes on the main nerves of the lower leaf surface. Racemes with > 8 flowers. Coastal forests of Kenya or NE Tanzania (Fig. 23) *C. pwaniensis*

- 47* Leaves subcordate to 3- or 5-lobate, subsessile or distinctly petiolate (Fig. 6). Lower leaf surface glabrous or nerves with short (wart-like) to long trichomes. Racemes < 10 flowers (Fig. 39). SE Tanzania, C and N Mozambique, or Malawi (Fig. 23) ***C. senensis***
- 46* Calyx lobes < 2.5 mm or if longer, then not pointed (may be lineal though) or petiole and lower leaf lamina puberulous, or plant from NE Africa (Kenya, Ethiopia, Somalia) **48**
- 48 Plant with long (> 0.5 mm) trichomes or with short, narrowly conical trichomes, calyx lobes > 2 mm, lineal. NE Africa..... **49**
- 49 Apex of the cordate leaf or central lobe tapering into a long, acute tip. Male flowers solitary or in racemes with a long common peduncle. Plant from high elevations (Fig. 20)..... ***C. abyssinica***
- 49* Apex of leaf or central lobe retuse, obtuse, or rather abruptly tapering into a short acute tip (Fig. 9). Male flowers solitary, clustered or if in racemes, then peduncle short. Plant rather from lower elevations (Fig. 20)... ***C. megarrhiza***
- 48* Plant glabrous or with short trichomes (< 0.8 mm), if with longer trichomes then not from NE Africa. Calyx lobes < 2.5 (–3.5) mm long. Taxa in E Africa not easily distinguishable (complex around *C. adoensis*) **50**
- 50 Plant with long (> 0.8 mm) trichomes (Figs 3a, 5c), calyx lobes 1.5–3.5 mm long but not with pointed tip. Malawi, C, and S Tanzania, maybe also N Mozambique (Fig. 23) ***C. adoensis* var. *jeffreyana***
- 50* Plant glabrous or with short (< 0.8 mm) trichomes..... **51**
- 51 Lower leaf surface and usually also herbaceous stems, petioles, and upper leaf surface densely covered with short (< 0.5 mm) trichomes. Peduncle often shorter than pedicelled part. Calyx lobes < 2 mm. Corolla orange, rarely yellow? E Africa (C Tanzania; Fig. 23) ***C. adoensis* var. *aurantiaca***
- 51* Lower leaf surface glabrous to densely covered with trichomes, but if so then peduncle longer than pedicelled part. Calyx lobes usually < 2 mm. E, NE, or NC Africa (Fig. 21) ***C. adoensis* var. *adoensis***
- 34* Plant with female flowers, fruits or vegetative..... **52**
- 52 Plant with female flowers **53**
- 53 Flowers solitary or in ebracteate racemes. Calyx lobes lineal, spreading in buds, reflexed in mature flowers, > 2 mm long. W African rainforests (W of Dahomey Gap) (Fig. 32) ***C. keayana***
- 53* Plant not as above. If with spreading to reflexed calyx lobes, then not from rainforest regions or < 2 mm long **54**
- 54 Ovary cylindrical. Calyx lobes broader than 0.75 mm at base, corolla urceolate. W African rainforests..... ***C. longicarpa***
- 54* Ovary shortly elliptical, (ob-)ovoid or globose, if cylindrical then not from W African rainforests. Calyx lobes narrower at base **55**
- 55 Flowers in bracteate or ebracteate racemes or solitary. Ovary globose or (ob-)ovoid, if longer then from E Africa. Calyx lobes < 2 mm long. W and C Africa but also in rainforest relicts or mountain forests in E Africa **56**

- 56 Female flowers solitary or in racemes. Corolla cup-, urn- or funnel-shaped, not open campanulate. Ovary globose to (ob-)ovoid. W or C (or western E) Africa. Lowland or in relict rainforests in highlands ***C. barteri***
- 56* Female flowers solitary. Corolla cup-shaped to campanulate. Ovary long spindle-shaped to oblong. Mountain forests of Kivu Mts, Eastern Arc Mts, Livingstone Mts, also introduced in Kenyan high mts ***C. mildbraedii***
- 55* Flowers solitary. Calyx lobes > 2 mm or if shorter then plant not from rainforests from regions as above.....57
- 57 Calyx lobes spreading to reflexed, lower leaf surface glabrous with pale glands between main veins. Leaf margin with colored teeth (Fig. 7a). Calyx lobes lineal, in buds spreading, later reflexed. Corolla white or buff. Plant from NW (Senegal) to NE and E Africa, southern Arabia, S and SE Asia, naturalized or likely to become so in (sub)tropical regions worldwide but natively not from C or S Africa.... ***C. grandis***
- 57* Plant with trichomes or if glabrous, then with darkish glands or without glands on lower leaf surface. Calyx lobes not reflexed. Corolla in various colors but not snow-white.....58
- 58 Calyx lobes erect with recurved tips, lower leaf lamina with dark glands between veins, sometimes with white pustules on veins and petiole. Margin of mature leaves with colored teeth. Plant from woodlands or savannas of W Africa.... ***C. intermedia***
- 58* Plant not as above and not from W Africa or if so, then not with white pustules and dark teeth.....59
- 59 Calyx lobes subulate to narrowly triangulate with pointed tip, > 2.5 mm (Fig. 39); petiole and lower leaf surface not puberulous. E Africa (Tanzania, Mozambique, Malawi; Fig. 23).....60
- 60 Leaves 3-lobate, distinctly petiolate, often with few short trichomes on the main nerves of the lower lamina. Coastal forests of Kenya or NE Tanzania ..
.....***C. pwaniensis***
- 60* Leaves cordate to 3- or 5-lobate, subsessile or distinctly petiolate. Lower leaf lamina glabrous or nerves with short (wart-like) to long articulate trichomes. SE Tanzania, C and N Mozambique, or Malawi ***C. senensis***
- 59* Calyx lobes < 2.5 mm or if longer, then not with pointed tip or then petiole and lower leaf lamina puberulous, or plant from NE Africa (Kenya, Ethiopia, Somalia).....61
- 61 Plant from S Africa (S Angola, Zimbabwe, C Mozambique and further S)...62
- 62 Plant (esp. stem and petioles) with or without white speckles, perianth tube/hypanthium with long (> 0.7 mm) trichomes or if glabrous then calyx lobes > 2 mm (Fig. 10a)..... ***C. rehmannii***
- 62* Plant without white speckles, perianth tube/hypanthium with short (< 0.7 mm) trichomes or if glabrous then calyx lobes < 2 mm (Fig. 21).....
..... ***C. adoensis* var. *adoensis***
- 61* Plant from NC, NE or E Africa63
- 63 Plant with long (> 0.5 mm) trichomes or straight, narrowly conical trichomes or if trichomes short (< 0.2 mm) then ovary globose. Plant from NE Africa (Ethiopia, Kenya, Somalia, N Tanzania).....64

- 64 Apex of leaf or central lobe tapering into a long, acute tip. Plant from high (> 800 m) elevations of Ethiopia (Fig. 20) *C. abyssinica*
- 64* Apex of leaf or central lobe retuse, obtuse, or rather abruptly tapering into a short, acute tip. Plant rather of dry habitats in lower elevation.....65
- 65 Plant rather densely covered with long (> 0.5 mm) trichomes that appear articulate when dry. Leaf apex retuse, obtuse, or rather abruptly tapering into a short acute tip (Fig. 9). Leaf margin of mature leaves with dark glands. Ovary ellipsoid, never globose. N Kenya, Ethiopia and likely also Somalia..... *C. megarrhiza*
- 65* Plant rather laxly covered with trichomes, if denser then trichomes usually minute (< 0.2 mm), if longer then not appearing articulate when dry. Leaf apex rarely obtuse (e.g., around the Usambaras), often abruptly tapering into a short acute tip. Leaf margin never with dark glands. Ovary globose, rarely (ob-)ovoid. N Tanzania, Kenya, Ethiopia and likely also Somalia (Fig. 2a) *C. microphylla*
- 63* Plant glabrous or with short (< 0.5 mm) trichomes, if with longer trichomes then not from NE Africa, ovary not globose66
- 66 Leaves on upper lamina with short trichomes (Fig. 8b). Calyx lobes 2–5 mm long. N Tanzania or Kenya..... *C. trilobata*
- 66* Leaves on upper lamina glabrous (but with white pustules) or if with short trichomes, then from C Tanzania. Calyx lobes < 2 mm or if longer, then lower leaf lamina with long (> 0.8 mm) trichomes or puberulous. Taxa in E Africa not easily distinguishable (complex around *C. adoensis*)67
- 67 Plant with long (> 0.8 mm) trichomes that appear articulate when dry (Figs 3a, 5c), calyx lobes 1.5–3 mm long but not pointed (as in Fig. 39). C and S Tanzania, Malawi, maybe also N Mozambique (Fig. 23)
..... *C. adoensis* var. *jeffreya*
- 67* Plant glabrous or with short (< 0.8 mm) trichomes only.....68
- 68 Lower sometimes also upper leaf lamina densely covered with short trichomes. Ovary densely covered with short (< 0.5 mm) trichomes. Calyx lobes < 2 mm. Corolla orange, rarely yellow? E Africa (C Tanzania; Fig. 23).....
..... *C. adoensis* var. *aurantiaca*
- 68* Lower leaf lamina glabrous or covered with short trichomes (Fig. 21). Ovary glabrous or only laxly (inconspicuously) covered with short trichomes. Corolla yellowish, pinkish or orange. E, NE, or NC Africa (Fig. 22) *C. adoensis* var. *adoensis*
- 52* Plant with fruits only or vegetative.....69
- 69 Plant with fruits.....70
- 70 Fruit long elliptical to cylindrical (> 8 cm). Forest species.....71
- 71 Plant from West African rainforests *C. longicarpa*
- 71* Plant from mountain forests of Kivu Mts, Eastern Arc Mts, Livingstone Mts, also introduced in Kenyan high mts..... *C. mildbraedii*
- 70* Fruit globose to oblong (< 8 cm)72
- 72 Fruit globose. Plant from dry habitats.....73
- 73 Plant from N Tanzania, Kenya, S and SE Ethiopia or Somalia... *C. microphylla*
- 73* Plant from S Africa *C. rehmannii*

- 72* Fruit obovoid to oblong, if (sub-)globose, then from humid habitats.....74
- 74 Leaf margin at maturity with colored teeth (blackening when dry), lower leaf surface glabrous and with pale glands between main veins, petioles and veins at maturity often with white pustules. Fruit (ob-)ovoid to elliptical. Plant natively not from C or S Africa..... ***C. grandis***
- 74* Lower leaf surface without glands or with darkish glands or if with pale glands, then mature leaves without colored teeth on leaf margin.....75
- 75 Fruit subglobose to obovoid-elliptical, in raceme or if solitary, then rainforest species.....76
- 76 Fruit in ebracteate raceme or solitary. Plant from W Africa (W of Dahomey Gap) (hardly distinguishable in shared distribution range) ***C. keayana* or *C. barteri***
- 76* Fruit in bracteate or ebracteate raceme. Plant from W and C Africa and in relict rainforest patches along the Western Rift..... ***C. barteri***
- 75* Fruit solitary or 1–3 clustered but not in raceme. Plant not from rainforests .77
- 77 Plant glabrous, at maturity often with white speckles on stem, petiole, and lower leaf lamina. Fruit subglobose to obovoid-elliptical, solitary or 1–3 clustered. Semi-humid savannas and woodlands of W Africa..... ***C. intermedia***
- 77* Plant different and not from W Africa. If from (eastern) W Africa then fruit often with sterile apex (“beak”)78
- 78 Fruit elliptical to oblong, often with sterile apical tip (“beak”). Unripe with dark green/light green longitudinal stripes or mottling. Seeds rather lenticular and with symmetrical shape (Fig. 14a). Lower leaf lamina glabrous or with trichomes, if trichomes appearing articulate and > 0.5 mm, then from C and S Tanzania, maybe also Malawi or N Mozambique79
- 79 Leaves 3-lobate. Leaf surface glabrous but veins and petiole often with few short trichomes. Plant from coastal forests of SE Kenya to E Tanzania (Fig. 23) ***C. pwaniensis***
- 79* Plant different or from different region (hardly distinguishable)80
- 80 Plant with long (> 0.8 mm) trichomes (Figs 3a, 5c) or if reduced, then stiff but not bent (as in Fig. 21), warty or subglabrous. C and S Tanzania, Malawi or C to N Mozambique (Fig. 23)..... ***C. senensis* or *C. adoensis* var. *jeffreya***
- 80* Plant glabrous or with short (< 0.8 mm), but not warty or articulate appearing trichomes.....81
- 81 Stem, petiole, lower leaf lamina, and ovary/young fruit densely covered with short trichomes. C Tanzania (Fig. 23) ***C. adoensis* var. *aurantiaca***
- 81* Stem, petiole, lower leaf lamina glabrous or with short trichomes, but young fruit only with lax indumentum. S, E, NE, or NC Africa (Fig. 22) ... ***C. adoensis* var. *adoensis***
- 78* Fruit obovoid, shortly to long elliptical, but not oblong and not with conspicuous sterile apical tip (“beak”). Unripe fruits with whitish longitudinal mottling that often has a dark green halo. Seed face rather flat, shape often asymmetrical (Fig. 14b, c). S Africa or NE Africa, incl. Kenya and N Tanzania..... 82
- 82 Plant from S Africa ***C. rehmannii***
- 82* Plant from NE Africa, incl. Kenya and N Tanzania.....83

- 83 Upper leaf lamina with fine, short trichomes. Lower leaf lamina with rather shortly (< 0.8 mm) articulate (Fig. 8b) or with narrowly conical trichomes. Plant from higher elevations of N Tanzania and S to C Kenya... ***C. trilobata***
- 83* Upper leaf lamina glabrous (but with pustules), rarely with narrowly conical trichomes. Lower leaf lamina with often long (> 0.8 mm), articulate or with narrowly conical trichomes **84**
- 84 Apex of leaf or central lobe tapering into a long acute tip. Plant from high elevations (> 900 m) (Fig. 20) ***C. abyssinica***
- 84* Apex of leaf or central lobe retuse, obtuse, or rather abruptly tapering into a shortly acute tip (Fig. 9). Plant rather from low elevations **85**
- 85 Plant rather densely covered with trichomes that appear articulate when dry. Leaf apex retuse, obtuse, or rather abruptly tapering into a shortly acute tip (Fig. 9). Leaf margin of mature leaves with dark teeth. N Kenya, Ethiopia and likely also Somalia (Fig. 20) ***C. megarrhiza***
- 85* Plant rather laxly covered with trichomes or if densely, then trichomes minute (< 0.5 mm). Leaf apex, rarely obtuse (e.g., around the Usambaras), often abruptly tapering into a short acute tip. Leaf margin never with dark teeth. Fruit globose to elliptical. N Tanzania, Kenya, Ethiopia and likely also Somalia (Figs 2a, 33)... ***C. microphylla***
- 69* Plants with vegetative characters only **86**
- 86 Plant glabrous, lower leaf surface with pale glands (if strongly oxidized then also dark, but then leaf margin also with black markings/teeth) between veins, veins at maturity often with white speckles. Leaf margin at maturity with colored teeth (Fig. 7a). Plant from W to NE Africa, incl. Kenya and N Tanzania or from outside of Africa ***C. grandis***
- 86* Plant with trichomes or if glabrous, then with darkish glands between veins or without glands on lower leaf surface **87**
- 87 Plant from W Africa **88**
- 88 Leaves cordate to 5-lobate, rarely broader than 10 cm, rather papery, lobes triangulate to narrowly lanceolate or oblong. Lower leaf surface without white speckles, glabrous or with often short, bent trichomes (Fig. 21). Semi-humid savannas and woodlands of N Cameroon, N Nigeria, distribution in the area imperfectly known ***C. adoensis* var. *adoensis***
- 88* Leaves cordate to subhastate to 3- or 5-lobate, mature often > 10 cm wide. Leaf lobes triangulate to broad lanceolate, but not narrowly lanceolate or oblong. Rainforest species or if from semi-humid savannas or woodlands (rarely dry forests), then margin of mature leaves with conspicuously colored teeth and lower leaf lamina often with white speckles **89**
- 89 Plant glabrous, at maturity often with white speckles on stem, petiole, and lower leaf surface. Margin of mature leaves with conspicuously colored teeth. Semi-humid savannas and woodlands of W Africa ***C. intermedia***
- 89* Plant not as above, from rainforests or gallery forests. Species not confidently distinguishable ***C. barteri*, *C. keayana* or *C. longicarpa***
- 87* Plant not from W Africa **90**

- 90 Leaves coriaceous. Plant glabrous or puberulous on abaxial side of petiole. Rainforest or mountain forest plant from C Africa or western E Africa (along the Western Rift, Livingstone Mts, Eastern Arc Mts), plants vegetatively hardly distinguishable **91**
- 91 Lowland rainforest (in relict sites of western E Africa also in mountains) plant from C Africa, incl. areas around Kivu Mts, Chimanimani Mts, and forests (mountain ranges) along the Western Rift, incl. Uganda *C. barteri*
- 91* Plant from mountain forests of Kivu Mts, Eastern Arc Mts, Livingstone Mts, also introduced in Kenyan high mts..... *C. mildbraedii*
- 90* Plant not from C African rainforests **92**
- 92 Plant from S Africa **93**
- 93 Lower leaf surface with usually bent trichomes (Fig. 21), never white speckled, rarely subglabrous..... *C. adoensis* var. *adoensis*
- 93* Lower leaf surface with straight, in herbarium collections often articulate appearing trichomes, often with white speckles towards maturity or glabrous
..... *C. rehmannii*
- 92* Plant from E, NE, or NC Africa **94**
- 94 Plant from NE or NC Africa **95**
- 95 Teeth on leaf margin conspicuously colored. NE Africa..... **96**
- 96 Apex of leaf or central lobe tapering into a long acute tip. Plant from high elevations (> 900 m; Fig. 20) *C. abyssinica*
- 96* Apex of leaf or central lobe retuse, obtuse, or rather abruptly tapering into a shortly acute tip (Fig. 9). Plant rather from lower elevation (< 1200 m) (Fig. 20).....
..... *C. megarrhiza*
- 95* Teeth on leaf margin not conspicuously colored. Plant from NC or NE Africa.. **97**
- 97 Lower leaf surface glabrous or with short (often bent) trichomes, cordate to deeply lobate (Figs 2a, 21). Plant from NC and NE Africa (incl. N Tanzania)..... **98**
- 97* Lower leaf surface with long (> 0.5 mm) trichomes that appear articulate when dry (such as in Figs 8b, 9), or with narrowly conical trichomes. Plant from NE Africa..... **99**
- 98 Stem sometimes pustulate. Leaf shape variable, if lobate then lobes extending and not pointing forward, lobes not oblong to linear. Leaves usually with trichomes, often also on upper lamina and then minute (< 0.2 mm). Trichomes on lower lamina often not only restricted to the veins. (Fig. 2a) Plant rather from low elevation drylands (rarely in higher elevations) of NE Africa (incl. N Tanzania) ... *C. microphylla*
- 98* Stem glabrous or with short (< 0.5 mm) trichomes but not pustulate. Leaf shape variable (cordate to deeply lobate) but also with oblong to linear lobes that point forward. Leaves glabrous or with (often bent) trichomes, on upper leaf lamina rarely beset with trichomes in this distribution area. Trichomes on lower leaf lamina usually restricted to the veins (Fig. 21). Plant from woodlands and semi-humid habitats..... *C. adoensis* var. *adoensis*
- 99 Leaves cordate to lobate but not lobulate. If profoundly lobate, then central lobe lanceolate or ovate tapering into an acute tip. Lower leaf surface with long (> 0.5 mm) trichomes that appear articulate when dry, or with narrow

- conical trichomes. Plant from higher elevations (> 900 m) or cultivated, from Ethiopia (Fig. 20) *C. abyssinica*
- 99* Leaf reniform to lobate, rarely lobulate. Apex of leaf or central lobe retuse, obtuse, rather abruptly tapering into a shortly acute tip (Fig. 9), if longer tapering into an acute tip then lobes lobulate. Lower leaf surface with long (> 0.5 mm) trichomes that appear articulate when dry, only rarely with conical trichomes. Plant rather from lower elevations (< 1200 m) (Fig. 33) *C. microphylla*
- 94* Plant from E Africa..... **100**
- 100 Leaves 3- or 5-lobate. Lobes extending, not pointing towards apex, broadly triangulate, elliptical, ovate or somewhat angulate but not narrow, oblong, or lineal. Upper and lower leaf surface with short, white trichomes that appear articulate when dry (Fig. 8b). Plant from higher elevations of N Tanzania and Kenyan highlands (Fig. 33)..... *C. trilobata*
- 100* Upper leaf surface glabrous (but pustulate) or if with trichomes then leaf shape different or from different region..... **101**
- 101 Stem sometimes pustulate. Leaf shape variable, if lobate then lobes extending and not pointing forward, lobes not oblong to linear. Leaves usually with trichomes, often also on upper lamina and then minute (< 0.2 mm). Trichomes on lower lamina often not only restricted to the veins. (Fig. 2a) Plant rather from low elevation drylands (rarely in higher elevations) of E Africa (Kenya, N Tanzania) ... *C. microphylla*
- 101* Stem not pustulate, leaves with or without oblong to elliptical lobes. If upper lamina densely covered with minute trichomes then from C Tanzanian woodlands. If plant with long articulate appearing trichomes then from highlands of C to S Tanzania, Malawi or N Mozambique..... **102**
- 102 Leaves 3-lobate (rather small auriculate), upper surface glabrous (but pustulate), lower lamina glabrous, but often with short trichomes on main veins. Coastal forests of SE Kenya and NE to E Tanzania (Fig. 23) *C. pwaniensis*
- 102* Plant not as above or from different area (in some cases hard to distinguish) ... **103**
- 103 Lower leaf surface with long (> 0.5 mm) trichomes that appear articulate when dry or reduced to warts, rarely almost glabrous; sometimes leaves sessile (Fig. 23) *C. senensis* or *C. adoensis* var. *jeffreyana*
- 103* Lower leaf surface glabrous or with short, thin, straight or bent trichomes **104**
- 104 Lower leaf surface (also often upper lamina), petiole and stem rather densely covered with short trichomes. C Tanzania *C. adoensis* var. *aurantiaca*
- 104* Lower leaf lamina glabrous or covered with trichomes, if densely then upper lamina glabrous (but pustulate) or with few straight trichomes but not tomentose. Widespread in E Africa *C. adoensis* var. *adoensis*

Taxonomic treatment

Herbarium abbreviations follow Index Herbariorum (<http://sciweb.nybg.org/science2/IndexHerbariorum.asp>). Digital collections were accessed from the homepages of the corresponding herbaria, except for “JPS” (= JStor Plant Science; <http://plants.jstor.org/>) and “CVH” (= Chinese Virtual Herbarium; <http://www.cvh.org.cn/>).

Species concepts in this treatment mainly follow the morphospecies concept but also include ecological aspects (habitats) and biogeography. Apart from easily recognizable distinct forms, it was tried to include molecular data (plastid and nuclear; Figs 17, 18) from as many forms as possible to check whether they cluster together or not. Accessions in polytomies are treated as one species as long as they are not morphologically or ecologically (habitat) distinct or are distantly distributed, if not contra-indicated otherwise (e.g., full crossing compatibility in Asian and African *C. grandis*). Names have been synonymized if no character was found to separate confidently the collections from the type material. Names have been changed in status (in this treatment to varieties), when characters to separate the collections change in degree, rather than absence/presence.

The minimum leaf size and petiole length were taken from leaves on the same node as open flowers or fruits. Leaf length is measured from the attachment point of the petiole on the lamina to the apex.

***Coccinia* Wight & Arn., Prodr. fl. Ind. orient.: 347. 1834.**

Cephalandra Schrad. ex Eckl. & Zeyh., Enum. pl. afric. austral. 2: 280. 1836.

Type species: *Bryonia quinqueloba* Thunb.

Physedra Benth. & Hook.f., Gen pl. 1(3): 827. 1867. Indirectly lectotypified by Jeffrey (1962: 361)

Type species: *Physedra heterophylla* Hook.f.

Staphylosyce Benth. & Hook.f., Gen pl. 1(3): 828. 1867.

Type species: *Staphylosyce barteri* Hook.f.

Type species. see *Bryonia grandis* L.

Description. Dioecious. Perennial climbers or creepers. Stems up to 20 m, glabrous or covered with simple smutty-white to yellowish trichomes. Leaves alternate, simple, paired with a tendril. Leaves sessile (*C. sessilifolia* var. *sessilifolia*), subsessile to distinctly petiolate. Petioles up to 16.5 cm. Petioles glabrous or covered with simple trichomes. Leaves 0.7–20 × 1.1–23 cm, reniform, cordate to deeply palmately 3- to 7-lobate, sometimes lobulate. Lobes triangulate, ovate, elliptical to linear. Margin entire to more or less densely serrate, dentate. Teeth inconspicuous or colored. Leaf apex obtuse, acute to acuminate. Upper leaf surface with clear or whitish pustules, sometimes with trichomes emerging from the lamina or from pustules. Nerves glabrous or with simple trichomes. Lower leaf surface paler than upper surface, glabrous or with simple trichomes. Probracts caducous or persistent, ovate, up to 4.5 mm long. Lower surface keeled or bulging outwards, often with extraxillary glands. Tendrils simple or unequally bifid. Flowers and inflorescences emerging from leaf axils. Male flowers solitary, fascicled or in up to 20-flowered racemes. If solitary flowers and racemes are developed, then solitary flowers occurring before the racemes (within the plant and per node). Common peduncle of raceme 0.5–10 cm, pedicel of flowers in racemes 0.3–1.8 cm, glabrous or with indumentum as on stem but often less dense. Bracts ovate, up to 4 mm long or missing. Pedicel of solitary flowers 0.2–8.5 cm, glabrous or with simple trichomes. Perianth tube glabrous or more or less densely covered with

trichomes. Calyx connate, campanulate, rarely cupulate or urceolate, glabrous, puberulous or with long, simple trichomes. Calyx lobes 0.5–15 mm, triangulate, lineal or subulate; reflexed, spreading to erect. Corolla connate, campanulate, urn-shaped or tubular, 0.7–6.2 cm long; white, dull yellow to orange, salmon; lobes 0.3–4.7 cm, inside densely covered with multicellular trichomes, of which some end with a glandular endcell. Filament column (greenish-)white or orange, anther head pale yellowish green to orange, pollen sacs S-shaped. Female flowers solitary, in pairs or in racemes. Common peduncle 0.3–2.1 cm, glabrous or puberulous. Pedicel of flowers in racemes 0.3–1 cm, glabrous or with simple trichomes, pedicels of solitary flowers 0.7–5 cm, glabrous or puberulous. Calyx and corolla as in males but with hypogynous ovary. Calyx in few cases urn-shaped. Style columnar, greenish yellow, yellow, or orange. Stigmas bulging or 2-lobed, greenish yellow, yellow, or orange. Staminodes 3, attached to the perianth, white (also yellowish or orange?), anthers reduced. Ovary glabrous or with simple, short to long trichomes that then appear articulate when dry. Fruits 1.8–30 × 1.4–5 cm, globose, ovoid, elliptical, or cylindrical; glabrous or with sparse trichomes. Unripe fruits glaucous green to green, sometimes with white, white-and-green or rarely green longitudinal mottling. Ripe fruits orange-red to scarlet red; unicolored or rarely with white to yellowish longitudinal mottling. Seeds enclosed in a hyaline hull, 4.5–7 × 2–3.5 × 1–1.5 mm (L/W/H), symmetrically or asymmetrically obovate, apex round, base narrowed, obtuse, round or square-edged. Face flat to lenticular. Seed surface, depending on the extraction mode, rugulose or filamentose.

1. *Coccinia abyssinica* (Lam.) Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 536. 1881.

Bryonia abyssinica Lam., Encycl. 1(2): 497. 1785.

Type: Cultivated. Unknown, from seeds sent by Bruce (Jeffrey 1962) from Ethiopia, cultivated in Paris Royal Botanical Garden, male, fl, *Anon. in herb. J.-B. Lamarck s.n.* (Holotype: P-LAM! [P00307815, digital image: P-LAM]).

Bryonia macrophylla Ser. in DC., Prodr. 3: 308. 1828.

Type: without location [probably Ethiopia]. Male and female, fl, 1815, *Anon. in coll. [E.]Thibaud s.n.* (Holotype: G-DC!).

Cucumis striatus A.Rich., Tent. Fl. Abyss. 1: 295. 1847.

Type: Ethiopia. [Tigray]: Mt Sholada near Adwa, fr, [Aug], *R. Quartin-Dillon s.n.* (Lectotype, designated here: P! [from “Sholada”]).

Type: Ethiopia. [Tigray]: [Mt] Selleuda [Mt Sholada near Adwa], fr, *R. Quartin-Dillon s.n.* (Syntype: P! [P05621224, digital image: P]).

Cucurbita exanthematica Fenzl ex A.Rich., Tent. Fl. Abyss. 1: 296. 1847.

Type: Ethiopia. Without detailed location, female, fl, *G.H.W. Schimper 1418* (Lectotype, designated here: W! [digital image: WU]; isoelectotypes: BM!, G!, P! [P05621261, digital image: P], TUB! [TUB004727], non TUB-004726!).

Coccinia diversifolia Naudin ex C.Huber, Cat. Print. 1864: 6. 1864. *Cephalandra diversifolia* (Naudin ex C.Huber) Naudin, Ann. Sci. Nat. Bot. ser. 5, 5: 19. 1866.

Type: Cultivated. From seeds sent by Schimper from Ethiopia, cultivated in Paris Botanical Garden and Huber’s Garden in Olbia [Hyères, France], *C.V. Naudin s.n.*

(Lectotype, designated by Jeffrey (1962: 347): P! [P06745719, digital image: P]; isoelectotypes: P [P06745720, digital image: P], G-DC (2)!, K).

Coccinia diversifolia var. *glabrescens* Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 537. 1881. Type: Ethiopia. Chaqou-Choad, 2000 m, in thicket, male, fl, 21 Jul 1852, *G.H.W. Schimper* 250 (Lectotype, designated here: P! [sheet with descriptive text]; isoelectotypes: P (2)!).

Description. Perennial climber. Stems up to 5 m, covered with more or less dense, articulate, dirty-white to yellowish trichomes, rarely glabrous. Petioles 1.5–14 cm, at least on nerves more or less densely covered with articulate trichomes, rarely glabrous. Leaves 7.5–12 × 6.5–12 cm, often cordate to profoundly 3- or 5-lobate. If lobate then central lobe dominating, over-all shape rather (long) cordate (Fig. 4a). Lobes triangulate, ovate to elliptical. Margin more or less densely serrate, dentate. Teeth rarely (if so then small) pale brownish colored in living state or blackish when dried. Leaf apex acute, or if leaf lobate then central lobe acute to long acuminate. Upper leaf surface with clear or whitish pustules, sometimes with some trichomes. Lower leaf surface with soft trichomes articulate appearing when dry or sparsely with stiff narrowly conical trichomes, which can appear warty when short or broken off. Probracts up to 3 mm long. Tendrils simple. Male flowers solitary or in long-pedicelled few-flowered racemes. Pedicel with indumentum as on stem. Common peduncle of raceme 2.5–10 cm, pedicel of flowers in racemes up to 1.5 cm, indumentum as on stem or less dense. Bracts up to 1.7 mm long or missing. Solitary flowers with up to 5 cm long pedicel with trichomes as on stem. Perianth tube more or less densely covered with articulate trichomes. Calyx lobes 2–4 mm, lineal-subulate, upright. Corolla c. 1.4 cm long, yellow to slightly orange, darker on the lobes, lobes up to 5 mm. Filament column white, anther head pale yellowish green, pollen sacs yellow. Female flowers solitary. Pedicel up to 3.5 cm long, indumentum as on stem to glabrous. Style not seen. Stigma shape not seen, yellow. Staminodes not seen. Ovary with long trichomes, often appearing articulate when dry. Hypanthium more or less densely covered with articulate trichomes, calyx lobes and corolla as in males. Fruits 5.5–6 × 3.5–4 cm, short elliptical, glabrous, orange-red sometimes with yellow longitudinal mottling. Seeds 5–6 × 3 × 1.5 mm (L/W/H), slightly asymmetrically obovate, face flat (Fig. 14b).

Phenology. Flowering time: June–October.

Distribution. Fig. 20. Ethiopia (Amhara, Oromia, Southern Nations, Nationalities and People's Region, Tigray). Elevation 1300–2800 m. On limestone, sandstone, black soil, chromic nitisol (Mengesha et al. 2012), loam, on deep to shallow soil. Along lake shores among *Typha* sp., in *Podocarpus-Celtis* forest (clearings) and degraded forms of these, evergreen shrubs (e.g., *Euclea* sp.).

Use. Edibility of fruits is disputed and may differ between wild and cultivated forms (*E. Westphal & J.M.C. Westphal-Stevens* 1951 and 1953). Tuberous roots boiled for food (*T. Ebba* 250), young shoots and leaves are eaten when cooked (Hora 1995). For details see chapter Use, economic potential, and phytochemistry.

Vernacular names. Dawuro: shushe, ushushe (Hora 1995); Galinya [Oromo]: anchote (Getahun 1974b); Kefinya [Kaffa]: ajjo (Hora 1995); Tigrinya: wouchich (*G.H.W. Schimper* 1048); Wollamo [Wolleyta]: ušuşe (*W. Kuls* 681). The Kefinya name is not exclusive for *C. abyssinica* but also used for another crop, *Plectranthus edulis* (Vatke) Agnew.




Figure 20. Distribution map of *C. abyssinica* (pale yellow dots; based on 23 collections) and *C. megarrhiza* (blue dots; based on 28 collections). For Ethiopia the borders of the regions are given.

Remarks. The occurrence of monoecy has been reported by *W.J.J.O. de Wilde et al.* 7805, but the seen specimens contained male flowers only. If both sexes are found on the same individual, this is likely to be a case of leaky dioecy (see also section on Chromosomes and sex determination).

Taxonomic remarks. The *C. abyssinica* specimen in the Lamarck herbarium must be the holotype, since there is only one specimen of *Coccinia abyssinica* in the herbarium of Lamarck in Paris and none in the herbarium of Sonnerat, which he has seen, too. The specimen in the Linnaean herbarium was not annotated with a corresponding name.

Cucurbita exanthematica Fenzl ex A.Rich. is commonly recognized as a synonym of *C. grandis* with a K.G.T. Kotschy collection as type. However, the label on the

Kotschy 308 specimens merely state the species name, the locality, and “frutices scandens” (= climbing on shrubs; W. Greuter – pers. comm.), which cannot be regarded as a diagnostic feature. The label is printed and therefore effectively published but not validly so. Valid publication of that name was effected by Achille Richard (1847), but he chose a different specimen (*G.H.W. Schimper* 1418), which belongs to *C. abyssinica*. The *Schimper* 1418 specimens bear printed labels on which Fenzl designated a variety of his invalid name with the phrase “var. foliis superioribus integris (non lobatis)”. The phrase, however, is also not a validation since the species to which this variety is supposed to belong, is not validly published either (Art. 41.3a and b ICN). Naudin (1859) suggested that Eduard Fenzl mixed-up some specimens. He accepted the *Kotschy* 308 specimen as a synonym of his *C. schimperi* and recognized the similarity of the Schimper specimen to Lamarck’s *Bryonia* (*Coccinia*) *abyssinica* and *Cucumis striatus*.

The identity of *Cucumis striatus* A.Rich. is not obvious. There are two original specimens with this name in P herbarium: one from Selleuda (P05621224) and the other one from Sholada, both names for the same mountain near the city of Adwa. The P05621224 specimen consists of a ripe fruit, a drawing of the fruit, and a tiny fragment of a leaf. Cogniaux identified this specimen as *C. adoensis*. However, the fruit is ovoid, which would be unusual for that species in which fruits are long ovoid to short cylindrical and often have a sterile apex (“beak”). Since there are no seeds, which would help to clear this problem up easily, the fruit shape is the only usable character. The leaf fragment might be *C. adoensis* but it is too small to be certain, and it is loose so it might also be debris from another specimen. The other original specimen (with a number “26” from “Sholada”) contains much leaf material and fruits. The fruits are darker than in the first type specimen. The indumentum of the lower leaf lamina matches certain *C. abyssinica* collections, as does the leaf shape (cf. *G. Negri* 703, *G.H.W. Schimper* 250) although they are not very typical. This specimen is not close to *C. adoensis*, therefore the present author chose it to be the lectotype and to synonymize the name *Cucumis striatus* with *C. abyssinica*.

Specimens examined. (Selection, in total: 57) Ethiopia. Amhara: Sankaberr [vicinity of Reb river] and Begemder [highland], *G.H.W. Schimper* 1446 (E [E00303229], S [S08-12052], S [S08-12057], W, Z (3)). Oromia: 32 km from Addis Abeba on road to Debre Zeit [Debre Zeyit], *E. Westphal & J.M.C. Westphal-Stevens* 1951 (BR [BR0000008914613], EA, MO, PRE, WAG [WAG0225550], WAG [WAG0225551], WAG [WAG0225552]) & 1953 (MO, WAG [WAG0225546], WAG [WAG0225547]). SNNPR: Bonga, near Roman Catholic Mission, *W.J.J.O. de Wilde & B.E.E. de Wilde-Duyffes* 7805 (MO, WAG [WAG0225537], WAG [WAG0225538], WAG [WAG0225539]). Tigray: 18 km along road from Adu Abun to Axum, 14°09'N, 38°49'E, *J.J.F.E. de Wilde* 7059 (M, WAG [WAG0225544], WAG [WAG0225545]).

2a. *Coccinia adoensis* (Hochst. ex A.Rich.) Cogn. var. *adoensis*

Momordica adoensis Hochst. ex A.Rich., Tent. Fl. Abyss. 1: 293. 1847. *Coccinia adoensis* (Hochst. ex A.Rich.) Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 538. 1881.

Type: Ethiopia. [Tigray]: Adwa, near church “Eta Mariam”, mixed specimens, male and female, fl, fr, 5 Jun 1837, *G.H.W. Schimper Iter Abyss. sect. 1 no. 166* (Lectotype, designated here: P! [P00346261, digital image: JPS, P]; isolectotypes: BM! [BM001010004], BR [BR0000008351036], BR! [BR0000008351039, digital image: BR, JPS], BR! [BR0000008886781, digital image: BR, JPS], G! [G00301597], G! [G00301598], G-DC!, HBG! [HBG506429, digital image: JPS], K! [K000542642, digital image: JPS, K], K! [K000542643, digital image: JPS, K], K! [K000542644, digital image: JPS, K], L! [L0057303, digital image: JPS, L], LG! [LG0000090028250; digital image: JPS], M! [M0105779, digital image: JPS], M! [M0105780, digital image: JPS], P! [P00346261, digital image: JPS, P], P! [P00346262, digital image: JPS, P], S! [S10-19076, digital image: S], TUB [TUB004719, digital image: JPS], TUB [TUB004720, digital image: JPS], W! [W0011091, digital image: WU], W! [digital image: WU]).

Type: Ethiopia. [Tigray]: near Adwa, in thicket, *R. Quartin-Dillon s.n.* (Syntype: P!).

Type: Ethiopia. [Amhara?]: Ouodgerate Province, *A. Petit no.?* (Syntype: P?).

Bryonia convolvuloides A.Rich., Tent. Fl. Abyss. 1: 289. 1847.

Type: Eritrea/Ethiopia. [Gash-barka/Tigray]: Chiré [borderland between both regions, between Tekezé river and Mareb/Gash river], male, fl, *R. Quartin-Dillon & A. Petit s.n.* (Lectotype, designated here: P! [P05621227, digital image: P]; isolectotype: P! [P05621226, digital image: P]).

Type: Ethiopia. No location, male, 1844, *R. Quartin-Dillon & A. Petit s.n.* (Syntype: P! [P05621255, digital image: P]).

Bryonia jatrophiiifolia A.Rich. [sphalm. *Bryonia jatrotrophæfolia* A.Rich.], Tent. Fl. Abyss. 1: 289. 1847. *Coccinia jatrophiiifolia* (A.Rich.) Cogn. [sphalm: *Coccinia jatrophæfolia* (A.Rich.) Cogn.] in A.DC. & C.DC., Monogr. Phan. 3: 535. 1881.

Type: Ethiopia. [Tigray]: in valley near Adwa, male, fl, Aug 1839, *R. Quartin-Dillon & A. Petit s.n.* (Lectotype, designated here: P! [P00346263, digital image: JPS, P; K neg. 2990], syntype: P! [P05621222, digital image: P]).

Type: Ethiopia. [Tigray]: near Tchélatchekanné [Djeladjeranné, in Tekezé river valley (Gillett 1972), on label as “Tchessu Hetchequenné”], male, fl, *R. Quartin-Dillon s.n.* (Syntype: P! [P00346260, digital image: JPS, P; K neg. 4467], P! [P05621223, digital image: P]).

Cephalandra pubescens Sond. in Harv. & Sond., Fl. Cap. 2: 493. 1862. *Coccinia pubescens* (Sond.) Eyles, Trans. Roy. Soc. South Africa 5(4): 498. 1916. *Coccinia pubescens* (Sond.) Cogn. ex Harms in Fries, Notizbl. Bot. Gart. Berlin-Dahlem 8: 491. 1923. nom. superfl.

Type: South Africa. [North West]: Magaliesberg, male and female, fl, Dec, *J. Burke 408* (Lectotype, designated here: K! [K000313229, digital image: JPS, K]; isolectotype: BM! [BM000815207], K! [K000313227, digital image: JPS, K], NBG, Z!).

Type: South Africa. North West/Gauteng: at Magalies river, male and female, fl, *C.L.P. Zeyher 588* (Syntype: K! [K000313228, digital image: JPS, K], S! [S08-12187, digital image: S]).

Coccinia hartmanniana Schweinf., Reliq. Kotschy.: 42, t. 27, t. 28. 1868.

Type: drawing in protologue, t. 27 (Lectotype, designated here).

Type: Sudan. Sinnar Province: no detailed location given, 1860, *R. Hartmann s.n.* (Syntype: ?, if B, then destroyed).

Type: South Sudan. At White Nile in the region of the Tschier people [a nilotic tribe also called: Chir, Kir, Mandari, Mondari, Mundari, Shir], [possibly the area between Tombe and Mongalla in Central Equatoria state], 1861, *W. von Harnier s.n.* (Syntype: B destroyed, ?).

Type: South Sudan. [al-Qadarif]: Gallabat at Gendua [river], Jun 1861, *H. Steudner 843* (Syntype: ?, if B, then destroyed).

Type: Ethiopia. [Amhara]: near Matamma at border to Sudan, after beginning of Jun 1865, *G.A. Schweinfurth Flora of Gallabat 62* (Syntype: ?, if B, then destroyed).

Coccinia rigida Cogn., Bot. Jahrb. Syst. 21: 210. 1895.

Type: Tanzania. [Tabora]: Ugunda, near Gonda [Igonda], on ground in wet corn fields, *R. Böhm 176* (Holotype: B destroyed, lectotype, designated here: BR! [BR0000008886804, digital image: BR, JPS]).

Coccinia djurensis Schweinf. et Gilg, Bot. Jahrb. Syst. 34: 357. 1904.

Type: South Sudan. [West Bahr al-Ghazal or Warab]: Seriba Ghattas, male, fl, 24 May 1869 or Jun 1869, *G.A. Schweinfurth 1878* (Lectotype, designated here: P!; isolecotypes: B [destroyed], E! [E00303230], G!, K! [K000542640, digital image: JPS, K], PRE [PRE0592944-0, digital image: JPS], S! [S08-12060, digital image: S], Z! [Z-000073403, digital image: Z]).

Type: South Sudan. [West Bahr al-Ghazal or Warab]: Seriba Ghattas, male, fl, 24 May 1869, *G.A. Schweinfurth 1867* (Syntype: if B, then destroyed).

Type: South Sudan. [West Bahr al-Ghazal]: Djur realm, Seriba Agad Wau [Waw], fr, May, *G.A. Schweinfurth 1688* (Syntype: B [destroyed], K! [K000542641, digital image: JPS, K], L!).

Coccinia princeae Gilg, Bot. Jahrb. Syst. 34: 358. 1904.

Type: Tanzania. Iringa: Uhehe highland, no detailed location given, fr, *M. von Prince s.n.* (Holotype: B, destroyed).

Type: Tanzania. Morogoro: Uluguru Mts, NW slopes, male, fl, 19 Jan 1933, *H.J.E. Schlieben 3271* (Neotype, designated here: B!; isoneotypes: G!, HBG!, M! [M-0210964], P! [P05621242, digital image: P], S! [S08-12183]).

Coccinia parvifolia Cogn. in Schinz, Vierteljahresschr. Naturforsch. Ges. Zürich 52: 433. 1907.

Type: South Africa. Limpopo: [Mopani District], [Leydsdorp area], Mt Marovounge [Mt Marovougne], male, fl, May 1904, *H.A. Junod 2491* (Holotype: Z! [Z-000004444, digital image: Z], isotype: BR! [BR0000008886811, digital image: BR, JPS]).

Coccinia homblei Cogn., Bull. Jard. Bot. État Bruxelles 5: 114. 1916.

Type: D. R. Congo. Katanga: Lualaba region, Kapanda confluence plains, male, fl, Dec 1912, *H. Homblé 992* (Lectotype, designated here: BR! [BR0000008887146, digital image: BR, JPS]; isolecotype: BR! [BR0000008887474, digital image: BR, JPS]).

Type: D. R. Congo. Katanga: Kapiri valley, fr, Feb 1919, *H. Homblé 1198* (Syntype: BR! [BR0000008886927, digital image: BR], BR! [BR0000008886859, digital image: BR, JPS]).

Type: D. R. Congo. Katanga: Kapiri valley, male, fl, Feb 1919, *H. Homblé 1199* (Syntype: BR! [BR0000008887177, digital image: BR, JPS], BR! [BR0000008887504, digital image: BR, JPS], BR! [BR0000008886842, digital image: BR, JPS, K neg. 4875]).

Coccinia subspicata Cogn., Bull. Jard. Bot. État Bruxelles 5: 115. 1916.

Type: D. R. Congo. Katanga: Lualaba region, Kapanda confluence plains, male, fl, Dec 1912, *H. Homblé 992a* (Holotype: BR! [BR0000008887528, digital image: BR, JPS, K neg. 4876]).

Coccinia roseiflora Suess., Proc. Trans. Rhodes. Sci. Ass. 43: 134. 1951.

Type: Zimbabwe. [Mashonaland East]: Marandellas [Marondera], Cave Tatooma, 31 Nov 1941, *G. Dehn 188A* (Holotype: M! [M0105778, digital image: JPS]).

Description. Perennial climber or creeper. Stems up to 6 m, glabrous to densely covered with trichomes. Indumentum whitish to beige. Trichomes < 0.5 mm, sometimes curved (Fig. 21). Petioles 0.6–3.5 cm, glabrous, pubescent to tomentose with indumentum like on stem. Leaves 4.2–13.5 × 4.7–16 cm, cordate, shallowly to deeply 3- or 5-lobate, rarely elliptical with hastate base. If profoundly to deeply lobate, then lobes often (slightly) pointing towards apex (as in *C. keayana*; Fig. 32). Lobes triangulate, lanceolate, lineal, elliptical to obovate. Leaf margin entire to serrate (especially on lateral sides) to lobulate. Tips of lobes retuse, obtuse to acute, often with apical tooth. Upper leaf surface clear to white pustulate, rarely with whitish trichomes. Lower leaf surface soft pubescent with short, whitish trichomes, rarely glabrous, often with black to dark brown glands between nerves towards the base, rarely also along main nerves. Probracts missing or obovoid up to 2.5 mm long. Tendrils simple, rarely bifid. Male flowers in few to many-flowered racemes, often accompanied by one solitary flower. Common peduncle 1–11 cm, glabrous or with short trichomes. Pedicel of racemous flower 0.4–1.3 cm, indumentum like peduncle, pedicel of solitary flower 1.5–4 cm glabrous or with short, white trichomes. Bracts missing (caducous?) or up to 2 mm. Perianth tube glabrous or with short (< 0.5 mm), white trichomes. Calyx lobes 1.2–2.5(–3) mm, linear to broadly triangular, adpressed to spreading, apex obtuse to acute. Corolla 0.9–1.6 cm, yellow, salmon-pink, orange, maroon, veins sometimes purplish-brown, lobes 2–5 mm. Filament column whitish to orange, anthers whitish?, yellowish to orange, pollen sacs yellow to orange. Female flowers solitary. Pedicel 0.8–3.5 cm, often glabrous or with short, white trichomes. Hypanthium glabrous or with short (< 0.5 mm), white trichomes. Calyx lobes and corolla like in male flowers. Ovary glabrous. Style and stigmas not seen. Fruits 3–7 × 1–1.5 cm, ovoid, oblong to shortly cylindrical, while ripening (usually?) with dark green longitudinal mottling, ripe orange-red to red, often with sterile tip (“beaked”). Seeds 4–6 × 3–4 × 1–1.7 mm (L/W/H), symmetrically obovate, face lenticular (Fig. 14a, 21).

Phenology. Flowering time: January–May, August–December.

Distribution. Fig. 22. Angola? (likely in the South and East since the species occurs close-by), Botswana (North-West District), Burundi?, Cameroon (Extreme North), N Central African Republic, S Chad, Democratic Republic of Congo (Katanga, along the Western Rift), Eritrea (Gash-Barka, likely wider distributed as relicts



Figure 21. Male inflorescence and leaf of *C. adoensis*; picture taken from lectotype (G.H.W. Schimper 166 (P00346261)). Note the short bent trichomes, which are a good indicator for this species (but glabrous collections or other kinds of trichomes may occur in this species, too).

in the highlands), Ethiopia (except the dry southeast), Kenya (in the west and central highlands), Malawi, Mozambique, N Namibia, Nigeria (only known from Adamawa State, but likely more widely distributed), Rwanda (Eastern Province, maybe wider distributed), South Africa (Gauteng, KwaZulu-Natal, Limpopo, Mpumalanga, E North West, E Free State), South Sudan, Sudan (West Darfur, maybe also in woodland relict sites in other provinces), Swaziland, Tanzania, Uganda (Northern Region), Zambia, Zimbabwe. Elevation 130–3450 m. On sandy and silty soils, clay loam, loam, laterite, syenite soils, dolerite soil, dolomite soil, limestone. *Hyparrhenia cymbaria* savanna; *Crossopteryx* tree savanna; *Acacia-Combretum-Stereospermum-Cussonia* woodland; *Pseudoprosopis fischeri* woodland, *Anogeissus leiocarpus* woodland, *Brachystegia* woodland, sourveld grassland, *Melhania rehmannii-Enneapogon scoparius* mixed bushveld (Siebert et al. 2010).

Use. Roots are boiled and drunk for fever (J.C. Lovett & C.J. Kayombo 3434). The potato-like tubers are eaten (F.W. Andrews 1310), also raw (T. Scudder 56). The greens are used as spinach, among others by the Venda (N.J. van Warmelo s.n. Mar 1960, J. Gerstner 5838) and also eaten by the Luo (Johns and Kokwaro 1991). Ripe fruits are edible (J.C. Lovett et al. 3842, J. Pawek 11008, T. Scudder 56). According to de Boer et




Figure 22. Distribution map of *C. adoensis* var. *adoensis* (based on 311 collections).

al. (2005), the Pare people in Tanzania use an infusion of leaves and stems for abortions, uterus cleansing, and against chickenpox. *Coccinia adoensis* is quite variable and so there are likely many varieties, of which some might not be edible. Rehm et al. (1957) reports the cytotoxic cucurbitacin B and traces of cucurbitacin D in immature fruits, but edible ripe fruits are known from South Africa. Gradé et al. (2009) report that the usability or toxicity of the tuberous roots is disputed, suggesting chemical variability.

Vernacular names. Bokora tribe [Karamojong?]: edaldalakisín (Gradé et al. 2009); Kiluo: mutkuru (Johns and Kokwaro 1991), nyatunduguwoqe (J.G.B. Newbould 5745); Kipare: mlushi (de Boer et al. 2005); Kisafwa: tandandala (J.C. Lovett & C.J. Kayombo 3434), ndandala (J.C. Lovett & C.J. Kayombo 3773); Kisagara [Kisagala]: lutetere (F. Haerdi 563/0); Nhungoé [Cinyungwe]: mugwingwi (L. Macuácuá 1441); Sotho [most likely Northern Sotho]: sephu (J. Gerstner 5838); Tigrinya: entatakh (fide G.H.W. Schimper (Schweinfurth 1893)), entota (fide A. Richard (Schweinfurth 1893)); Tshivenda: tshiphu (N.J. van Warmelo s.n. Mar 1960); Zande: bawiriokoro (F.W. Andrews 1601).

Remarks. *Coccinia adoensis* is widespread and morphologically variable. Some populations or local forms appear to be distinct, but there are intermediate individuals or similar-looking collections from different parts of the overall distribution range. In East Africa (C Tanzania, Malawi), one can find forms linking to *C. aurantiaca*, which is treated here as variety of *C. adoensis*, and to *C. senensis*. The latter forms have a similar plastid haplotype with *C. senensis*, but lack a specific deletion in the *trnS*^{GCU}–*trnG*^{UCC} intergenic spacer. These forms from central and southern Tanzania are discussed here under the name *C. adoensis* var. *jeffreya*, while another form from Kenya remains in var. *adoensis*. The non-monophyly in the plastid tree (Holstein and Renner 2011b) makes *C. adoensis* even more peculiar. The scenario given in the chapter Evolution and phylogeny might explain this pattern, but without phylogeographic analysis and crossing experiments, this will remain speculative.

Some specimens from the Kingupira area (Lindi, Tanzania; K. Vollesen 3182, 3212, 3384, 4320) have an unusual morphology by having veins that run along the leaf margin, which is unique in *Coccinia*. Except for this character, they match *Coccinia adoensis* var. *adoensis* well (sympetalous, obovate probracts). They strongly resemble *Eureiandra* species in vegetative characters, but not in generative traits.

Taxonomic remarks. The lectotypification of *Momordica adoensis* by Meeuse (1962) is not effected, as he did not specify which specimen was supposed to be the lectotype. However, the present author follows his suggestion and chose among the two Schimper 166 specimens from P.

The types of *Bryonia jatrophaefolia* are not too obvious as such. The protologue states “Tchélathekanné”, but Paris Herbarium holds two Quartin-Dillon and Petit specimens (P00346260 and a non-barcoded one) with a location “Tchessu Heckequenné”. Although the spelling has some similarities, they are quite different. On the other hand, the Ge’ez letter sat [S] and läwe [L] look similar and might have been mistranscribed if the name was written down in Ge’ez script. Additionally, the two specimens bear the species name in Richard’s handwriting (C. Bräuchler, pers. comm.). Gillett (1972) notes that Achille Richard, the author of the botanical treatment of Quartin-Dillon’s and Petit’s journey, wrote consistently “Tchelatchekanné”, although Schimper calls it “Djeladjerranné”. However, there is a locality with that name [Tchelatchekenneh at c. 13°43’N, 38°22’E] in Johnston’s atlas (Johnston 1861), to which Gillett (1972) refers.

The G.A. Schweinfurth 1668 specimen in L is not obviously a type specimen as it lacks the original label. However, the Herb. D’Alleizette label mentions “*Coccinia djurensis*”, and the location is the same as the K duplicate with the original label. Ad-

ditionally, the specimen is a fruiting female, like the specimen in K. Hence, d'Alleizette must have obtained a duplicate of the type.

The placement of *C. hartmanniana* as a synonym of *C. adoensis* is done with a high level of confidence although no type specimens were found. The protologue contains drawings showing lenticular seeds and short calyx lobes, which match well other collections of the *C. adoensis* complex. According to Ascherson (in Schweinfurth 1867), von Harnier's collections consisted of two duplicates, one of them in B. Other duplicates of von Harnier are in BM and K but not this collection apparently. Since no type specimen was seen, despite extensive search, a drawing from the protologue was chosen as lectotype.

For *C. princeae*, a neotype was selected because the holotype was destroyed. The leaves of the chosen specimen, *H.J.E. Schlieben* 3271, match the description well, and the specimens have been identified as *C. princeae* when the original material was still existing. The specimens differ in the generative characters (fruiting in the holotype, male flowers in the neotype), but Gilg referred strongly to the distinctive leaves, so the neotype appears to be a good match.

The holotype of *C. roseiflora* is the drawing in M, which has the number 188a. The protologue states: “descriptio sec. tabulam cl. Dehniae” (described following/based on the illustration of Ms. Dehn). There are (at least) two specimens with the number 188 (K, SRGH), which partially might have served as basis for the drawing, but they are not types. The drawing contains all necessary characters to synonymize it with certainty with *C. adoensis*: seed shape, fruit maculation and calyx lobe morphology.

Specimens examined. (Selection, in total: 483) Botswana. North-West District: Ngamiland, Motantanyane, *H.H. Curson* 784 (M). Central African Republic. Nana-Grébizi: Gribingui (Ft-Crampel) [=Kaga-Bandoro], *A.J.B. Chevalier* 6325 (P [P05621208]). D. R. Congo. Orientale: Faradje (Kibali-Ituri), *J. Lebrun* 3406 (BR, WAG [WAG0225534]). Eritrea. Gash-Barka: Seraé [Seraè, a former province], in Tukul region, *A. de Benedictis* 519 (FT). Ethiopia. Tigray: Edaga Sciaba, *E. Chiovenda* 581 (FT). Malawi. Northern Region: Karonga district, 17 mls [27.2 km] N of Chilumba, *J. Pawek* 11008 (DSM, MO, PRE, WAG [WAG0234129]). Mozambique. Zambézia: Massingire [Morrumbala district], M'bôbo [river at Morrumbala], on road to Mopeia, *A.R. Torre* 5337 (M). Nigeria. Adamawa: 10 miles [16 km] from Mubi to Toyola, *P. Wit et al.* 1797 (BR, MO (2), P [P05621253], WAG [WAG0041392], WAG [WAG0041393]). Rwanda. Eastern Province: Kibungo prefecture, Rusumo [Ruzumo Falls], savanna park, on opposite slope of A.I.D.R. [Association internationale de développement rural] camp, *J. Lambinon* 74/1568 (M, MO, WAG [WAG0225516]). South Africa. Gauteng: [City of Johannesburg], N of Eikenhof, Johannesburg, Walkerville rd., *L.E. Davidson* 3781 (B, M). Sudan. West Darfur: Zalingei, *G.E. Wickens* 1800 (K). Tanzania. Lindi: Selous Game Reserve, Kingupira, 8°28'S, 38°33'E, [38°34'E], *K. Vollesen* MRC 3384 (DSM, EA, WAG [WAG0234135]). Ruvuma: Songea, *E. Milne-Redhead & P. Taylor* 7876 (EA); *ibid.*, *E. Milne-Redhead & P. Taylor* 7950 (B, EA, K, P [P05621243], S [S08-11819]). Zambia. Lusaka: c. 10 km S of

Chilanga and 23 km S of Lusaka, c. 1 km N of Kafue Road near Chilanga Cement housing; Shimabala Cave, 15°39'01"S, 28°14'15"E, *D.K. Harder & M.G. Bingham* 2584 (K, MO). Zimbabwe. Matabeleland North: Matetsi Safari Area Headquarters House no. 2, *P. Gonde* 256 (E, G, MO, P [P05621269]).

2b. *Coccinia adoensis* var. *aurantiaca* (C.Jeffrey) Holstein, stat. nov.

urn:lsid:ipni.org:names:77148917-1

Coccinia aurantiaca C.Jeffrey, Kew Bull. 17: 169. 1963.

Type: Tanzania. Dodoma: Kondoa District, Great North road, 15 miles S of Kondoa, 1310 m, fl, fr, 19 Jan 1962, *R. Polhill & S. Paulo* 1221 (Holotype: K! [K000313236, digital image: JPS, K], isotypes: B! [B 10 0154922, digital image: B, JPS], B! [B 10 0154923, digital image: B, JPS], BR! [BR0000008887252, digital image: BR, JPS], EA (2)!, PRE! [PRE0592951-1, digital image: JPS], PRE! [PRE0592951-2, digital image: JPS]).

Type: Tanzania. Dodoma: 1 ml. [1.6 km] S of Dodoma, Imagi hill, *R. Polhill & S. Paulo* 1274 (Paratypes: B! [B 10 0154921, digital image: B, JPS], BR!, EA!, K (3)!, P! p.p., PRE!, S! [S08-11841, digital image: S]).

Type: Tanzania. Mwanza: Mwanza, Ilemera, Butimba, *R.E.S. Tanner* 1902 (Paratypes: BR!, EA!, K (2)!).

Type: Tanzania. Mwanza: Mbarika [chiefdom], Buzomo, *R.E.S. Tanner* 1068 (Paratypes: BR!, COI (2)!, EA!, K!, NY!).

Type: Tanzania. Mwanza: Mwanza, *R.E.S. Tanner* 646 (Paratype: K!).

Type: Tanzania. Shinyanga: near Shinyanga, *R.D. Bax* 57 (Paratypes: K (2)!).

Type: Tanzania. Shinyanga: hills near Shinyanga, *B.D. Burt* 2517 (Paratype: K!).

Type: Tanzania. Shinyanga: Shinyanga, *H. Koritschner* 1823 (Paratypes: EA (2)!, K!).

Description. Perennial climber. Stems up to 10 m, almost tomentose with short (< 0.5 mm), stiff, whitish trichomes. Petiole 1.5–3.5 cm, indumentum similar to stems. Leaves 5.2–12.5 × 6.4–14.5 cm, cordate, shallowly to profoundly 3- or 5-lobate. Lobes triangulate, ovate, elliptical to obovate. Margin serrate to lobulate. Apex obtuse, rarely acute, with final tooth. Upper leaf surface white-pustulate sometimes more or less pubescent with short, whitish trichomes. Lower leaf surface usually densely covered with curved or short, straight trichomes on nerves. Probracts up to 1.5 mm, often caducous. Tendrils simple. Male flowers solitary or in racemes. Common peduncle 0.4–3 cm, pedicels in racemes up to 0.5 cm, pedicel of solitary flowers 1–2 cm, indumentum as on stem. Bracts up to 1.5 mm, persisting. Perianth tube densely covered with short (< 0.5 mm) trichomes. Calyx lobes 1–3 mm, narrow triangular to dentate, spreading. Corolla 1.6–2.4 cm long, pale yellow-brown to orange, rarely? yellow, with green to orange venation, lobes 0.6–1 cm. Filament column, anther head, and pollen sacs more or less pale orange, rarely yellowish? (Fig.




Figure 23. Distribution map of *C. adoensis* var. *aurantiaca* (pale yellow triangles; based on 18 collections), *C. adoensis* var. *jeffreyana* (pale yellow dots; based on 18 collections), *C. pwaniensis* (blue triangles; based on 11 collections, including a natural hybrid), and *C. senensis* (blue dots; based on 30 collections). For Tanzania the borders of the regions are given.

10b). Female flowers solitary, pedicel 1.2–4 cm, indumentum as on stems. Hypanthium densely covered with short (< 0.5 mm) trichomes, calyx lobes, and corolla as in male flowers. Style shape not seen, green. Stigma shape not seen, yellow to orange. Ovary with short trichomes. Unripe fruits pale green with irregular lighter spots and

dark green longitudinal lines. Fruits 5–9 × 1.5–3.5 cm, long ovoid, apex sometimes beaked, when ripe orange-red. Seeds 6–6.5 × 3.5–4 × 1.5–1.7 mm [L/W/H], slightly asymmetrically obovate, face flatly lenticular.

Phenology. Flowering time: January, March, July, October, December.

Distribution. Fig. 23. Tanzania (Dodoma, Iringa, Manyara, Morogoro, Mwanza, Shinyanga). Elevation 600–1200 m. Red sandy soil, red clay, granite. White clay. Gray sand. Brown sandy loam. Dry Miombo woodland, *Acacia tanganyikensis*-*A. tortilis* subsp. *spirocarpa*-*Adansonia digitata*-*Maerua crassifolia*-*Balanites aegyptiaca* woodland, long grass savannas, dry (*Commiphora*-*Acacia*) bushland, thickets (e.g., *Combretum* thickets), among rocks on hills.

Use. Leaves are boiled and eaten (*J.L. Newman* 62). Fruits edible when ripe and dry (*E.S. Macha* 600).

Vernacular names. Sandawe language: koba (*J.L. Newman* 62).

Remarks. The status of this taxon as species is unclear, therefore it is treated as a variety of the polymorphic *C. adoensis*. *Coccinia adoensis* var. *aurantiaca* specimens as listed here are usually more densely covered with trichomes than *C. adoensis* var. *adoensis*. Jeffrey segregated this species from the polymorphic *C. adoensis* because of the non-beaked fruits and flat seeds with a hyaline girdle. The beak is a sterile part of the ovary with variable length, but it does not occur in all populations. Two of the paratypes (*R. Polhill* & *S. Paulo* 1274 (BR, P)), which match other *C. adoensis* var. *aurantiaca* collections vegetatively, have a slightly beaked fruit, although most other collections do not. The seeds are also hardly distinct from *C. adoensis*, perhaps somewhat larger. Seeds in *Coccinia* are enclosed in a hyaline aril. Jeffrey only observed the dry collapsed aril, which is not part of the seed, as a “hyaline girdle”. The orange color of the petals, even with purple venation also occurs in individuals of *C. adoensis* var. *adoensis* that have a less dense indumentum. The corolla is thus not a good distinguishing character either. However, this variety seems to occur in a drier part of the range of the overall *C. adoensis* distribution (*Holstein and Renner* 2011b).

Taxonomic remarks. The flowers in the *R. Polhill* & *S. Paulo* 1274 specimen in P do not belong to *Coccinia*. The calyx appears to be *Momordica foetida* Schum. & Thonn. The HEID specimen (HEID779579) of that collection is also mistaken, eventually a mix-up while mounting the specimen. It has a completely different indumentum and a narrow, almost cylindrical perianth tube.

Specimens examined. (Selection, in total: 29) Tanzania. Dodoma: Dodoma–Kondoa road, c. 20 km S of Kondoa, 05°16'31.5"S, 35°53'01.1"E, *N. Holstein et al.* 85 (DSM, M), and 86 (M). Iringa: Iringa Rural District, along road Iringa–Morogoro road and Lukosi River, at bottom of Kitonga Gorge, c. 6 km W of Mahenge village at milepost 253 km from Morogoro, 7°38'S, 36°14'E, [7°34'S, 36°19'E], *C.M. Taylor et al.* 8485 (K, MO); [Ruaha National Park], Msembe [near airfield], *P.J. Greenway* & *K. Kanuri* 14811 (EA (2), K, M). Manyara: Tarangire National Park, road Tarangire camp–Babati, 1 ml. from camp, *H.M. Richards* 24817 (EA, K). Morogoro: Kilosa district, Elphon's Pass, 7°22'S, 36°42'E, *J.C. Lovett* & *T.C.E. Congdon* 2931 (K, MO).

2c. *Coccinia adoensis* var. *jeffreyana* Holstein, var. nov.

urn:lsid:ipni.org:names:77148914-1

Type: Tanzania. Iringa, Mufindi District, Ngwazi, 8°30'S, 35°15'E, 1830 m, female, fl, fr, 25 Feb 1987, *J.C. Lovett 1597* (Holotype: MO!, isotype: EA!).

Diagnosis. This variety has affinities with *C. adoensis* and *C. senensis*. The abaxial side of the petiole and the lower leaf surface bears simple trichomes with long cells, which appear crumpled or articulate when dry. Most of the trichomes, especially on the nodes, exceed 0.8 mm (–1.2 mm), whereas trichomes of *C. adoensis* var. *adoensis* and var. *aurantiaca* are shorter < 0.5(–0.8) mm. The calyx lobe length often exceeds 2 mm (in contrast to other *C. adoensis* varieties), but the lobes are not subulate or narrowly acute as in *C. senensis* but rather linear or if narrowly triangulate, then not with a pointed tip.

Description. Perennial creeper or climber. Stems up to 3 m, more or less densely covered with long (at least on the nodes > 0.8 mm, Figs 3a, 5c) trichomes that appear articulate when dry. Petiole 0.25–3.5 cm, subsessile to distinctly petiolate, with long patent trichomes. Leaves 3.2–10.5 × 2.6–12 cm, shallowly to profoundly 3- or 5-lobate, lobes triangular, ovate to elliptical, margin dentate, slightly serrate, apex acute to obtuse with apical tip. Upper leaf surface glabrous or with few trichomes, hyaline to white pustulate. Lower leaf surface more or less densely covered with articulate trichomes, rarely almost glabrous with white pustules on veins. Probracts up to 3 mm. Tendrils simple. Male flowers in racemes, often accompanied by a single flower or one solitary flower. Common peduncle 5–5.5 cm, with short articulate trichomes. Pedicel of racemous flowers 5–9 mm, with short articulate trichomes. Bracts up to 1 mm, caducous. Pedicel of solitary flower 2.2–7.8 cm, with short articulate trichomes. Hypanthium with short trichomes. Calyx lobes 1–3.5 mm, narrowly triangular but not subulate, erect. Corolla 1.1–1.65 cm, yellow, orange, to dark crimson with darker veins outside, lobes 4–7 mm. Color of filament column pink, anther head orange-yellow to orange, color of pollen sacs not seen. Female flowers solitary. Pedicel 0.6–1.7 cm long, puberulous. Ovary with short to long, articulate trichomes. Fruit size c. 2–6 × c. 1 cm long, elliptical, often with sterile apical tip (“beaked”), glabrous, green with white spots when unripe, red when ripe. Seeds 4–5.5 × 3–3.5 × 1.5 mm (L/W/H), symmetrically obovate, face lenticular (Fig. 14a).

Phenology. Flowering time: January–March, November, December.

Distribution. Fig. 23. Malawi (Northern Region, Southern Region), Tanzania (Dodoma?, Iringa, Mbeya, Morogoro?, Singida), Kenya (southern Rift Valley Province). 1300–2600 m. Soil preferences unknown. With *Dodonea viscosa*; under pines; in *Eucalyptus* plantation, highland grassland, in open woodland with *Combretum* sp., *Grewia* sp., *Strophanthus emenii*, *Acacia tortilis*, *Tapiphyllum obtusifolium*, *Burttia* sp., *Cassia* sp.

Etymology. The epithet was chosen to honor Charles Jeffrey, who worked extensively on the Cucurbitaceae and the flora of East Africa.

Use. Unripe and ripe fruits are reported to be edible (*C.J. Kayombo 296*, *P. Kuchar 22631*), roots taken to make stomach medicine (*P. Kuchar 22631*).

Vernacular names. Kihehe: mtumbulansoka (*W. Carmichael* 171); Kinyaturu: mukunguhi (*P. Kuchar* 22631).

Remarks. Morphologically, this variety closely matches *C. senensis* (with rather short petiolate to subsessile leaves, and a *C. senensis*-like indumentum), but it has the calyx lobes rather of *C. adoensis* var. *adoensis*, with the lobe length being intermediate between *C. senensis* and *C. adoensis* var. *adoensis*. The sequenced specimens do not cluster with most other *C. adoensis* haplotypes from East Africa or southern Africa, and lack the typical deletion of *C. senensis* in the *trnS*^{GCU}–*trnG*^{UCC} intergenic spacer (Holstein and Renner 2011b). A *C. adoensis* var. *adoensis*-like collection (*S.A. Robertson* 1925) also clusters with this variety, but it lacks the long trichomes. Long trichomes also appear in populations of *C. grandiflora* or *C. mackenii* in higher altitudes or in areas with higher precipitation. The collections of this variety are distributed above 1300 m and thus receive higher amounts of rainfall, so the long trichomes could be an adaptation. On the other hand, very similar trichomes regularly occur in *C. senensis*, sometimes short though, but that species does not occur in such high altitudes. As the collections of this variety differ from the “typical” *C. adoensis*, but still belong to *C. adoensis*, they are treated as a new variety.

The collection *R.E. Gereau & C.J. Kayombo* 3582 (K, MO; *C. adoensis* 4 in Fig. 17) is morphologically inseparable from this variety, and the plastid haplotype clusters within East African *C. adoensis*. This collection has a normal-sized corolla, and therefore seems to be fertile, which supports the hypothesis that the var. *jeffreyana* is not reproductively isolated from var. *adoensis*. This is also why the present author refrains from designating it as a paratype, namely in order to avoid confusion about the genetic definition of this variety.

Phylogenetically, it is uncertain whether this variety retains an ancestral morphology of the common ancestor of *C. adoensis* var. *adoensis* and *C. senensis* or whether the longer trichomes are homoplastic due to an adaptive nature or this is a case of incomplete lineage sorting. Given the strong impact of aridification caused by the ice ages, the ancestor of *C. adoensis* and *C. senensis* presumably survived during an arid era in more humid coastal “forests” and woodlands of East Africa, where it evolved to *C. senensis* and *C. pwaniensis*. Other morphs evolved in woodlands rather in the inland, and are now pooled as *C. adoensis*. Interestingly, the distribution of *C. adoensis* var. *jeffreyana*, *C. senensis*, and the allied *C. pwaniensis* (shares the subulate calyx lobes with *C. senensis*) is very similar to that of the Apocynaceae species *Carvalhoa campanulata* K.Schum. (Leeuwenberg 1985), which suggests shared ecological preferences.

The collections from Singida occur in drier habitats than those from C and S Tanzania. Collections with an indumentum like *C. adoensis* var. *jeffreyana* also occur in NE D. R. Congo (*A. Taton* 128, *G. Troupin* 570), but it is uncertain whether these are also genetically linked to *C. adoensis* var. *jeffreyana*, so they are listed here under *C. adoensis* var. *adoensis*.

Specimens examined. (selection, in total: 26) Kenya. Rift Valley Province: Nanyang, cultivated in Munich Botanical Garden, *N. Holstein* 125 (M) and 130 (M). Malawi. Northern Region: Mzimba Dist., 3 mi. W of Mzuzu at Katoto, *J. Pawek* 10404 (MO, WAG [WAG0234128]). Tanzania. Iringa: Ludewa district. Livingstone Mountains, E slope of Msalaba Mountain, above stand of *Acacia abyssinica* on foot trail

from mission at Luana, 09°59'S 34°36'E, *R.E. Gereau & C.J. Kayombo* 3535 (DSM, EA, MO, NHT, PRE); Great North Road, Sao Hill, 61 ml. S of Iringa, *R. Polhill & S. Paulo* 1722 (B, EA, P [P05621244], PRE). Mbeya: Nyassa Hochland, Station Kyimbila, *A.F. Stolz* 504 (JE, M, U, W). Singida: 8½ km along road from Singida to Sepuka, 04°46'35"S 034°40'00"E, *P. Kuchar* 23919 (MO, S [S08-12129]).

3. *Coccinia barteri* (Hook.f.) Keay, *Kew Bull.* 8: 82. 1953.

Staphylosyce barteri Hook.f. in Oliv., *Fl. trop. Afr.* 2: 554. 1871. *Physedra barteri* (Hook.f.) Cogn. in A.DC. & C.DC., *Monogr. Phan.* 3: 525. 1881.

Type: Nigeria. Nupe [Niger State]: exact locality not specified, male, fl, *C. Barter* 1525 (Lectotype, designated by Jeffrey (1967: 60): K!, digital image! [K]).

Type: Equatorial Guinea. [Fernando Po] Bioko Island, female, *C. Barter* no. ? (Syntype: K?), see Taxonomic remarks.

Coccinia macrocarpa Cogn., *Bull. Jard. Bot. État Brux.* 5: 113. 1915–1919. Pro parte [except *E. Luja* 125].

Type: D. R. Congo. Sankuru river [tributary of Kasai river], no detailed location given, on farmland and in bushland, male, fl, Jul 1904, *E. Luja* 205 (Lectotype, designated by Jeffrey (1967: 61): BR! [BR0000008888570, digital image: BR, JPS, photo: K!]).

Type: Sankuru river [tributary of Kasai river], no detailed location given, female, fr, Nov 1903, *E. Luja* 125 (Syntype: BR! [BR0000008888228, digital image: BR, JPS]).

Coccinia subhastata Keraudren, *Fl. Cam.* 6: 131. 1967.

Type: Cameroon. South Region: Bitye, male, fl, 1917, *G.L. Bates* 1469 (Holotype: BM!).

Description. Perennial climber. Stems up to 10 m, glabrous or puberulous. Petioles 1–3.5(–8.5) cm, glabrous to puberulous, adaxial side rarely with trichomes. Leaves 3.5–20 × 4–23 cm, cordate, subhastate, shallowly to deeply 3- or 5-lobate. Lobes triangulate, ovate to oblong. Margin entire with few to many teeth to serrate. Apex obtuse to acute, with apical tooth. Upper leaf surface glabrous with clear or white pustules, lower leaf surface glabrous to puberulous on main nerves, esp. towards base, with or without small dark glands. Probracts ovate to elliptical, up to 5 mm long or missing. Tendrils simple or bifid. Male flowers in few- to many-flowered racemes. Common peduncle up to 3–8 mm long, glabrous to puberulous. Pedicel < 8 mm, indumentum like peduncle. Flowers without or with up to 1.5 mm long bracts. Perianth tube glabrous to puberulous. Calyx lobes 1–2.5 mm, subulate, lineal, rarely somewhat lanceolate, reflexed, spreading or erect and adpressed to corolla, sometimes seemingly fleshy. Corolla 1.1–2.4 cm, salmon, yellow to orange-yellow, lobes up to 3–10 mm. Filament column, anther head, and pollen sac color not seen. Female flowers in racemes, sometimes accompanied with a solitary flower or one solitary flower only. Peduncles and petioles in racemes like in males. Solitary female flowers with up to 1.5 cm long glabrous to puberulous pedicel. Ovary glabrous. Hypanthium glabrous to puberulous, calyx lobes and corolla as in males. Style not seen. Stigma

shape not seen, more or less dark yellow. Fruit 1.5–2.5 × 1.5 cm, shortly elliptical to subglobose, unripe green with pale spots, ripe red. Seeds 5.5 × 2.5–3 × 1–1.5 mm (L/W/H), more or less symmetrically obovate, face flat to flatly lenticular.

Phenology. Flowering time: January–June, August–November.

Distribution. Fig. 24. Humid tropical West Africa, Angola (Cabinda, Cuanza Norte), Burundi?, C and S Cameroon, C and S Central African Republic, D. R. Congo, R. Congo, Equatorial Guinea, Gabon, Mozambique (Manica), S South Sudan?, Uganda (Western, Central), Rwanda?, W Tanzania, Zambia (Northern Province), Zimbabwe (Manicaland). Elevation from sea level to 1650 m. Soil preference not well known, on loam soil, on granite (*J.B. Gillett 15298*). (*Newtonia*) rainforest; forest margins; near open water with *Pandanus candelabrum*, *Oxystigma mannii*, and *Raphia vinifera*; near river with *Saba comorensis*; on border of gallery forest and *Terminalia glaucescens* woodland; in riverine bushes, on river islands with *Alchornea cordifolia*; in fallows.

Use. The Turumbu people mash young leaves, mixed with white argil, and put the paste onto the heads of ill children (*W. Kesler 1034*).

Vernacular names. Lissongo [Mbatl]: makpo (*C. Tisserant (Équipe) 2250*); Twi: isamaŋ kyěkyěa (*F.R. Irvine 2604*); Turumbu: eliki e litoko (*J. Louis 2253*), ndombo di ilo (*W. Kesler 1034*).

Remarks. *Coccinia barteri* is treated here in a wide sense as it contains several forms (see also Holstein and Renner 2011b). This is because data on these forms are scarce and do not unambiguously allow to refer to these as species. Therefore, the present author refrains from creating an intraspecific classification as a phylogeographic treatment and crossing experiments appear to be necessary to clarify this problematic taxon.

There are collections in Gabon that are of intermediate morphology between *C. barteri* and *C. racemiflora* (*M.A. van Bergen 490* (WAG) = *C. barteri* 6 in Fig. 17). Holstein and Renner (2011b) suggested that hybridization occurs between these species. Whether the hybrids are fertile or sterile is not known.

Taxonomic remarks. *Coccinia barteri* (Hook.f.) Keay is type species of the genus *Staphylosyce* Hook.f.

Joseph Dalton Hooker mentions collections from Fernando Po [Bioko Island] and Nupe in the protologue of *C. barteri*. He only gives the name of Barter, whose Nupe specimen is in K, but there are no *Coccinia* specimens by Barter from Fernando Po. However, there are two specimens collected by G. Mann (*Mann N199!* and *N1166!*) in Hooker's herbarium (now in K). These were collected on that island, and they contain drawings that were most likely the basis for Hooker's description of *Staphylosyce barteri*. Possibly, Hooker mistakenly left out Mann's name in the protologue, whose collections contain many type specimens (Hooker 1871).

Keay published in error *Coccinea barteri* [*sic*] in his new combination, but accepted the species as belonging to *Coccinia* in Hutchinson and Dalziel's Flora of Tropical Africa (1954).

The syntypes of *Coccinia macrocarpa* certainly belong to different taxa. The present author concurs with Kéraudren, who placed the male specimen *É. Luja 205* into the polymorphic *Coccinia barteri* (1967). However, the female plant *É. Luja 125* is clearly not part of *Coccinia*. *Coccinia* seeds are up to 7 mm long, at the base attenuate to trun-




Figure 24. Distribution map of *C. barteri* (based on 99 collections).

cate and with a rounded apex. In contrast, the seeds of *É. Luja* 125 are subquadratic as Jeffrey already pointed out on the type specimen. A placement in *Momordica* by Jeffrey (on the sheet) seems to be correct, whether this is *M. multiflora* Hook.f. (1871) as identified by Jeffrey or *M. parvifolia* Cogn. (1916) as identified by Kéraudren is beyond the present author's knowledge.

Coccinia subhastata was described under the assumption that *C. barteri* has long calyx lobes, as it can be seen in *Flore du Cameroun* (Kéraudren 1967). Monique Kéraudren in her research on western Central African Cucurbitaceae (Kéraudren-Aymonin 1975a; Kéraudren 1967) treated *Coccinia/Physedra barteri* and *P. heterophylla* as synonymous. However, she confused the long subulate calyx lobes of *P. heterophylla* as a character for *C. barteri*, describing a specimen with short calyx lobes and flowers in long racemes as a new species, *Coccinia subhastata* Keraudren. She also described several differences of *C. subhastata* to *C. barteri*, which are not supported when carefully examined. *Coccinia subhastata* should only have simple tendrils, but the holotype of *C. subhastata* also has a bifid tendril. Furthermore, the *C. barteri* lectotype *C. Barter* 1525 has a subhastate leaf and simple tendrils. The description of *C. subhastata* thus is thus wrong and the species is a synonym of *C. barteri*, as it has been pointed out by Holstein and Renner (2010). In addition to the confusion of *C. heterophylla* and *C. barteri*, Kéraudren separated the western Central African specimens with few-flowered racemes as *C. keayana* R.Fern. (Kéraudren

1967). *Coccinia keayana*, however, does in fact not occur in this region but only in West Africa, and her *C. keayana* specimens from Cameroon belong to the polymorphic *C. barteri*.

Specimens examined. (Selection, in total: 153) Angola. Cuanza Norte: Cazengo municipality, near Agricultural Station Cazengo, *J. Gossweiler* 5492 (COI, LISU), and 5507 (LISU). Benin. Atlantique: Allada commune, Dahoukpa (Niaouli), 6°44'N, 2°07'E, *A. Akoègninou & F. Bada* 2992 (WAG [WAG0314946]). Cameroon. South Region: Bipinde [Bipindi], *G.A. Zenker* 1657 (E, G (4), HBG, P [P05621189], P [P05621192], W, Z). Central African Republic. Lobaye: Boukoko, *C. Tisserant (Équipe)* 2176 (BM, P [P05620598], P [P05621175]). D. R. Congo. Katanga: [Haut Katanga district], 40 km on road from Lubumbashi to Sakanja, *A. Schmitz* 4465 (EA (2), WAG [WAG0225507], WAG [WAG0225508]). Equatorial Guinea. Bioko Norte: Malabo–Punta Hermosa km 9, 32NMK8114, *F.J. Fernández Casas* 12077 (BM, MA n.v., MO, WAG [WAG0069018]). Gabon. Haut-Ogooué: 21 km on road from Okonja to Akiéni, 0°45.84'S, 13°47.01'E, *J.J. Wieringa et al.* 6387 (M, WAG [WAG0250956], WAG [WAG0250957], WAG [WAG0250958]). Ghana. Volta: Agumatsa Wildlife Sanctuary, at town Wli-Agorviefe, W of Park Guard HQ, 7°06'46"N, 0°35'25"E, *H.H. Schmidt et al.* 2192 (K, MO). Guinea. Nzérékoré: [Beyla préfecture], Bola [Boola], Famondou [Famodougou], 8.48068°N 8.70129°W, *E. Achigan-Dako* 07 NIA 899 (GAT (3)). Ivory Coast. Lagunes: Abidjan, Banco Forest Reserve, in marshy valley, near the entrance, *J. de Koning* 6144B (WAG [WAG0099439]). Nigeria. Rivers: Old GRA [Government reservation area], Port Harcourt, *B.E. Okoli* 150 (IFE (cited and picture of plant in Okoli 1984)). R. Congo. Sangha: 36.89 km E-SE Bomassa, 2°11.85'N, 16°31.10'E, *S.T. Ndolo Ebika* 401 (E [E00486113]). Sierra Leone. Kono: Tingi Mts, top of E ridge, *J.K. Morton & D. Gledhill* SL1886 (WAG [WAG0225480], WAG [WAG0225481]). Tanzania. Kigoma: Kigoma Rural District, Gombe Stream National Park, Mitumba Valley, research staff houses, 4°39'11"S, 29°38'09"E, *G. Gobbo et al.* 471 (MO). Zimbabwe. Manicaland: Chirinda Forest, *B. Goldsmith* 39/62 (COI, K, NY).

4. *Coccinia grandiflora* Cogn., Bot. Jahrb. Syst. 21: 211. 1895.

Coccinia grandiflora Cogn. ex Engl., Abh. K. Preuss. Akad. Wiss.: 34. 1894. nom. nud.
Type: Tanzania. [Tanga]: Mlalo, dry hill range. *C.H.E.W. Holst* 506a (Holotype: B, destroyed).

Type: Tanzania. Tanga: Usambara, near Amani, male, fl, [*H.J.P.?*] *Winkler* 3611 (Neotype, designated here: BR!).

Coccinia engleri Gilg, Bot. Jahrb. Syst. 34: 354. 1904.

Type: Tanzania. [Tanga]: West Usambara, Sakare [Sakarre], at waterfall in primeval forest, 1100 m, fl, fr, Sep, *A. Engler, Reise nach Ostafrika* 948 (Holotype: B, destroyed).

Type: drawing in protologue (Lectotype, designated here).

Description. Perennial climber. Stems up to 20 m, glabrous or (when from higher altitudes) sparsely covered with long, whitish trichomes. Petioles 2.5–13 cm, indumentum

as on stem. Leaves 12–20 × 11–20 cm, profoundly 5-lobate. Lobes triangulate, ovate to oblong. Leaf margin smooth to slightly serrate, dentate. Apex obtuse to acute with final tooth. Upper leaf surface glabrous with small hyaline pustules. Lower leaf surface glabrous, rarely with few trichomes on the main nerves esp. at base, with blackish glands scattered esp. along main nerves. Probracts up to 5 mm long (Fig. 8a). Tendrils bifid. Male flowers solitary or in (usually few-(–10-)flowered) racemes. Common peduncle 4–12 cm, glabrous. Pedicels of flowers in racemes 0.2–1.3 cm, glabrous. Bracts up to 3 mm or missing. Pedicel of solitary flowers 4–15 cm, glabrous. Perianth tube glabrous. Calyx lobes (2–)4–13 mm, lineal, narrowly lanceolate to triangulate, tip subulate to sub-acute. Corolla 4–6.5 cm long, apricot, salmon, yellowish-buff to yellow, lobes 2–4.7 cm. Filament column and anther head not seen. Pollen sac yellow-orange. Female flowers solitary, rarely in racemes, glabrous. Common peduncle < 1 cm, pedicel in racemes, pedicel of solitary flowers 3–4 cm. Ovary long cylindrical, glabrous or with hyaline–whitish pustules. Hypanthium glabrous, calyx lobes and corolla like in male flowers. Style columnar, yellowish to buff. Stigmas 2-lobed, yellow. Fruits < 30 × 2–4 cm, long cylindrical, glabrous, when unripe green, ripe (orange-)red. Seeds 4.5 × 2.5 × 1–1.2 mm (L/W/H), symmetrically obovate, face flatly lenticular.

Phenology. Flowering time: January–December.

Distribution. Fig. 25. Southeastern Kenya (Coast Province: Taita Hills, coastal forests), Tanzania (SE Dodoma; Iringa: Uzungwa Mts; Kilimanjaro; Lindi; Morogoro: Nguru Mts, Ukaguru Mts, Uzungwa Mts; Mtwara; Tanga: Usambara Mts), Malawi (Northern Region: Misuku Hills; Central Region: Nchisi Mts; Southern Region: Lisao Hill), Mozambique (Manica: East African Highlands with foothills), Zimbabwe (Manicaland: East African Highlands with foothills). Elevation 30–1900 m. On rich red-brown clay, over limestone, on diabase outcrops. Coastal and lowland forests and forest margins (*Parinari* sp.-*Newtonia buchani* forests and others), riverine forests (with *Cola clavata*, *Synsepalum msolo*, *Sorindeia madagascariensis*), rarely in montane forests (*Cassipourea malosana*-*Teclea simplicifolia*-*Teclea nobilis*-*Olea mildbraedii*-*Tabernaemontana* forest), or Miombo from degraded forests.

Use. Fruits are reported to be either poisonous (*A. Peter* 56598) or edible (*W.J. Kindeketa* 630). Leaves cooked in water used against fever (*K. Braun* 714).

Vernacular names. Kihehe: mudesselema (*F. Haerdi* 617/0), Kipare: hotwe (*W.J. Kindeketa* 630), Kishamba: matombo shanga (*G.R. Williams* G43), Kisamba: matombo ya nyoka “snake breasts” (*M.A. Mwangoka & A. Kalage* 1578).

Remarks. The southern distributed individuals in Zimbabwe and C Mozambique often bear short trichomes, and the leaves are rather shallowly lobate, just as in *C. schliebenii*. These populations may represent hybrids or descendants of a non-differentiated common ancestor.

It is difficult to distinguish between *C. grandiflora* and *C. mildbraedii* in the Central Tanzanian highlands (Eastern Arc Mts). Both species also occur in high altitude forests and are clearly delimited by flower size. *Coccinia grandiflora* also has larger probracts than *C. mildbraedii*, but this is rarely well visible. *Coccinia grandiflora* may also be confused vegetatively with *C. barteri* in Mozambique and Zimbabwe.




Figure 25. Distribution map of *C. grandiflora* (based on 62 collections). For Tanzania the borders of the regions are given.

Taxonomic remarks. The *C. grandiflora* holotype by Holst was destroyed in the fire of the Berlin herbarium in 1944. The Winkler specimen was chosen as neotype because it was already designated as type in December 2008. There is no annotation on the type label, however, and it seems that this neotypification was not published. However, the Winkler specimen label bears Cogniaux' handwriting. Strangely, the Winkler specimen also states "mars 1892", with the 92 crossed out. This is the date when Holst collected his specimen; but H. J. P. Winkler collected in Tanzania in 1910.

As the holotype of *C. engleri* also was destroyed, the original material left is a drawing in the publication of the protologue. The drawing is of sufficient quality to synonymize unambiguously *C. engleri* with *C. grandiflora*.

Specimens examined. (Selection, in total: 105) Kenya. Coast Province: Taita-Deveta District, Taita Hills, Mbololo Forest, 3°19'S, 38°27'E, Mwambirwa Forest Station, *R.B. Faden et al.* 799 (EA, WAG [WAG0234141]). Malawi. Southern Region: Chiradzulu district, Lisau Hill above Njuli P.O., *R.K. Brummitt & I.H. Patel* 18534 (K, WAG [WAG0361192]). Mozambique. Manica: Chimoio, Garuzo mountain ridge, *J.G. Garçia* 5224 (LISC, MO). Tanzania. Arusha: above Saje, N side of Ngurdoto Crater rim, *P.J. Greenway & K. Kanuri* 13444 (EA, K, M, PRE). Kilimanjaro: Chome [ward], Njokava forest, *R. Abdallah* 814 (EA, NHT). Lindi: 40 km W of Lindi, Lake Lutamba, Mirola Valley, *H.J.E. Schlieben* 5905 (B, HBG, G, M, S [S08-12068], Z (2)). Morogoro: Kilosa district, Ukaguru Mts, along track between Mandege and Ihanga rock, c. 6°24'S, 36°56'E, *M. Thulin & B.E. Mhoro* 2877 (DSM, EA, K, MO). Tanga: western Usambara Mts, Shagai forest, forestry house near Sunga, *R.B. Drummond & J.H. Hemsley* 2783 (B, EA, K, S [S08-12075]); [eastern Usambara Mts], Amani, c. 50 m before gate of headquarters, 05°05'58.3"S, 38°39'11.2"E, *N. Holstein et al.* 96 (B, DSM, M), 97 (BR, DSM, M), 98 (DSM, M, WAG), 99 (DSM, M), and 100 (M). Zimbabwe. Manicaland: Chipinge district, outskirts of Chirinda [forest] above Msilizwe river, *B. Goldsmith* 2/63 (B, BR, S [S08-12072]); [near Mutare], S slope of Murahwa's Hill, *N.C. Chase* 8008 (COI, K, LISC, MO).

5. *Coccinia grandis* (L.) Voigt, Hort. suburb. Calcutt.: 59. 1845.

Bryonia grandis L., Mant. Pl.: 126. 1767. *Coccinia indica* Wight & Arn., Prodr. fl. Ind. orient.: 347. 1834. nom. illeg. [nom. superfl. as epithet has not been adopted]. *Coccinia grandis* M.Roem., Syn. Pepon.: 93. 1846. nom. illeg. [nom. superfl.] *Cephalandra indica* Naudin, Ann. Sci. Nat. Bot. ser. 5, 5: 16. 1866. nom. illeg. *Cephalandra grandis* (L.) Kurz, J. As. Soc. Beng. 46(2): 103. 1877.

Type: Sri Lanka. *Bryonia foliis subrotundis angulosis, momordicae facies* Burm., Thes. zeylan.: 49, t. 19, fig. 2. 1737 (Type: drawing in l.c.).

Type: Sri Lanka. *Vitis alba indica* Rumphius [G. E. Rumpf], Herb. Amboin. 5: 448, t. 166, fig. 1. 1747 (Type: drawing in l.c.).

Type: No detailed information, female, fl, *Anon. in Herb. Linn.* 1153.2 (Typelectotype, designated by Nazimuddin and Naqvi (1984): LINN! [digital image: LINN]).

Type: India. Gottori, male and female, fl, *Anon. in Herb. Linn. 1153.3* (Type?: LINN! [digital image: LINN]).

Type: India. [Gujarat]: Suratt [Surat], female, fl, *Anon. in Herb. Linn. 1153.12* (Type?: LINN! [digital image: LINN]).

Type: India. No detailed information, female, fl, *Anon. in Herb. Linn. 1153.13* (Type?: LINN! [digital image: LINN]).

Cucumis sativus var. *arakis* Forssk., Fl. aegypt.-arab.: 169. 1755.

Type: Yemen. [Al-Hudaydah Governorate]: Lohaja [Al-Luhayyah] [region?], Môr [Mawr], male, fl, *P. Forsskål 660* (Holotype: C [C10002122, digital image: JPS, microfiche IDC: 35 II, 7–8]).

Turia moghadd Forssk. ex J.F.Gmel., Syst. nat. 2(1): 403. 1791. *Turia moghadd* Forssk., Fl. aegypt.-arab.: 166. 1755. nom. inval. *Coccinia moghadd* (Forssk. ex J.F.Gmel.) Asch. in Schweinf., Beitr. Fl. Aethiop.: 251. 1867. *Cephalandra moghadd* (Forssk. ex J.F.Gmel.) Broun et Massey, Fl. Sudan: 105. 1929.

Type: Yemen. [Al-Hudaydah Governorate]: Lohaja [Al-Luhayyah], female, fl, *P. Forsskål 663* (Lectotype, designated here: C! [microfiche: IDC 110 I, 1–2]).

Type: ibid., male, fl, *P. Forsskål 662* (Syntype: C! [microfiche: IDC 109 III, 7–8]).

Type: ibid., *P. Forsskål 666* (Syntype: C! [microfiche: IDC 110 I, 3–4]).

Bryonia alceifolia [sphalm. *alceaefolia*] Willd. in Rottler, Neue Schriften d. Ges. Naturf. Freunde Berlin 4: 223. 1803. *Coccinia cordifolia* (L.) Cogn. var. *alceifolia* [sphalm. *alceaefolia*] (Willd.) Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 531. 1881.

Type: India. [Tamil Nadu]: Tiruchinapally [Tiruchirappalli], male and female, fl, Nov 1793, *Anon. [J.G. Klein, B. Heyne or J.P. Rottler] in Herb. J.G. Klein 177* (Lectotype, designated here: B-W! [B-W 18065], isoelectotype: K!).

Momordica covel Dennst., Schlüssel Hortus malab.: 23. 1818. *Cucumis pavel* Kostel., Allg. med.-pharm. Fl. 2: 738. 1833. nom. illeg. [nom. superfl.]

Type: *Covel* Rheede, Hort. malab. 8: 27, t. 14. 1688 (Holotype: drawing in l.c.).

Momordica bicolor Blume, Bijdr. fl. Ned. Ind.: 928. 1825–26.

Type: Indonesia. [Java], Kuripan, in calcareis [on calcareous ground?], *K.L. Blume 1012* (Holotype: L! [L 0587745]).

Momordica bicolor var. *a* Blume, Bijdr. fl. Ned. Ind.: 928. 1825–26. nom. inval. Indonesia. Maluku Province, Timor, *A. Zippelius s.n.* (L! [L 0587743]).

Momordica bicolor var. *b* Blume, Bijdr. fl. Ned. Ind.: 928. 1825–26. nom. inval. [Indonesia]. [Java], Parang Mts, calcareous mountains. *K.L. Blume 1016* (L! [L 0587744]).

Bryonia moimoi Ser. in DC. Prodr. 3: 305. 1828. *Moi-moi* Adans., Hist. nat. Sénégal: 159, Paris. 1757. nom. inval. *Coccinia moimoi* (Ser.) M.Roem., Syn. Pepon.: 93. 1846.

Type: Sri Lanka. *Bryonia folio anguloso acuto glabro* Burm., Thes. zeylan.: 48, t. 19, fig. 1. 1737 (Holotype: drawing in l.c.).

Momordica monadelphæ Roxb., Fl. ind. [2nd ed.], 3: 708. 1832. nom. illeg. [nom. superfl.] pro parte. *Bryonia foliis cordatis oblongis angulatis dentatis glabris*. Fl. Zeyl. 356. 1747. nom. inval.

Type: Sri Lanka. *P. Hermann Mus. Zeyl.* 2: 37 (Type: BM [BM000621642], BM [BM000621643]) [these two specimens are *Cucumis maderaspatanus* L.].

Type: Sri Lanka. *P. Hermann Mus. Zeyl.* 5: 225 (Lectotype, designated here: BM [BM000595000]).

Type: Sri Lanka. *P. Hermann Mus. Zeyl.* 5: 321 (Type: BM [BM000621089]).

Type: Sri Lanka. *Bryonia folio anguloso acuto glabro* Burm., *Thes. zeylan.*: 48, t. 19, fig. 1. 1737. (Type: drawing in l.c.).

Type: Sri Lanka. *Bryonia foliis subrotundis angulosis, momordicae facies* Burm., *Thes. zeylan.*: 49, t. 19, fig. 2. 1737 (Type: drawing in l.c.).

Type: Sri Lanka. *Vitis alba indica* Rumphius [G. E. Rumpf], *Herb. Amboin.* 5: 448, t. 166, fig. 1. 1747 (Type: drawing in l.c.).

Coccinia loureiriana M.Roem., *Syn. Pepon.*: 93. 1846. *Bryonia grandis* Lour., *Fl. cochinch.* 1(2): 595. 1790. nom. illeg. and *Fl. cochinch.* 2: 731. 1793. nom. illeg.

Type: Sri Lanka. *Bryonia foliis subrotundis angulosis, momordicae facies* Burm., *Thes. zeylan.*: 49, t. 19, fig. 2. 1737. (Type: drawing in l.c.).

Type: Sri Lanka. *Vitis alba indica* Rumphius [G. E. Rumpf], *Herb. Amboin.* 5: 448, t. 166, fig. 1. 1747 (Lectotype, designated here: drawing in l.c.).

Coccinia wightiana M.Roem., *Syn. Pepon.*: 93. 1846. *Coccinia cordifolia* (L.) Cogn. var. *wightiana* (M.Roem.) Cogn. in A.DC. & C.DC., *Monogr. Phan.* 3: 531. 1881. *Coccinia grandis* var. *wightiana* (M.Roem.) Greb. in R. Mansfeld & J. Schultze-Motel, *Verz. Landwirtsch. u. Gaertn. Kulturpfl.* 2: 929. 1986. *Coccinia indica* Wight et Arn. β , *Prodr. fl. Ind. orient.*: 347. 1834.

Type: India. [Tamil Nadu]: [= Wall. Cat 6711a], [*J.G. Klein, B. Heyne* or *J.P. Rottler*] in *Herb. Madras s.n.* (Syntype: E! [E00174668, digital image: E, JPS]).

Type: India. [Tamil Nadu]: [Chennai, Saidapet], Nopalry [= in Wall. Cat. 6711b or e], female, fl, *R. Wight 1124* (Lectotype, designated here: E! [E00174667, digital image: E, JPS]).

Type: India. [Tamil Nadu]: Negapatam [Nagapattinam], female, fl, fr, *R. Wight 1124* (Syntype: E! [E00174666, digital image: E, JPS]).

Type: unknown. *R. Wight 1124* (Syntypes: E! [E00174664], NY! [00172358, digital image: NY]).

Coccinia grandis var. *quinqueangularis* Miq., *Fl. Ned. Ind.* 1(4): 673. 1856.

Type: [Indonesia]. [Central Java]: near Soerakarta [Surakarta], *T. Horsfield s.n.* (Holotype: U!).

Coccinia schimperi Naudin, *Ann. Sci. Nat. bot.*, ser. 4, 8: 366. 1857. *Cephalandra schimperi* (Naudin) Naudin, *Ann. Sci. Nat. Bot.* ser. 5, 5: 16. 1866.

Type: Ethiopia. In Semen [Semien Mts], female, fl, 1854, *G.H.W. Schimper Herb. Abyss.* 1215 (Lectotype, designated here: P! [the specimen with thick branch and fruit], isotype: P!).

Type: Ethiopia. Biria Dekeno et Dschadscha, 5000', female, fl, 1853, *G.H.W. Schimper Herb. Abyss.* 1215 (Syntype: P!).

Type: Ethiopia. Dschadscha, 5000'–5500', male and female, fl, fr, 13 and 21 Jul 1853, *G.H.W. Schimper Herb. Abyss.* 1215 (Syntype: P!).

Type: Ethiopia. Dschadscha, 5000', male and female, fl, fr, 13 and 21 Jul 1853, *G.H.W. Schimper Herb. Abyss. 1215* (Syntype: P!).

Type: Ethiopia. without details, *G.H.W. Schimper Herb. Abyss. 1215* (Syntypes: BR! [BR0000005113958, digital image: BR, JPS], BR! [BR0000005114641, digital image: BR], BR! [BR0000008251053, digital image: BR, JPS], BR [BR0000008351050], G-DC!).

Type: Ethiopia. Dschadscha, female, 1853, *G.H.W. Schimper Herb. Abyss. 1215* (Syntype: BR! [BR0000005111923, digital image: BR]).

Coccinia palmatisecta Kotschy, Sitzungsber. K. Akad. Wiss. Math.-Naturwiss. Cl. Abt. 1, 51: 360–361. 1865.

Type: [South Sudan], Kyk [Ciec (a Dinka subtribe) realm, S of confluence of Bahr al-Ghazal and White Nile], male, fl, no date given, *M.L. Hansal s.n.* (Lectotype, designated here: W! [K neg. 4837]).

Type: [South Sudan]. marshes in Noer [Nuer] realm [S to E of Malakal], no detailed location given, *F. Binder s.n.* (Syntype: W! [K neg. 4838]).

Cephalandra indica var. *palmata* C.B. Clarke in Hook.f., Fl. Brit. Ind. 2(6): 621. 1879.

Bryonia alceifolia [sphalm. *alceaefolia*] Willd. in Rottler, Neue Schriften d. Ges. Naturf. Freunde Berlin 4: 223. 1803. *Coccinia wightiana* M. Roem., Syn. Pepon.: 93. 1846.

Type: India, [Tamil Nadu]: Tiruchinapally [Tiruchirappalli], male and female, fl, Nov 1793, *Anon. [J.G. Klein, B. Heyne or J.P. Rottler] in Herb. J.G. Klein 177* (Syntypes: B-W! [B-W 18065], K!).

Type: India. [Tamil Nadu]: [= Wall. Cat 6711a], *[J.G. Klein, B. Heyne or J.P. Rottler] in Herb. Madras s.n.* (Lectotype, designated here: E! [E00174668, digital image: E, JPS]).

Type: India. [Tamil Nadu], [Chennai, Saidapet], Nopalry [= in Wall. Cat. 6711b and e], female, fl, *R. Wight 1124* (Syntype: E! [E00174667, digital image: E, JPS]).

Type: India. [Tamil Nadu]: Negapatam [Nagapattinam], female, fl, fr, *R. Wight 1124* (Syntype: E! [E00174666, digital image: E, JPS]).

Type: unknown. *R. Wight 1124* (Syntypes: E! [E00174664], NY! [00172358, digital image: NY]).

Coccinia cordifolia (L.) Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 529. 1881. pro parte majore, non *Bryonia cordifolia* L.

Coccinia cordifolia (L.) Cogn. var. *genuina* Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 531. 1881. nom. invalid.

Description. Perennial climber or creeper. Stems up to 5 m, glabrous, when older often white pustulate. Petioles 0.5–5.5 cm, glabrous, rarely some trichomes on adaxial side. Leaves 3–11 × 3–13 cm, cordate to 3-lobate or 5-edged to 5-lobate, sometimes lobulate. Lobes triangulate, ovoid, oblong, to obovoid. Leaf margin dentate, teeth usually with yellowish-reddish to brownish gland (Fig. 7a), becoming black when dried. Margin rarely with short (< 1 mm), whitish trichomes. Apex obtuse to acute with final tooth. Upper leaf surface glabrous, more or less dense hyaline to white pustulate. Lower leaf surface glabrous, with glands that are usually framed with lighter color between major nerves, nerves sometimes with white pustules. Probracts < 1.5 mm or missing.

Tendrils simple. Male flowers 1(–3) solitary, rarely in short racemes. Peduncle 0.3–1.5 cm, glabrous, pedicels of flowers in racemes up to 3.2 cm, glabrous, pedicels of solitary flowers up to 4.5 cm, glabrous. Bracts inconspicuous (< 1 mm), often absent. Perianth tube glabrous. Calyx lobes 1.2–3.5 mm long, lineal, spreading to reflexed, often with reddish to brownish gland on lower side at the acute tip (Fig. 7a). Corolla 1.7–4 cm long, yellowish buff (mostly African collections) to snow-white (esp. outside Africa). Corolla lobes 0.7–1.7 cm. Filament column and anther head pale greenish, pollen sacs yellow. Female flowers 1(–2) solitary. Pedicel up to 1 cm, glabrous. Hypanthium glabrous, calyx and corolla like in males. Ovary glabrous. Style columnar, yellowish-green. Stigmas 2-lobed, greenish. Fruit globose to ellipsoid, cultivated also shortly cylindrical, 3–4.5 × 1.5 cm, glabrous, unripe greenish with few pale spots and/or lines, becoming scarlet red when ripe. Seeds 5–7 × 2.5–3.5 × 1.2 mm (L/W/H), asymmetrically obovate, face flat (Fig. 14f).

Phenology. Flowering time: All over the year, but not at the end of the dry season or in cold seasons. Seems to need 1–2 weeks of at least 10 hours daylight with sunny weather for flower induction (pers. observ. from greenhouse cultivation).

Distribution. Figs 26, 27. Benin, N Cameroon, S Chad, D. R. Congo (in Great African Rift valley), Djibouti, Egypt (along Nile, Elba Mts), Eritrea, Ethiopia, Kenya, Mali (except N), S Mauritania, Niger, Senegal, Somalia, South Sudan, S and E Sudan, N Tanzania (Arusha, Dar es Salaam, Manyara, Morogoro, Mwanza, Pwani, Tanga, Zanzibar), Uganda, mountains and oases of the southern and western Arabian Peninsula, tropical and subtropical India, subtropical Nepal, Pakistan, Sri Lanka, South East Asia, S China incl. Hainan. Occurrence likely due to recent introduction in: Australia (Northern Territory, Queensland, Western Australia), R. China, Maldives, Mauritius, Mozambique, USA (Florida, Guam, Hawaii, Wake Island), many tropical Pacific islands, Caribbean area, Central and tropical South America. 0 to 1600 m. Black clay soil, black cotton soil, sand, on limestone. There seems to be a certain salt tolerance (Bharathi 2007). Grasslands, bushlands, (semi-arid) savannas, thickets, along rivers, ruderal sites, seemingly avoiding closed canopies (woodlands and forests). *Coccinia grandis* is, especially in tropical Asia, often reported from sand or from calcareous grounds (karst areas), which are well-drained. Although precipitation in South East Asia is much higher than in Africa, the quick loss of surface water allows *C. grandis* to survive there.

Use. Fruits (raw and cooked) and shoots (cooked) edible. The Luo eat the leaves as spinach (Orech et al. 2005). The sap is used against diabetes (Ramachandran and Subramaniam 1983) due to its hypoglycemic activity (Chopra and Bose 1925; Shibib et al. 1993). It is used in traditional Indian medicine in India for various diseases, and seems to have a general antibacterial effect (see also Use, economic potential, and phytochemistry).

Vernacular names. Agau or Chomir [most likely: Khamir/Xamtanga language of Agaw language family]: amballa gosa (*G.H.W. Schimper* 365, Schweinfurth 1893); Arabic: mogad (Getahun 1974b); Bagirmi: na odio (*A.J.B. Chevalier* 9527); Bengali: talakucha (Nadkarni and Nadkarni 1976), tela kucha [tēlākucā] (*J. Sinclair* 4414); Canarese [Kannada]: tonde-konde, tonde-kayee [tonḍe kāyi] (Nadkarni and Nadkarni 1976); Chinese: hong gua [Hóng guā] (Wu et al. 2011); Dassanetch [Daasanach]: dali (*C.J.*




Figure 26. Partial distribution map (Africa to Australia) of *C. grandis* (based on 629 collections).




Figure 27. Partial distribution map (Pacific Ocean and Americas) of *C. grandis* (based on 629 collections). State borders are given for the USA.

Carr & *C. Metolo* 867); Galinya [Oromo]: raho (*T. Ebba* 624; Getahun 1974b); Gujarati: gholi (Nadkarni and Nadkarni 1976); Hindi: kan-duriki-bel (Nadkarni and Nadkarni 1976), kundru [kandrū] (Bharathi 2007); Karen [in Mae Hong Son Province, Thailand]: khae-do (de Wilde and Duyfjes 2008); Khmer: (sloek') bah (Kéraudren-Aymonin 1975b); Kizigua: lukewja (*R.E.S. Tanner* 2691), pondwa (*R.E.S. Tanner* 2030); Lao: (khua 'phāk) tām ling, tām nhing (Kéraudren-Aymonin 1975b), phak tam nin (*J.-M. Dubost* 200); Luo: nyathud gush (*S.H. Padwa* 71); Mahrati [Marathi]: ran-tondula, ton-dli (Nadkarni and Nadkarni 1976); Malayalam: kova (Nadkarni and Nadkarni 1976), kovakka (Bharathi 2007); Masai: olaposhi (*C.H. Pecler* 3); Nepali: gol kankri (Aryal 2007); Ngadha (Baba-Ngadha): `uta kala (*J.A.J. Verheijen* 5415); Ngadha (Mataloko-Ngadha): uta kobho (*J.A.J. Verheijen* 5415); Persian [Farsi]: kabare-hindi (Nadkarni and Nadkarni 1976); Punjabi: kanduri (Nadkarni and Nadkarni 1976); Sanskrit: vimboshta,

vimbaja, bimba, tundika [tuṇḍikā] (Nadkarni and Nadkarni 1976); Serer: ayandyaloup (*J.-F. Ezanno Cucurbitaceae* 9); Sinhala: kowakka (*D. Philcox et al.* 10454, *D. Philcox et al.* 10458); Somali: masskar (*T. Ebba* 804); Songhai: lombaria (*A.J.B. Chevalier* 2996); Sunda: aroi papassang (Miquel 1855); Tamasheq [Touareg]: seffel (*H. Lhote* 2). Tamil: kovai [kawai] (Nadkarni and Nadkarni 1976), kovakkai (Bharathi 2007); Telugu: dondakaya [doṇḍa kāya] (Bharathi 2007), dondatiga, kakidonda (Nadkarni and Nadkarni 1976); Thai: in Central Region: phak tamlueng, in Northern Region: phak khaep (de Wilde and Duyfjes 2008); Tigrinya: asumbek (*G.A. Schweinfurth & D. Riva* 1007); Vietnamese: northern: hoa bát, southern: (dây) bai bác, bình bát, ma'nh bát, lá bát (Kéraudren-Aymonin 1975b); Wolof: barbouf (*J. Trochain* 663).

Remarks. With the exception of South and South East Asia, *C. grandis* is easily recognizable, especially by the lack of an obvious indumentum and the pale framed (in living state) glands in the axils of the nerves at the base of the lower leaf lamina. In NE Africa, collections with finely dissected leaves can be similar to *C. ogadensis*. When compared to collections from South Africa, *C. sessilifolia* var. *variifolia* and some forms of *C. mackenii* are also similar, but *C. sessilifolia* var. *variifolia* is glaucous, *C. mackenii* plants have bifid tendrils, and both South African species lack colored leaf teeth and have erect to spreading calyx lobes instead of spreading to reflexed calyx lobes. In South and South East Asia, some vegetatively similar *Gymnopetalum* species (e.g., *G. chinense*) can be mistaken for *C. grandis*, as in both taxa the leaf shape is 5-edged to cordate, and glands on the lower leaf lamina can be found. However, *Gymnopetalum* species are rather densely beset with trichomes, have ribbed fruits and are monoecious, whereas *C. grandis* is glabrous, has smooth fruits, and is dioecious, at least in wild populations.

Asian and (at least most of the) African *C. grandis* differ genetically (e.g., in a short sequence in the 5'-beginning of the *LEAFY*-like 2nd intron) and in petal color (white in Asian, buff in African individuals). Hence, the distribution in Asia (at least India) is not due to human impact. Whether *C. grandis* is introduced or native to Malesia, northern Australia and southern China and Taiwan is not known. Crossing experiments by Naudin (1862) indicate full compatibility between the African and Asian morphs though.

Taxonomic remarks. Up to the 21st century (e.g., Bulbul et al. 2011; Hussain et al. 2011), there has been quite a lot of confusion about the valid name of the species that is now called *Coccinia grandis*. Wight and Arnott (1834) established the genus *Coccinia*, with the name based on the scarlet-red fruits of a species that Wight had collected several times during his 13-year stay in South India (Stafleu and Cowan 1988). The name of the only species they described in their new genus, *Coccinia indica* Wight & Arn., is illegitimate since they included *Bryonia grandis* L., of which they ought to have adopted the epithet (Art. 52.1 ICN). The illegitimacy of *Coccinia indica*, however, does not affect the legitimacy of the genus name *Coccinia* (Art. 42.2), although there was also some confusion about which genus name to use. In 1845, Voigt published the correct combination *Coccinia grandis* (L.) Voigt with a description exactly matching *Bryonia grandis*.

Although *C. indica* is not valid, and the problem seemed to have been solved, a third species name was brought into discussion by Cogniaux. He thought that *Bryo-*

nia cordifolia L. and *Bryonia grandis* L. referred to the same species as Linnaeus cited Rumphius' *Vitis alba indica* (*C. grandis*) under *Bryonia cordifolia* (Linnaeus 1763). Hence, Cogniaux created a more broadly circumscribed *Coccinia cordifolia* (L.) Cogn. (1878). However, Linnaeus described *Bryonia cordifolia* beforehand (1753) citing the description from his *Flora Zeylanica* number 354 (1747), and only in the 2nd and 3rd edition of his *Species Plantarum* did Linnaeus synonymize Rumphius' *Vitis alba indica* (1747). Cogniaux therefore erred, when he stated that Linnaeus had based his *B. cordifolia* on Rumphius' figure, and the epithet *cordifolia* is hence misapplied in *Coccinia*.

There are four specimens in Linnaeus' herbarium, which belong to *Coccinia grandis* but it is unclear which are type material. The number 1153.2 is designated (by Linnaeus?) as *Bryonia grandis* on the sheet and is therefore the best choice for lectotypification as it has been done by Nazimuddin and Naqvi (1984). The specimen 1153.13 bears the note "*Bryonia foliis subrotundis angulosis, momordicae facie*" on the specimen flip side referring to one of the citations to drawings in the protologue. Hence, this specimen is also original material. The other two specimens (1153.3 and 1153.12) do not bear indications that Linnaeus referred to them as *Bryonia grandis*, but there is no contraindication either, so they might be original material, too.

The drawing to *Bryonia foliis subrotundis, angulosis, momordicae facie* Burm. (Thes. zeylan.: 49, t. 19, fig. 1. 1737), which is original material of *Bryonia grandis* L. appears to be a product of artistic freedom. The calyx lobes of the three uppermost and the lowermost flowers are reflexed and match *Coccinia/Bryonia grandis* well, whereas the calyx of the other two flowers appears to consist of almost free elliptic petals, quite like in *Momordica foetida* (except for the soft spines that are missing in the drawing). The addition "*momordica facies*" seems to relate to this. Strangely, the drawing of *Bryonia folio anguloso acuto glabro* Burm. (Thes. zeylan.: 48, t. 19, fig. 1. 1737) matches the current definition of *Bryonia grandis* also well, but has not been cited by Linnaeus. Eventually, the synonyms that Burman used, which also include *Bryonia cordifolia* L. (*Cucumis maderaspatanus* L.), made it hard to interpret the species and also lead to the confusion of Cogniaux.

The Forsskål 660 specimen (C10002122) has a hand-written field label by Forsskål (according to notes in JStor Plant Science) on the flip side, stating "*Cucumis incerta. Arakis, Mour.*" (*incerta* meaning "uncertain") and two hand-written identifications "*Cucumis inedulis* Fl. Yemen CXXII nr. 580" and "*c. s. Arakis cent VI nr. 61 p. 169*". The location in the text is the same as on the field label: Môr/Mour [Mawr, a small town about 30 km E of al-Luhayyah]. The former identification is a nom. nud. with the number 580 on page CXXII of Forsskål's *Flora Ægyptiaco-Arabica* (1775). The second identification links to a *Cucumis sativus* variety that is validly described in that book on page 169. The description matches well *Coccinia grandis* except for the tuberculate ovary. Additionally the collection is supposed to be from Loheja [al-Luhayyah], but might only indicate the region, in which Mawr is localized. However, the Arabic name of *Cucumis inedulis* and *Cucumis sativus* var. *arakis* are both: Arakis [3raqīs], so they can be cross-referenced. The description of the variety also mentions that the plant is not edible, just as the supposed species epithet.

The genus name *Turia* has been created by Forsskål in his *Flora Ægyptiaco-Arabica* (1775). There is a debate, however, whether it is validly published (Friis 1984). The descriptions of this part of Forsskål's book (page number in Latin) must be used in consideration with the corresponding parts in the lists of local floras (Roman page numbers), which is on page CXXI in the case of *Turia*. The first species there is *Turia sativa* (no. 550), which is called "turia" in Arabic and is cultivated according to the epithet. This matches exactly the first description in the descriptive part of the book (p. 165). Forsskål lists five *Turia* species in the floristic part and describes five species in the descriptive part. Therefore, the genus *Turia* lacks a description (Jeffrey 1962), because Forsskål does not mention any character to be typical for the genus. The name *Turia* was legitimately described by Gmelin in an extension of Linnaeus' *Systema naturae* (1791), and so was the name *Turia moghadd*.

The typification of *Turia moghadd* is not straight-forward, because Forsskål added little, if any marks on the sheets (Friis 1983). The flip side of the *Forsskål* 663 specimen has three notes: "Cucumis glandulosus", a second one with a different handwriting: "Bryonia Turia 35 Forsk" and a third one with another handwriting "Turia gijef Forsk Cent. 6 no. 38" with all words except for "Turia" crossed out. It can be hypothesized that these are different trials to identify the specimen, but not by Forsskål himself. Aside from these, the specimen lacks any written marks, but it matches exactly the description of *Turia moghadd*, as do the specimens *Forsskål* 662 and 666, which are syntypes. The *Forsskål* 663 specimen is chosen to be the lectotype, because the original description mentions the occurrence of fruits and this specimen is the only female of the three.

Willdenow describes *Bryonia alceifolia* in a travel report by J. P. Rottler from 1799, but he only mentions that he separates Rottler's *Bryonia epigaea* from another new species, viz. *B. alceifolia*, so the type was not necessarily collected in 1799. Willdenow knew *B. alceifolia* from Klein's specimens in his herbarium. Rottler was missionary in Tranquebar, the same place in which J. G. Klein was surgeon (Jensen 2005). Both are known to have collected together (with B. Heyne). The lectotype collection is in the describer's herbarium under the Klein number 177, but it is not certain, whether Klein collected the specimen himself, or if it was by Rottler or even Heyne. A duplicate with the same label data as in B-W is deposited in K. For Willdenow described the species, the specimen in his herbarium was chosen to be lectotype.

The name *Momordica monadelphica* Roxb. is superfluous, because Roxburgh synonymized *Bryonia moimoi* Ser. in total by citing the only element of that name and *Bryonia grandis*. The other elements of *M. monadelphica* are also interesting, though. Roxburgh cited "*Bryonia foliis subrotundis*" with the citation of Burman's *Thes. zeylan.*: t. 19, fig. 1. 1737 and fig. 2, *Vitis alba indica*, all *Coccinia grandis*, and Herman's *Musæum Zeylanicum* 356. The latter one consists of two specimens (2: 37), which are *Cucumis maderaspatanus* L., however, there are also two drawings (5: 225 and 5: 321) with the number 356. Both drawings represent *Coccinia grandis* because of the fruit size, fruit shape, and the flower morphology (calyx lobe length and position, corolla size), rather than *Cucumis maderaspatanus* L., *Diplocyclos palmatus* L. or *Cayaponia*

laciniosa (L.) C. Jeffrey. That the drawing 5: 225 (BM000595000) shows a plant with male and female flowers on one individual might be explained best by artistic freedom.

The name *Coccinia indica* var. *palmata* C.B. Clarke is valid and legitimate but not obvious to typify. Despite *C. indica* being illegitimate as a nomen superfluum, the variety is legitimate and validly described. Clarke cites *Bryonia alceifolia* Willd. and the *C. indica* protologue with page 348 although *C. indica* was described on page 347 (Wight and Arnott 1834). Page 348, however, only comprises the β variety, which itself was the basis for *C. wightiana* M. Roem. This unnamed variety consists of several elements, of which the literature citations of *Bryonia palmata* are mentioned with a question mark and are thus not eligible for typification (Art. 52.2 N1). The element “*Bryonia palmata* Linn.? herb. Madr.!” relates to a collection in Herbarium Madras that was identified a *Bryonia palmata*, but Wight and Arnott appear to have been in doubt whether the name was used sensu Linnaeus, hence the question mark after the name. This specimen (or a duplicate) is found in E and represents a deeply lobed *C. grandis* just as the protologue of the β variety says. Clarke obviously takes the epithet of the variety from this misidentification and not from *Bryonia palmata* L., which is why the name was typified with this collection. That Clarke meant a deeply lobed *C. grandis* is evident since he also cites *B. alceifolia*, which is also deeply lobed. It is thus not a new combination and status change but a new variety.

The name *Coccinia cordifolia* var. *genuina* Cogn. cannot be regarded as intended to represent a new variety, because Cogniaux divides all specimens he had seen into three varieties. Hence, the epithet “*genuina*” and his “*genuina*” variety is just synonymous with the autonym.

Specimens examined. (Selection, in total: 1066). Australia. Queensland: Weipa, Awonga Court, 12°37'28"S, 141°52'39"E, B.M. Waterhouse 7428 (BRI, CANB [CANB 682857]). Bangladesh. Sylhet: Sillet, F. de Silva in Wall. list 6700g (M [“439”], M [“440”]). Barbados. St. Michael: Bridgetown, old railway yard, Sep 1940, E.G.B. Gooding s.n. (BAR00002591). Brazil. Minas Gerais: No detailed location, < 1840, P.C.D. Clausen s.n. (G-DC). Cameroon. North Region: Pitowa [Pitoa], c. 17 km NE of Garoua, W.J.J.O. de Wilde et al. 4932 (MO, WAG [WAG0225471]). Chad. Chari-Baguirmi: Baguirmi et région du Lac Fitri, Tjecna, A.J.B. Chevalier 9527 (G, P [P05620497]). P. R. China. Hainan: Ch’ang-kiang District [Changjiang country], [Bawangling Nature Reserve], Ka Chik Shan [Qicha hill], S.K. Lau 3008 (GH n.v., P [P06394498], S). Hong Kong: near Ouang-nei-Tchong [Wang nai chung] – Happy Valley, E.M. Bodinier 1367 (E). Yunnan: Nan-chiao [Meng che], C.W. Wang 76988 (A n.v., IBSC [0207527], KUN n.v., PE [01178321], PE [01178322]). R. China. Pingtung County: Hsinpi [Sinpi], Y.-P. Cheng 4603 (BR [BR0000005164189], TAIF). Cuba. Havana: Santiago de las Vegas, A. O’Donovan 5106 (G, S [S08-12050]). D. R. Congo. Orientale: Mahagi territory, Mahagi-Port, J. Lebrun 3800 (BR (2), EA, WAG [WAG0225470]). East Timor. Dili: just W of Dili, Tasutolu area, 8°33'57"S, 125°30'05"E, L.D. Cowie 10658 (L). Egypt. Red Sea Governorate: [Hala’ib triangle], Gebel Shendib, G.W. Murray 3857 (K [K000037307]). Eritrea. Gash-Barka: Mai Mentai [at Sciagol River, SE of Agordat], N. Beccari 118 (FT).

Northern Red Sea Region: [NE of Ghimda/Gimda], Pianura Sabarguma, *A. Pappi* 3970 (G, MO, P [P05620558], S [S08-12133], W). Ethiopia. Oromia: 85 km NE of Nazareth, along road to Awash, c. 5 km W of Metahara, 8°55'N, 39°55'E, *J.J.F.E. de Wilde* 6870 (BR, MO, WAG [WAG0225448], WAG [WAG0225449], WAG [WAG0225450]). Tigray: near Djeladjeranné, 23 Apr 1841, *G.H.W. Schimper* 1570 (P [P05620544], specimens from BM, G, MO, S [S08-12147], and W [W 0011063] might also be from this location, for details see under *Cucurbita schimperiana*). Guyana. Demerara-Mahaica: Georgetown, *W. Hahn* 4810 (MO, US). India. West Bengal: Sibpur [Shibpur in Howrah city] near Calcutta [Kolkata], *J.G. Hallier s.n.* (M [“999”]). Indonesia. East Java: Pasoeroean [Pasuruan], 17 Jul 1928, *J.J. Ochse s.n.* (B, BR [BR0000009938076], L [L 0587578], P [P06394462], U). Mali. Kayes: Bafoulabé Cercle, Bamafélé arondissement, 3.2 km ESE of Manantali, 13°10.847'N, 10°26.260'W, *C.S. Duvall* 130 (MO (2)). Mozambique. Maputo: Vila Luísa [Marracuene], along Incomáti river, *A. Balsinhas* 1273 (COI). Nepal. Madhya Pashchim: [Rapti], Tulsipur, *O. Polunin et al.* 5906 (E). Nicaragua. Chontales: St Tomas [Santo Tomàs], 1841, *E. von Friedrichsthal s.n.* (W). Niger. Agadez: [Aïr Mountains], Mt Bagwezan [Mt Bagzan, Idoukal-n-Taghès], 31 May 1920, *A. Buchanan s.n.* (MO [acc. no. 1667128]). Pakistan. Sindh: 7 mls [c. 11.2 km] from Hyderabad to Tando M.[Mohammad] Khan, *S. ul-Abedin* 3839 (B, KUH n.v., MO). Senegal. Dakar: Jof [Yoff], *J.-G. Adam* 1806 (MO (2), [non P [P00694028/P05590411], which is a *Citrullus* sp.]). Singapore. South East District: Potong Pasir off Serangoon Road, *J. Sinclair* 5434 (E, L [L 0587600], P [P0639450]). Somalia. Togdheer: [c. 54 km SSW of Berbera], foot of Sheikh Pass, *P.R.O. Bally* 11827 (G (3), K, PRE). Sri Lanka. North Central Province: Anuradhapura, *F.R. Fosberg & N. Balakrishnan* 53431 (MO, PDA n.v.); *ibid.*, *W. Forstner s.n.* W13705 (W). Sudan. North Kurdufan: Chursi [Khursi], *K.G.T. Kotschy Iter nub.* 308 (BM, BR (2), E, G (2), G-DC, GOET, K, L, M [“400”], M [“401”], M [“782”], MO, P [P05620478], P [P05620484], P [P05620578], PRC, S [S08-12152], TUB [TUB004721], TUB [TUB004722], W [W 0011066], W (2), WAG [WAG0234161]). Tanzania. Tanga: Tanga town, near public beach, 200 m E of 05°03'19.2"S, 39°07'31.3"E, *N. Holstein et al.* 94 (B, DSM, M, WAG), and 95 (DSM, M). Zanzibar Urban/West: Chukwani, *H.G. Faulkner* 2782 (BR, COI, K n.v., P [P05620529]). Thailand. Bangkok: Bangkok, *R. Zimmermann* 66 (B (2), BR [BR0000009938878], G (2), H, L [L 0587657], M, MO, P [P06394565], PR [PR 801377], PRC, S, U, W). USA. Florida: along Indian River [lagoon], Coco [Cocoa], 9 May 1918, *J.K. Small s.n.* (FLAS [FLAS 29152], G, MO [acc. no. 1161454], US). Hawaii: O'ahu Island, Dole St, behind U Hawaii Manoa dorms, *B. Kennedy* 47 (MO, US [00674433]). Midway Atoll: Sand Island, 4208 Commodore Ave., *F. Starr & K. Martz* 080601-01 (BISH, PTBG [PTBG1000013099]). Vietnam. Da Nang: Tourane [Da Nang] and vicinity, no detailed location, *J. Clemens & M.S. Clemens* 3257 (A n.v., G, MO, P [P06394668], U, W, Z). Yemen. Al-Hudaydah: Mor [Mawr], 15°40'N, 28–30 Mar 1825, *F.W. Hemprich s.n.* (P [P06394550], S [S08-12138]).



Figure 28. Reconstruction of the habit of a female raceme of *Coccinia heterophylla* based on *T.-J. Klaine 414* (P). Black bar equals 0.5 cm.

6. *Coccinia heterophylla* (Hook.f.) Holstein, Kew Bull. 65(3): 435–441. 2010 [published in 2011].

Physhedra heterophylla Hook.f. in D.Oliv., F. T. A. 2: 553. 1871.

Type: Angola. [Cuanza Norte]: Golungo Alto, along the banks of the stream Casaballa, at the base of the mountains in Sobato de Bumba, male, fl, Oct 1855, *F.M.J. Welwitsch* 791 (Lectotype: BM! [BM000948006, digital image: BM, JPS], selected in Holstein and Renner (2010: 440); isolectotype p.p.: LISU! [LISU214547, digital image: JPS]).

Type: Angola. No detailed location, female, fr, Jan 1856, *F.M.J. Welwitsch* 791 (Syntype: BM! [BM000948008, digital image: BM, JPS]).

Type: Angola. In rugged places at Delamboa river, with *Coffea melanocarpa*, no clear date given, *F.M.J. Welwitsch* 791 (Syntype: BM! [BM000948007, digital image: BM, JPS]).

Type: Angola. No detailed location and date, *F.M.J. Welwitsch* 791 (Syntype: COI! [COI00005515, digital image: JPS]).

Type: Angola. No detailed location, male, fl, no date, *F.M.J. Welwitsch* 791 (Syntype: K! [K000313234, digital image: JPS]).

Type: Angola. No detailed location, female, fl, no date, *F.M.J. Welwitsch* 791 (Syntype: LISU! [LISU214548, digital image: JPS]).

Type: Angola. No detailed location, female, fr, Jan 1856, *F.M.J. Welwitsch* 791 (Syntype: LISU! [LISU214549, digital image: JPS]).

Type: Angola. No detailed location, male, fl, no date, *F.M.J. Welwitsch* 791 (Syntype: LISU! [LISU214550, digital image: JPS]).

Type: Angola. In rugged places at Delamboa river, with *Coffea melanocarpa*, Sep 1855, *F.M.J. Welwitsch* 791 (Syntype: LISU! [LISU214551, digital image: JPS]).

Type: Angola. At Delamboa river, no date, *F.M.J. Welwitsch* 791 (Syntype: LISU! [LISU214552, digital image: JPS]).

Type: Angola. No detailed location and date, *F.M.J. Welwitsch* 791 (Syntype: LISU! [LISU214553, digital image: JPS]).

Type: Angola. No detailed location and date, *F.M.J. Welwitsch* 791 (Syntype: P! [digital image: JPS]).

Type: Angola. No detailed location and date, *F.M.J. Welwitsch* 791 (Syntype: G-DC!).

Physhedra heterophylla var. *hookeri* Hiern in Cat. Welw. Afr. Pl. 1(2): 400. 1898.

Type: Angola. [Cuanza Norte]: Golungo Alto, near Ponte de Felix Simões, female, fl, Dec 1855, *F.M.J. Welwitsch* 792 (Holotype: BM! [BM000948009, digital image: BM]).

Description. Perennial climber. Stem up to 5–6 m, glabrous, sometimes whitish-speckled. Petiole 1–2 cm long, puberulous on abaxial side with few tiny trichomes, rarely with up to 0.8 mm long, yellowish trichomes, sometimes white-speckled, on adaxial side with small, yellowish-dirty trichomes. Leaves 7.5–12.5 × 10–14 cm, cordate, deltoid-subhastate to




Figure 29. Distribution map of *C. heterophylla* (based on 19 collections).

5-lobate, auriculate, or 7-lobate. Lobes triangulate to ovoid. Leaf margin dentate, often serrate. Apex acute with final tooth to acuminate. Upper leaf surface glabrous, small hya-

line pustulate. Lower leaf surface glabrous often with blackish glands, nerves glabrous, except for basis with up to 0.8 mm long, yellowish trichomes and sometimes white-speckled. Probracts up to 5 mm. Tendrils bifid. Male flowers in short racemes. Common peduncle 2–13 mm long, not exceeding the pedicel bearing part in length, puberulous with tiny yellowish-dirty trichomes (magnifying glass!). Pedicels < 4 mm, glabrous to puberulous. Bracts up to 3.5 mm. Perianth tube glabrous. Calyx lobes 5–7 mm, subulate, erect. Corolla up to 1.6 cm, dirty yellowish, dirty orange to reddish-orange. Corolla lobes 4–7 mm. Color of filament column, anther head, or pollen sacs not seen. Female flowers solitary or in short or long (up to 15 cm) racemes (Fig. 28). Raceme and bracts as in males, except raceme can be elongated (see remarks). Hypanthium glabrous, calyx lobes and corolla like male flowers. Ovary glabrous. Style columnar to 3-parted to the middle, color not seen. Stigma bulging, pale yellow. Fruits up to 4–6 × 2–2.5 cm, ovoid-elliptical to shortly cylindrical, glabrous. Fruits ripen from green? via orange-colored with longitudinal green bands, ripe fruits unknown. Seed size not known, rather symmetrically obovate, face flat.

Phenology. Flowering time: February–April, June, September–December.

Distribution. Fig. 29. Western Gabon (Estuaire), southern R. Congo (Kouilou), western D. R. Congo (Bas-Congo), western Angola (Cuanza Norte, Namibe). Elevation 10 to 900 m. Soil preference unknown. Transition between tropical lowland rainforest and woodlands, and its relict sites along the Angola highland escarpment, in secondary regrowths, on shrubs, along rivers.

Remarks. The long, subulate calyx lobes are the only good character for distinguishing this species, which otherwise can be easily confused with *C. barteri*. Female collections from Libreville (Gabon) and R. Congo have elongated racemes while the racemes are more condensed in the south. Whether this character shows affinity (introgression?) to *C. racemiflora*, which also has elongated racemes, defines an own species, or is just a coincidental observation of intraspecific variation is not known.

Taxonomic remark. This species is the type species of *Physedra* Hook.f. The genus was described by J. D. Hooker (Bentham and Hooker 1867) with three species belonging to it. However, in Oliver's *Flora of Tropical Africa* (Hooker 1871), Hooker only describes two species, *P. heterophylla* and *P. longipes*. When Jeffrey (1962) transferred *P. longipes* into a new genus, *Ruthalicia*, he indirectly lectotypified *Physedra*.

Monique Kéraudren regarded *P. heterophylla* as synonymous to *C. barteri* (Kéraudren-Aymonin 1975a; Kéraudren 1967) from which it, in fact, differs by the long calyx lobes. This, however, led to the creation of *C. subhastata* with short calyx lobes, which is a synonym to *C. barteri* (Holstein and Renner 2010).

Specimens examined. (Selection; in total: 37) Angola. Cabinda: Maiombe [Forest], Belize, *J. Gossweiler* 7653 (COI, K, LISU). Cuanza Norte: Cazengo municipality, near Agricultural Station Cazengo, *J. Gossweiler* 5655 (COI, LISU [LISC 031407], LISC [LISC 031408], LISC [LISC 031409], LISC [LISC031410], LISC [LISC 031411], LISU), and 5707 (COI, LISC [LISC 031412], LISU). Huíla: Chão da Chela, between [Lago] Tchivinguiro and Bruco, on middle slope of Serra da Chela escarpment, *E.J. Mendes* 925 (BM, COI, LISC); *ibid.* *L.A. Grandvaux Barbosa* 9448 (COI). D. R.

Congo. Bas-Congo: Lukula territoire, Temvo, *F.M.C. Vermoesen* 1824 (BR, EA, WAG [WAG0225504]). R. Congo. Kouilou: on left bank of Kouilou river, 4 km upstream of Kakamoeka, Sounda on path to level meter, *C. Farron* 4980 (P [P05621154], P [P05621155], P [P05621156]). Gabon. Estuaire: Libreville, *T.-J. Klaine* 414 (P [P05620605], P [P05620610], P [P05621146], P [P05621147], P [P05621149], P [P05621150], P [P05621152], P [P05621153]); Libreville, I.R.A.F. building, 0°25'N, 9°26'E, *J.M. Reitsma & B. Reitsma* 2120 (MO, NY, WAG [WAG0225493]).

7. *Coccinia hirtella* Cogn. in Schinz, *Bull. Herb. Boiss.* 4: 821. 1896.

Type: [South Africa]. KwaZulu-Natal: Howick, 3400', male, fl, 18 Feb 1895, *R. Schlechter* 6775 (Lectotype, designated by Meeuse (1962: 101): Z! [Z-000004443, digital image: Z]; isolectotypes: BR! p.p. [BR0000008887184, digital image: BR, see taxonomic remarks], GRA [GRA0002851-0, digital image: JPS], K! [K000542639, digital image: K]).

Description. Perennial creeper or climber. Stems up to 3 m, densely covered with long (> 0.5 mm), upright, whitish trichomes. Petioles 1.5–4 cm long, indumentum like on stem. Leaves 3–10 × 2.5–11 cm, 5-lobate. Lobes obovate or elliptical in outline, rarely ovate. Margin lobulate or coarsely serrate, (at maturity pale brownish) dentate (Fig. 13a). If margin serrate then lobe tips acute, else often rounded. Upper and lower leaf surface, esp. on the nerves, densely covered with long, upright, whitish trichomes. Probracts up to 3 mm. Tendrils simple or bifid. Male flowers solitary, rarely in few-flowered raceme. Common peduncle 1.8–4.5 cm, with white trichomes, pedicel of racemous flowers 0.2–3 cm, with white trichomes. Bracts up to 2 mm, caducous. Pedicel of solitary flowers 3–8.5 cm, with white trichomes. Perianth tube with white trichomes. Calyx lobes narrow triangulate to lineal, 3.5–6.5 mm long, spreading. Corolla 2–3.7 cm long, buff, lobes 1–2 cm. Filament column white, anther head pale yellowish, pollen sacs yellow to orange. Female flowers one solitary. Pedicel 0.5–5.5 cm, with long, white trichomes. Hypanthium with white trichomes, calyx lobes and corolla like in male flowers. Style columnar, whitish. Stigma bulging, yellow. Ovary with some long trichomes. Fruit oblong ovoid, 5–6 cm × 2.5–3 cm, sparsely covered with long trichomes to glabrous, ripening from green via green with longitudinal whitish mottling, via yellow, orange to red when ripe (Fig. 13a). Seeds 5.5–6.5 × 3–3.5 mm (L/W), seed height not seen, rather symmetrically obovate, face flatly lenticular.

Phenology. Flowering time: January–April, September, November, December.

Distribution. Fig. 30. Lesotho, South Africa (S KwaZulu-Natal, SE Free State). Afromontane scrubland and sourveld grassland. Elevation 110–1900 m. On sand, sandstone, loam, well-drained soils, full sun, grazing maybe tolerated.

Use. It is said that if a Masuto dreams unpleasantly about an ancestor (“balimo”), then relief is given after a bath with sun-dried *C. hirtella* roots and ironstone in a hole on the threshold of the “lelopa” (the circular fence around the hut) (Phillips 1933).




Figure 30. Distribution map of *C. hirtella* (pale yellow dots; based on 22 collections), *C. mackenii* (blue triangles; based on 50 collections), and *C. quinqueloba* (blue dots; based on 38 collections). For South Africa the borders of the provinces are given.

Vernacular names. Sotho: leraka-la-balimo (Jacot Guillarmod 1971; Phillips 1933), Sotho: monyaku (Jacot Guillarmod 1971).

Taxonomic remarks. The BR type specimen (BR0000008887184) contains two labels and is mixed. The female parts on the sheets have most likely been detached from the lectotype in Z, because it is a female branch with shallowly lobate leaves, just as part of the lectotype. The male parts on the BR specimen, however, are mixed. The leaves with the obtuse lobules are also likely type material, whereas the leaf with the acute lobules is very similar to the leaves of the *R. Schlechter* 6708 collection, which is not a type.

Specimens examined. (Selection; in total: 36). Lesotho. Leribe: Lérivé [Hlotse], *H. Dieterlen & A. Dieterlen* 145 (BR [BR0000008887467], P [P05620661], P

[P05620662], Z). Maseru: Roma, Map[h]otong, *M. Schmitz* 8039 (PRE). South Africa. Free State: Rooiberge area, Ross Kloof, *M. Jacobs* 8565 (L, LISU, PRE). Kwa-Zulu-Natal: 17 km from Nottingham Road on road to Loteni, *E. Retief* 1638 (MO, PRE); Albert Falls, *A.K. Meebold* 13160 (M); Giants Castle Game Reserve (Dinosaur Footprint area), *W.R. Trauseld* 930 (PRE).

8. *Coccinia intermedia* Holstein, *PhytoKeys* 7: 28. 2011.

Type: Benin. Atakora: Natitingou, Kouaténa (Perma), 10°12.00'N; 1°30.18'E, river bed, female, fl, fr, 3 Oct 2000, *A. Akoègninou et al.* 3625 (Holotype: WAG! [WAG0278370]; isotype: WAG [WAG0278369]).

Type: Ghana. Greater Accra: Shai Hills Game Reserve, monoecious, fl, fr, 25 May 1976, *J.B. Hall & J.M. Lock* GC 46016 (Paratypes: K! (4), MO!).

Type: Ivory Coast. Zanzan: Bouna, male, fl, 10 Aug 1967, *C. Geerling & J. Bokdam* 662 (Paratypes: MO!, WAG! [WAG0225492]).

Type: Togo. Maritime: between Lomé and Aného, female, fr, 25 Jun 1994, *L. Aké Assi* 18983 [typographical error in orig. publication stated 18982] (Paratype: MO!).

Description. Perennial climber. Stem length unknown, but likely several meters, glabrous, at maturity with clear to white pustules. Petioles 2.8–10.8 cm, glabrous, when older with clear to white pustules. Leaves 6–15 × 7–18 cm, shallowly to profoundly 5-lobate, more or less auriculate. Margin conspicuously dentate, blackening when dried. Apex acute. Upper leaf surface glabrous with clear to whitish pustules. Lower leaf surface glabrous, often with small dark glands near the leaf base. Probracts up to 2.5 mm long. Tendrils simple or bifid. Male flowers in few-flowered racemes, likely sometimes accompanied by a single flower. Common peduncle up to 1 cm, pedicels in racemose flowers 2–4 mm, each glabrous. Bracts up to 1.5 mm. Perianth tube glabrous, calyx lobes c. 1.5 mm, lineal to narrowly triangulate, erect with slightly recurved tips. Corolla 1.6 cm long, pale reddish-yellow to yellow, lobes 0.7 cm. Filament column and anther head not seen, pollen sacs yellowish. Female flowers 1–3 clustered (strongly reduced raceme). Pedicels 0.6–1.2 cm, glabrous. Hypanthium tube glabrous, calyx lobes and corolla like in male flowers. Ovary glabrous. Style and stigmas not seen. Fruit 4.5 × 2.5 cm, elliptical to oblong. Unripe fruit green with pale green longitudinal mottling, ripe orange?, more likely becoming red via orange ripening stage. Size of mature seeds unknown ($\geq 5.5 \times 3.5 \times 1.3$ mm (L/W/H)), symmetrically (to slightly asymmetrically) obovate, face flat.

Phenology. Flowering time: May, August, October.

Distribution. Fig. 31. NE Ivory Coast, SE Ghana (likely also in the north), S Togo (likely also in the north), NW Benin. Elevation sea level to 415 m. Wooded grasslands (semi-humid savanna), woodlands, dry forests, in riverbeds.

Remarks. This species is rather cryptic and imperfectly known. The leaves seem to develop conspicuous margin teeth during maturity, like e.g., *C. grandis*, but the darkish sublaminal glands differ from that species. The erect calyx lobes with slightly




Figure 31. Distribution map of *C. intermedia* (pale yellow dots; based on 4 collections), *C. keayana* (blue dots; based on 23 collections), and *C. longicarpa* (blue triangles; based on 22 collections).

recurved tips appear to be the most indicative character for *C. intermedia*. The clustered female flowers and the fruits link to *C. barteri*, from which it, among other characters, differs in ecology. Two *J.B. Hall & J.M. Lock GC 46016* specimens from K have male and female flowers/fruits on one twig and are thus monoecious. As all other *Coccinia intermedia* collections are dioecious, this could be a case of “leaky dioecy” (Baker and Cox 1984), which also has been observed in other Cucurbitaceae (Schaefer and Renner 2010).

9. *Coccinia keayana* R.Fern., *Bol. Soc. Brot. ser. 2*, 33: 191. 1959.

Type: Guinea-Bissau. Tombali: region Cacine, rainforest, fl, Aug 1933, *J.V.G. do Espírito Santo* 603 (Holotype: COI).

Type: *ibid.*, *J.V.G. do Espírito Santo* 631 (Paratypes: LISC [LISC 011640, digital image: IICT, JPS], LISC! [LISC 011641, digital image: IICT, JPS], LISC [LISC 011642, digital image: IICT, JPS], LISJC).

Type: Guinea-Bissau. Tombali: between Cacine and Guileje, plantation, 1 Aug 1945, *J.V.G. do Espírito Santo* 2151 (Paratypes: COI, LISC [LISC 002508, digital image: IICT, JPS], LISC! [LISC 002509, digital image: IICT, JPS]).

Type: Liberia. [Margibi County]: Firestone plantation, at Du River, 29 Jul 1926, *D.H. Linder* 121 (Paratype: K!).

Type: Sierra Leone. Jigaya, c. 350 m, 28 Sep 1914, *W. Thomas* 2844 (Paratype: K!).

Type: Sierra Leone. [Northern Province]: Bumban National Park, 30 Aug 1928, *F.C. Deighton* 1221 (Paratype: K!).

Type: Sierra Leone. [Southern Province]: Moyamba District, Moyamba, 25 Aug 1931, *F.C. Deighton* 2217 (Paratype: K!).

Coccinia sp. A Keay, *Fl. W. Trop. Afr.* 1: 216. 1954.



Figure 32. Reconstruction of the habit of *Coccinia keayana* based on C.C.H. Jongkind *et al.* 6542 (WAG). Black bar equals 1 cm.

Description. Perennial? climber. Stem up to 5 m, glabrous. Petiole 1.5–5 cm, with short, few-cellular trichomes on adaxial side, glabrous on abaxial side. Leaves 5–11 × 3.5–11 cm, (shallowly to) profoundly 3-(or 5-)lobate, auriculate, rarely long cordate. Margin rather remotely dentate to slightly serrate. Lobe apex acute or subacute with final tooth. Upper leaf surface tiny hyaline pustulate. Lower leaf surface with blackish glands, dried often with bluish-green tinge, glabrous or rarely with soft multicellular trichomes on nerves. Probracts up to 3 mm. Tendrils simple. Male flowers ebracteate, in lax racemes with up to 20 flowers, sometimes accompanied by a solitary flower (Fig. 32). Common peduncle up to 1.7 cm, shorter than racemous part, glabrous. Pedicels of racemous flowers up to 1 cm, pedicels of solitary flowers up to 1.5 cm long, each glabrous. Perianth tube glabrous. Calyx lobes linear, 2.5–3 mm, in buds spreading, later reflexed. Corolla 1.7–2 cm long, white, yellow, dirty orange, salmon to dull pinkish. Corolla lobes 3–5 mm long. Filament column and anther head not seen, pollen sacs pink–orange. Female flowers solitary or in few-flowered lax racemes. Common peduncle 1.2–2.1 cm, glabrous. Pedicel flowers in racemes up to 1 cm, glabrous, pedicels of solitary flowers up to 2.7 cm, glabrous. Hypanthium glabrous, calyx lobes and corolla as in male flowers. Ovary glabrous. Style and stigma not seen. Fruits 2–3 × 2 cm, subglobose to globose, un-

ripe glaucous green, ripening via yellow to pinkish [rather glaucous?] red. Seeds $4.5 \times 2.5 \times ?$ mm (L/W/H), rather symmetrically obovate, face flatly lenticular.

Phenology. Flowering time: March to November.

Distribution. Fig. 31. Tropical West Africa: Guinea-Bissau (Tombali), Sierra Leone, Liberia, Guinea (Nzérékoré), S Ivory Coast, S Ghana, S Togo? Elevation sea level to 1250 m. On sandy soil, lateritic soils. Tropical rainforest, in high trees, high bushes, on roadsides.

Vernacular names. Cf. Koranko: nala (*W. Thomas 2844*), Limba: ngolibwe (*W. Thomas 2844*), Mende: ndogbo-gojai (*F.C. Deighton 2217*), Temne: efosa (*W. Thomas 2844*).

Remarks. The long racemes with ebracteate flowers and the linear, reflexed calyx lobes are good characters for this species. It is barely distinguishable from *C. barteri* without flowers. *Coccinia keayana* collections often have a bluish green tinge and the lobes conspicuously point forwards (see Fig. 32), which might only rarely occur in *C. barteri*. The corolla is rather tubular, sometimes somewhat inflated.

Taxonomic remarks. The name *C. keayana* R.Fern. is misapplied for the Flora of Cameroon or western Central Africa in general (for details, see taxonomical remarks of *Coccinia barteri*).

Specimens examined. (Selection; in total: 33) Ghana. Eastern Region: Asiakwa district, Atewa Range Forest Reserve, Accra-Kumasi highway 5–6 km along forest road that intersects the highway at Sagyimase village, $6^{\circ}13'48''\text{N}$, $0^{\circ}32'42''\text{W}$, *M. Merello et al. 1179* (MO). Volta: Amedzofe, *J.B. Hall GC40053* (P [P05620653]). Western Region: Bia National Park, *J.B. Hall & J.M. Lock GC46493* (WAG [WAG0046501]). Guinea. Nzérékoré: Nzérékoré Préfecture, [WSW of] Nzérékoré, $7^{\circ}43'35.54''\text{N}$, $8^{\circ}51'21.28''\text{W}$, *E. Achigan-Dako 07 NIA 917* (GAT). Guinea-Bissau. Tombali: Bedanda sector, Cantanhez, *J. Alves Pereira 3172* (H, LISC). Ivory Coast. Lacs: Oroumba Boca [Orombo Bocca Mt.], *H.C.D. de Wit 5772* (WAG [WAG0234139], WAG [WAG0234140]). Lagunes: Abidjan, Banco Forest Reserve, *W.J.J.O. de Wilde 893* (BR, EA, WAG [WAG0044624], WAG [WAG0044625], Z). Liberia. Bong: 3 miles NE of Suacoco, *Z.D. Traub 256* (BR, G, MO). Nimba: Yéképa, Mt Nimba, *J.-G. Adam 21232* (MO, P [P00694038/P05590407], P [P05620652], PRE).

10. *Coccinia longicarpa* Jongkind, *Blumea* 49: 83. 2004.

Type: Ivory Coast. Lagunes: Forêt du Banco, S of Arboretum, near river, male and female, fr, 20 Jul 1973, *J. de Koning 1965* (Holotype: WAG! [WAG0099441, digital image: JPS], isotype: K! [K000035540, digital image: K]).

Type: Ivory Coast. Lagunes: Forêt du Banco, Route Martineau, secondary forest, 10 Oct 1974, *J. de Koning 4077* (Paratype: WAG! [WAG0099430]).

Type: Ivory Coast. Lagunes: Forêt du Banco, near swampy secondary forest, 8 Aug 1975, *J. de Koning 5904* (Paratypes: K [K000035541], MO!, WAG! [WAG0099438, digital image: JPS]).

Type: Ivory Coast. Lagunes: Forêt du Banco, N of center, near Banco river, in forest clearing on clear spot, 16 Jun 1975, *W.J. van der Burg* 551 (Paratype: WAG! [WAG0062627]).

Type: Ivory Coast. Lagunes: near Abidjan, 6 Sep 1967, *C. Geerling & J. Bokdam* 829 (Paratypes: WAG! [WAG0011282], WAG! [WAG0011283]).

Type: Ivory Coast. Lagunes: [W of Abidjan], Adiopodomé [Adiopo-Doumé], margin of bush pathway, 3 Aug 1956, *J.J.F.E. de Wilde* 183 (Paratypes: WAG! [WAG0044616], WAG! [WAG0044617], WAG! [WAG0044618], WAG! [WAG0044619]).

Type: Ghana. Eastern Region: near Kibi, Atewa Range Forest Reserve, Jun 1976, *J.M. Lock* GC 43991 (Paratype: K).

Type: Liberia. Gbapolu/Lofa: Gbanga, Sep 1926, *D.H. Linder* 576 (Paratype: K).

Type: Ghana. Nsuta, no detailed location, 1500 ft, May 1929, *C. Vigne* 1735 (Paratypes: K, P!).

Coccinia sp. A C. Jeffrey, Key to the Cucurbitaceae of West Tropical Africa. J. W. African Sci. Assoc. 9: 87 p.p., 1964.

Coccinia sp. B Keay, Fl. W. Trop. Afr. 1(1): 216 p.p., 1954. Nigeria. [Ogun]: Ijebu-Ode District, toward the head of extraction Rd. from Grace Camp [in Omo Forest Reserve] westwards toward the R. Omo [Shasha], about 1.5 mls from the Grace Camp, 2 May 1946, *S. Tamajong* FHI 16938 (K!); *D.H. Linder* 576 (K); *C. Vigne* 1735 (K, P!).

Coccinia sp. D Keay, Fl. W. Trop. Afr. 1(1): 216 p.p., 1954. Nigeria. [Oyo]: Lagos [Colony], Ibadan forest, 1 Dec 1900, *C. Punch* 46 (K!).

Description. Perennial climber. Stems up to 5 m, glabrous. Petiole 0.9–4.5 cm, glabrous. Leaves 7–12 cm × 2.5–11 cm, 5-angularly subcordate, subhastate, rarely 3-lobate, auriculate. Lobes elliptical. Margin suspiciously dentate, whitish in living state, blackening when dried. Apex acute with final tooth. Upper leaf surface pale to white pustulate. Lower leaf surface glabrous, nerves often white-speckled. Probracts < 1 mm, often missing. Tendrils simple, rarely bifid. Male flowers ebracteate, in glabrous racemes, occasionally accompanied a solitary flower. Common peduncle 0.7–1.5 cm, pedicel of flowers in raceme 2.5–8 mm, pedicel of solitary flower 5–10 mm. Perianth tube glabrous, calyx lobes 2–3 mm long, (0.75–)1.2–1.8 mm broad at base, erect. Corolla 0.8–1.5 cm long, yellow to (pale) orange, sometimes with greenish nerves. Corolla lobes up to 2 mm long. Filament column and anther head not seen, pollen sacs yellow to orange. Female flower solitary. Petiole 1.5–2.2 cm, glabrous. Hypanthium glabrous, calyx lobes and corolla not seen, but likely as in male plants. Ovary glabrous. Style and stigmas not seen. Fruit up to 20 cm long, c. 1 cm in diam., long cylindrical, unripe waxy green, ripe unknown. Seed size and shape unknown, face flat.

Phenology. Flowering time: May–December.

Distribution. Fig. 31. Tropical West Africa: SW Ivory Coast, S Ghana, SE Benin, SW Nigeria. Elevation sea level to 460 m. Soil preference not known. In primary or secondary forests, in disturbed places (roadsides, near rivers).

Remarks. The broad calyx lobes are, apart from the long cylindrical fruit, the best character for identifying this species. An urceolate corolla (Jongkind 2004) also occurs in *C. barteri*.

A single collection (*H.J. Beentje* 602 from M) mentions a lilac corolla color, which would be unique in *Coccinia*. Although this might be possible, since there are also pinkish flowers reported in *C. adoensis*, the fact that the WAG duplicate with the same collection number is a *Ruthalicia* makes it more likely that the observation is due to a mixed collection, eventually from a Convolvulaceae.

Specimens examined. (Selection; in total: 24) Benin. Plateau: Pobe, 6°57'52.56"N, 2°40'19.70"E, *E. Achigan-Dako* 07 NIA 731 (GAT). Ghana. Ashanti: Bobiri Forest Reserve, 6°41'N, 1°21'W, *C.C.H. Jongkind* 3970 (WAG [WAG0023747], WAG [WAG0023748]). Central Region: Twifo/Hemang/Lower Denkyira District, Kakum, 5°20'54.31"N, 1°23'1.39"W, *E. Achigan-Dako* 07 NIA 734 (GAT). Eastern Region: Atewa Range Forest Reserve, 06°15'N, 00°33'W, *C.C.H. Jongkind et al.* 1538 (MO, WAG [WAG0020265]). Western Region: Jomoro District, Fawoman, 5°19'32.63"N, 2°43'28.13"W, *E. Achigan-Dako* 06 NIA 050 (GAT); *ibid.*, *E. Achigan-Dako* 06 NIA 051 (GAT). Ivory Coast. Lagunes: Abidjan. Banco Forest Reserve, *J.J. Wieringa* 5386 (WAG [WAG015997]). Sud-Comoé: Aboisso, 5 km NNW of Nganda-Nganda [Forest Reserve], near lagoon Aghien [lagoon Aby?, because lagoon Aguien is NE of Abidjan], 5°14'N, 3°20'W, *H.J. Beentje* 602 (M, but not WAG). Nigeria. Ogun: Ijebu-Ode District, Omo Forest Reserve, Compartment 8, *J.R. Charter FHI* 38635 (K).

11. *Coccinia mackenii* Naudin ex C.Huber [sphalm. Mac-Kennii, after John M'Ken, ICN 60C.5], Cat. Print.: 5. 1865.

Cephalandra mackenii (Naudin ex C.Huber) Naudin [sphalm. mac kennii], Ann. Sci. Nat. Bot. 5: 17, ser. 5. 1866.

Type: Cultivated. Cultivated in Paris Botanical Garden from seeds from Huber's Garden in Olbia [Hyères, France] who obtained the seeds from M'Ken from near Port Natal [Durban, KwaZulu-Natal, South Africa], female, fl, 1864, *C.V. Naudin s.n.* (Syntypes: P!, G-DC! [G00211343, digital image: G]), P [P06745731, digital image: P], P [P06745732, digital image: P].

Type: Cultivated. Cultivated in Paris Botanical Garden from seeds from Huber's Garden in Olbia [Hyères, France] who obtained the seeds from M'Ken from near Port Natal [Durban, KwaZulu-Natal, South Africa], 1863, *C.V. Naudin s.n.* (Syntypes: P! [P06745737, digital image: P], P! [P06745739, digital image: P], P [P06745740, digital image: P]).

Type: Cultivated. Cultivated in Huber's Garden in Olbia [Hyères, France] who obtained the seeds from M'Ken from near Port Natal [Durban, KwaZulu-Natal, South Africa], male and female, fl, 1864, *C.V. Naudin s.n.* (Lectotype, designated here: P! [P06745735, digital image: P; K neg. 2993]; isoelectotypes: G-DC! [3

sheets, all G00211344, digital images: G], K! [K000542637, digital image: K], K! [K000542638, digital image: K], K (2)!, P! [P06745730, digital image: P], P! [P06745733, digital image: P]).

Coccinia palmata (Sond.) Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 540. 1881. Nom. illeg. *Momordica palmata* E.Mey. ex Drège, Zwei pflanzengeogr. Dokum. [addition to Flora 26(2)]: 156, 159, 202. 1843. Nom. nud. *Cephalandra palmata* Sond. in Harv. & Sond., Fl. Cap. 2: 493. 1862.

Type: [South Africa]. [KwaZulu-Natal]: near Port Natal [Durban; cited in l.c. p. 159], male and female, fl, fr, 7 Apr 1832, *J.F. Drège s.n.* (Lectotype, designated by Meeuse (1962: 96): S! [S08-12155, digital image: S], isolectotype: P! [P00748835, digital image: P]).

Type: [South Africa]. Without location and date, male, fl, *J.F. Drège s.n.* (Syntypes: G! [G00226835, digital image: G], HBG! [HBG506427], K! [K000313198, digital image: K], K! [K000313199, digital image: K], L!, P! [P00748833, digital image: P], P!, PRC!, W! [W 0026939, digital image: WU]).

Type: [South Africa]. [KwaZulu-Natal]: Omsamculo [Umzimkulu], between shrubs and thickets, near river mouth, female, fr, 5 Mar 1832, *J.F. Drège 4637* (Syntype: P! [P00748834, digital image: P])

Coccinia dinteri André, Rev. hort. [Paris] 72: 276. 1900.

Type: Unnumbered plate in l.c.

Description. Perennial climber or creeper. Stems up to 9.5 m, glabrous. Petioles 0.7–11 cm long, glabrous or with thin trichomes. Leaves 3–13.5 × 3–15.5 cm, shallowly to profoundly 5-lobate, in the latter case often weakly lobulate. Lobes triangulate, lanceolate, ovate to obovate. Margin smooth, dentate, sometimes serrate to lobulate, esp. towards the apex. Apex acute with final tooth. Upper leaf surface glabrous with clear to white pustules, rarely with few trichomes. Lower leaf surface glabrous or with thin, stiff or articulate trichomes, towards base usually with glands. Probracts up to 4 mm, oblong-lanceolate. Tendrils bifid, rarely simple. Male flowers solitary or ebracteate in few-flowered racemes. Common peduncle 5–6.5 cm, pedicel of flower in raceme up to 2.5 cm, pedicel of solitary flowers 6–9 cm, all glabrous, rarely with long trichomes. Perianth tube glabrous. Calyx lobes 1.5–6.5 mm, lineal, subulate to narrowly triangulate, when young erect, later spreading to reflexed. Corolla 1.3–2.7 cm long, cream to pale buff, corolla lobes subulate to triangulate, 0.7–1.1 cm. Filament column, anther head, and pollen sacs not seen. Female flowers one solitary. Pedicel 0.7–5 cm long, glabrous. Hypanthium glabrous, calyx and corolla like in male flowers. Ovary glabrous. Style columnar, color not seen. Stigma bulging, color not seen. Fruits elliptical to oblong, c. 10 × 2–2.5 cm. Unripe green with white mottling, ripe red-orange to red, sometimes? with white mottling. Seeds 6–7 × 4–4.5 × 1.5 mm (L/W/H), slightly asymmetrically obovate, face flatly lenticular.

Phenology. Flowering time: January–April, July, November, December.

Distribution. Fig. 30. South Africa (E Eastern Cape, KwaZulu-Natal, Mpumalanga, Limpopo), Swaziland. Elevation sea level to 1750 m. Clay, Berea red sand,

sandstone, quartzite, poorly drained soils. Afromontane forests, coastal forests, littoral forests, forest margins, sometimes grassland. In frost-free areas (Meeuse 1962).

Use. Leaves and fruits are eaten by Tsonga people (Shackleton et al. 1998).

Vernacular names. Xitsonga: Gomo, XipapaXipapana (Shackleton et al. 1998).

Remarks. Some collections with deeply lobate leaves and short petioles resemble the closely related *C. quinqueloba*, and some *C. quinqueloba* individuals have long petioles (*C.V. Naudin s.n.* 1863, *C.V. Naudin s.n.* 1863–1865, *E. Retief* 1215). However, Naudin (1866) reports considerable problems with seed production in interspecific crosses. It would be desirable to validate this observation.

Taxonomic remarks. The initial designation of the *C. mackenii* lectotype (Holstein and Renner 2010: 440) is not valid, because it erroneously designated a female specimen from Paris Botanical Garden. However, Naudin stated that all plants from Paris Botanical Garden were male (P06745733), so the former designation was ambiguous. The new lectotype was chosen from Olbia [Hyères] material. In contrast to Naudin's statement, that Olbia material was female, there is a male K specimen (K000542638), from Huber's Garden in Olbia. Eventually, this specimen is from Paris Botanical Garden but incorrectly labeled, because the lack of the opposite sex affected Naudin's crossing experiments.

Due to an overlooked published combination *C. mackenii* bore the illegitimate name *C. palmata* for more than 120 years. When Wight and Arnott published *Coccinia indica* they also included a specimen tentatively identified as *Bryonia palmata* L. Although without relevance for the genus *Coccinia* itself, it led to further complications. One year after Voigt's publication of the correct combination *Coccinia grandis*, Roemer (1846) also recognized the apparently missing combination and that Linnaeus' *B. palmata* and *B. grandis* indeed referred to different species. Roemer treated them, amongst other species, as *C. grandis* (L.) M.Roem. (nom. illeg.) and *C. palmata* (L.) M.Roem. In addition to the name *Coccinia palmata* (L.) M.Roem. another species from South Africa was described with the name *Cephalandra palmata* E.Mey. ex Sond. (Harvey and Sonder 1862). Cogniaux (1881) accepted this species in *Coccinia*, overlooking *Coccinia palmata* (L.) M.Roem. He thus created an illegitimate *Coccinia palmata* (E.Mey. ex Sond.) Cogn., which has since been used for this species. Holstein and Renner (2010) called attention to this erroneous usage by resurrecting the correct name, *Coccinia mackenii* Naudin ex C.Huber.

The drawing of *Coccinia dinteri* in the protologue shows a bifid tendril. Since all other characters match *C. mackenii* and the resemblance was already discussed in the protologue, it is feasible to synonymize it with that species. M. Proschowsky grew this plant in the Fabron quarter of Nice, France, but the origin of the seeds was not indicated. The label named it "*Coccinia dinteri*" after Moritz Kurt Dinter (in the protologue erroneously spelled as "Hurt Dinter"), who was curator in La Mortola (Giardini Botanici Hanbury, Liguria, Italy) where many South African plants were cultivated. Hence, it is reasonable to assume this origin as done by André there, which again would match *C. mackenii*. There is a specimen in K herbarium containing only seeds and a label indicating that they were sent from Hanbury, La Mortola in 1897. A note mentions that the seeds were sown in Kew Gardens. The identification is given as *Cephalan-*

dra mackenii with a question mark and a later note with the *Coccinia dinteri* citation. It is plausible to assume that these seeds come from the same plant stock that was used to grow and to describe *C. dinteri*. Although the seeds fit the description of *C. mackenii* seeds, it is not possible to use them to identify the species unambiguously.

Specimens examined. (Selection, in total: 71) South Africa. Eastern Cape: Port St. Johns, Jan 1933, *A.O.D. Mogg s.n.* (L, PRE [PRE 42990], Z). Kwazulu-Natal: Durban district, Isipingo North, *C.J. Ward 3747* (COI, PRE); Umzinto district, Vernon Crookes Nature Reserve, far end of Golden Valley, *K. Balkwill et al. 10930* (E [E00264193], MO); Pietermaritzburg, Ferncliffe Forest, *J. Bodenstein 92* (PRE); Nkandlha [Nkandla], Qudeni Forest, 5 mls [8 km] S of Qudeni P.O., *L.E.W. Codd 6991* (PRE). Limpopo: near Lydenburg, near Echo Cave, *R.G. Strey 3762* (M, PRE, WAG [WAG0234163]); Woodbush [Forest Reserve], 6 Aug 1925, *A.O.D. Mogg s.n.* (COI, L, PRE [PRE 43066], Z). Mpumalanga: Letaba district, E side of shoulder extending northwards from ridge above Weltevreden, *J.C. Scheepers 1110* (M, PRE). Swaziland. Hhohho: about 20 km N of Mbabane, Ngwenya Hills, Castle peak, north slopes, *B. Maguire 7553* (B [B 10 0019799], E). Manzini: Usutu Forests, *R.H. Compton 32287* (PRE).

12. *Coccinia megarrhiza* C.Jeffrey, Kew Bull. 15(3): 347. 1962.

Type: Kenya. Northern Province [Eastern Province]: Moyale, 3800 ft, male, fl, 28 Apr 1952, *J.B. Gillett 12967* (Holotype: K! [K000313235, digital image: JPS, K], isotypes: B! [B 10 0154926, digital image: B, JPS], S! [S08-12479]).

Type: Kenya. Northern Province [North Eastern Province]: [western Mandera District], Dandu, 2600 ft, 10 Apr 1952, *J.B. Gillett 12759* (Paratype: K! [K000354130]).

Type: Kenya. Northern Province [North Eastern Province]: [western Mandera District], Dandu, 2700 ft, 9 May 1952, *J.B. Gillett 13122* (Paratypes: EA!, K! [K000354129]).

Type: Kenya. Northern Province [North Eastern Province]: [western Mandera District], Dandu, 3000 ft, 14 May 1952, *J.B. Gillett 13191* (Paratypes: EA!, K!).

Type: Kenya. Northern Province [Eastern Province]: Moyale, 3200 ft, 3 Oct 1952, *J.B. Gillett 13986* (Paratype: K! [K000354128]).

Type: Kenya. Northern Province [Eastern Province]: Moyale, 3600 ft, 14 Oct 1952, *J.B. Gillett 14036* (Paratypes: B! [B 10 0154927, digital image: B, JPS], BR!, EA!, K! [K000354131], S! [S08-12482]).

Type: Kenya. *ibid.*, *J.B. Gillett 14037* (Paratypes: BR! [BR0000008914033], K! [K000354133], K! [K000354134], S! [S08-12481]).

Type: Kenya. *ibid.*, *J.B. Gillett 14038* (Paratypes: K! [K000354125], K! [K000354132]).

Type: Kenya. *ibid.*, *J.B. Gillett 14039* (Paratypes: K! [K000354126], S! [S08-12480]).

Description. Perennial climber or creeper. Stem up to 6 m, with long, whitish to beigeish patent trichomes, which appear articulate when dried. Petioles 1.5–5.6 cm, indumentum as on stem (Fig. 9). Leaves 3–11 × 4.2–17 cm, reniform to 3- or 5-lobate. Margin dentate (teeth at maturity brownish, when dried blackening),

serrate to lobulate. Upper leaf surface glabrous with pale pustules or with short, whitish to beigeish trichomes, lower leaf surface with indumentum as on stem, rarely glabrous. Probracts up to 3 mm long. Tendrils simple. Male flowers clustered. Pedicel < 1.5 cm, indumentum as on stem. Perianth tube with long, beigeish, upright trichomes that appear articulate when dried. Calyx lobes 2.5–4 mm, subulate to lineal, spreading. Corolla 1.2–1.3 cm, yellow to pale orange, lobes 4–6 mm. Filament column greenish, anther head pale greenish, pollen sacs orange-yellow. Female flowers 1(–2) solitary. Hypanthium with long, beigeish, upright trichomes that appear articulate when dried, calyx lobes and corolla like in male flowers. Ovary green with whitish spots. Style columnar, green. Stigma bulging, yellow (Fig. 11c). Fruit ovoid–ellipsoid, up to 6.5 cm long, unripe green with longitudinal white mottling. During ripening mottling partly developing a dark green corona (Fig. 13c). Ripe red (Fig. 12b). Seeds $7 \times 4 \times 1.5$ mm (L/W/H), asymmetrically obovate, face flat.

Phenology. Flowering time: March–May, August–October.

Distribution. Fig. 20. Ethiopia (Oromia, Somali Region), Kenya (Eastern Province, North Eastern Province). Elevation 800–1600[–2000] m. On granite outcrops and red lateritic soils, *Acacia-Commiphora* bushland, *Dracaena* vegetation, *Balanites* vegetation, *Euphorbia candelabrum* woodlands, dry *Juniperus* forest, grazing is tolerated.

Remarks. *Coccinia megarrhiza* and *C. abyssinica* form a species complex. Distinction between these two species can be difficult, especially in young plants, when the color of the marginal teeth of the leaf is not well developed. While the peduncle length differs, the earlier appearing solitary flowers can have the same length in both species. The broad leaves with an emarginate, obtuse to cuspidate tip (*C. megarrhiza*) versus rather long leaves with an acute tip (*C. abyssinica*) seems to be the best character. At maturity, the teeth coloration in *C. megarrhiza* is also much more conspicuous than in *C. abyssinica*. A phylogeographic analysis and crossing experiments would shed light on the question, whether these are ecologically differentiated forms or true species. Plants from the mountains near Yebelo with very large leaves are almost glabrous and occur, untypically, in dry *Juniper* “forests”. However, they have the typical cuspidate to obtuse central lobes and bear the colored leaf margin teeth. As larger leaves are also observed in high altitude individuals of *C. microphylla*, these forms might be regarded as mast specimens.

Specimens examined. (Selection; in total: 29) Ethiopia. Oromia: 38 km S of Neghelli [Negele Boran] on Wachelli road, *J.W. Ash* 814 (EA (2), K); Arero (Meta-Gafersa), *G. Cufodontis* 273 (FT, W); Bombal ca. 40 km on the way to Jijiga from Harar, *T. Ebba* 622 (K, WAG [WAG0285707]); c. 20 km NW of Moyale on the road to Mega, just after the turn off to Tuqa (and Sololo in Kenya) (3°39'N, 38°56'E), *I. Friis et al.* 8736 (K); c. 36 km from Harar to Jijiga and then c. 20 km to S, *J.J.F.E. de Wilde* 4793 (B, K, MO, WAG [WAG0285708], WAG [WAG0285709]). Somali Region: 95 km from Negele of Filtu road, 5°00'N, 40°12'E, *M.G. Gilbert & B.M.G. Jones* 110 (K).

13. *Coccinia microphylla* Gilg, Bot. Jahrb. Syst. 34: 357. 1904.

Type: Tanzania. [Kilimanjaro]: at base of Pare Mountains, between Kiswani [Kisiwani] and Maji ya Juu [Madji-ja-juu], mix of thornbush and wooded grassland [“gemischte Dornbusch- und Obstgartensteppe”], 700 m, fl, Oct, *A. Engler, Reise nach Ostafrika 1587* (Syntype: B destroyed).

Type: Tanzania. [Kilimanjaro]: between Kihuiro [Kihurio] and Gonja, thornbush steppe, fl, Oct, *A. Engler, Reise nach Ostafrika 948* (Syntype: B destroyed).

Type: Kenya. Coast Province: near Mariakani, NW of Mombasa, male, fl, 15 Oct 1955, *E. Milne-Redhead & P. Taylor 7104* (Neotype, designated here: LISU!, isoneotypes: B!, EA!)

Coccinia buikoensis Zimm., *Die Cucurbitaceen* 2: 177, 24, 51, 84, 96, 114, fig. 17 I–III, fig. 63 II, fig. 74 VII–XII, fig. 81 XVI, XVII. 1922.

Type: Tanzania. [Tanga]: Lushoto District, [S of Pare Mts], [between Hedaru and Mkomazi], near Buiko, steppe, male and female, fl, fr, Dec 1919, *P.W.A. Zimmermann G6595* (Holotype: B destroyed, lectotype designated by Jeffrey (1967: 68): EA [EA000002132, digital image: JPS]).

Coccinia sp. C in C. Jeffrey, F. T. E. A.: 69. 1967. Kenya, Northern Province: Furroli, lava plateau, semi-desert, *Acacia-Commiphora* shrub, on sand, female, fl, fr, 12 Sep 1952, *J.B. Gillett 13820* (B!, EA!, K!, P!, S! [S08-12180]) and *J.B. Gillett 13826* (K!).

Description. Perennial creeper or climber. Stems up to 4 m, glabrous or more or less densely covered with short, white trichomes, when older often densely white pustulate. Petiole 0.45–4 cm, with erect, often thick, when longer sometimes bent trichomes that are sometimes soft spiny (< 1 mm) or only wart-like. Leaves 0.7–7.5 × 1.1–12 cm wide, usually rather small, shallowly to deeply 3- or 5-lobate, sometimes lobulate, rarely reniform. Lobes narrow to broadly triangulate to lanceolate. Upper leaf surface more or less densely white pustulate, pustules sometimes with a short, thick trichome (Fig. 2a). Lower leaf surface glabrous or more or less dense with often stiff, pale yellowish trichomes, sometimes with darker glands between nerves, nerves usually with thick, erect trichomes, sometimes reduced to wart-like appearance. Leaf margin rather remotely denticulate, usually with minute, bent trichomes. Apex acute to obtuse, with final tooth. Probract usually absent, if present then < 1.5 mm. Tendrils simple. Male flowers 1–2 solitary, if fasciculate or in few-flowered racemes, then accompanied by 1 solitary flower. Common peduncle < 5 mm, glabrous or with minute to long, articulate trichomes. Pedicel of solitary flowers 2–7(–25 mm), pedicel of flowers in inflorescences < 5 mm, glabrous or especially at apex with long, yellowish, articulate trichomes. Perianth tube usually with long, yellowish, articulate trichomes, rarely almost glabrous. Calyx lobes 1.5–4 mm, narrowly lanceolate to lineal, when young erect, later spreading to reflexed. Corolla 0.7–1.2 cm long, greenish white to yellowish-orange with darker green veins, lobes 4–7 mm. Filament column and anther head pale greenish yellow. Pollen sacs yellow. Female flowers 1(–2) solitary (Fig. 2a). Pedicels up to 0.7 cm, glabrous or with white trichomes. Ovary glabrous, with some articulate trichomes to densely wooly




Figure 33. Distribution map of *C. microphylla* (blue dots; based on 49 collections) and *C. trilobata* (pale yellow dots; based on 51 collections).

with long (when dry articulate) trichomes. Style columnar, pale green. Stigmas bulging, greenish yellow. Ripe fruit globose to shortly obovate, 1.8–2.5 × 1.4–2.5 cm, glabrous or with few articulate trichomes, unripe green sometimes with longitudinal, whitish

mottling (Fig. 2a), which obtains a dark green corona during ripening, when ripe bright orange-red to red. Seeds $4.5\text{--}6 \times 2\text{--}3 \times 1\text{--}1.5$ mm (L/W/H), asymmetrically obovate to somewhat falcate, face flattened.

Phenology. Flowering time: January, April, May, July, October–December.

Distribution. Fig. 33. NE Tanzania (Arusha, Kilimanjaro, Manyara, Tanga), Kenya (Coast, Eastern, North Eastern, Rift Valley), S Ethiopia (S and C Oromia), Somalia. 70–1300(–1600) m. Savanna, open *Acacia-Commiphora* bushland, degraded *Combretum* bushland, open grassland, cultivated land, roadsides. Red sand, dark brownish-black soil. Limestone.

Vernacular names. [Akiek; Ogiek]: notoku (*A.S. Vincent 29, A.S. Vincent 221*), Maa [Maasai language]: ndegegeya (*A.S. Vincent 29*), sikuni (*Kiamba et al. KEFRI 112*).

Remarks. Some collections have a mixed (not intermediate) phenotype with *C. trilobata*: the calyx lobes are unusually long (up to 7 mm), which speaks for *C. trilobata*, but the indumentum matches *C. microphylla*. However, these do not occur in a single location, but are found in the Ndoto Mts (*O. Kerfoot 2644*), in Kiboko (*P. Kirika et al. 002/020/2011*), and around Voi (*M. Hucks 579, B. Verdcourt 3888*). Whether these are hybrids (F2 or later) or just a variety is not known. These collections also resemble *C. megarrhiza*, which occurs in northern Kenya and Ethiopia, however, the indumentum does not match either.

Despite the epithet, the leaves can become quite large, especially at higher altitudes. Then, collections may resemble *C. trilobata*, which has a denser indumentum. In dry low altitude areas, leaves and flowers emerge quickly, e.g., soon after a rain shower. The leaves are thus not well developed and small. Collections from more arid locations tend to be smaller in many characters, but whether these represent a new species is doubtful. There are only few collections of the proposed species (*C. sp. Burger 2947A, C. sp. Gilbert & Jones 129* (Jeffrey 1995)), but the natural scope of *C. microphylla* is hardly assessed. This will not be resolved without a phylogeographic analysis and more intensive collecting from Ethiopia.

Specimens examined. (Selection, in total: 72) Ethiopia. Oromia: 105 km on road from Negelli [Negele] to Filtu, *J.J.F.E. de Wilde & M.G. Gilbert 346* (K, WAG [WAG0285710], WAG [WAG0285711], UPS). Kenya. Coast Province: near Mariakani, NW of Mombasa, *E. Milne-Redhead & P. Taylor 7105* (B, EA, LISU). Eastern Province: E side of Lake Rudolf, between Koobi Fora and Shin (hill), $3^{\circ}57'\text{--}58'\text{N}$, $36^{\circ}12'\text{--}20'\text{E}$, *R.B. Faden & A. Evans 71/301* (EA, K). North Eastern Province: Wajir District, Catholic Girl's Town 2 km E of Wajir, *J.B. Gillett 21273* (EA, K, WAG [WAG0234120]). Rift Valley Province: Turkana District, *I. Ohta 24* (EA). Somalia. Togdheer: Malol [Mt Malool] near Sheikh, *J.R. Ironside 5/73/31* (K). Tanzania. Arusha: Monduli district, Longido division, SE of Longido, c. 100–300 m from Arusha Municipality, $2^{\circ}43'14''\text{S}$, $36^{\circ}42'02''\text{E}$, *C.J. Kayombo & K. Kitaba 4242* (MO). Dodoma: Tarangire National Park, Kalima Hill, *S. Chuwa et al. 5329* (NHT). Kilimanjaro: Same district, Mkomazi Game reserve, Ibaya Hill, $3^{\circ}58'\text{S}$, $37^{\circ}48'\text{E}$, *R. Abdallah & K. Vollesen 95/198* (BR, K, P [P05620649]). Tanga: Korogwe District, 2 km W of Mkomazi, under power line, $4^{\circ}38'53.7''\text{S}$, $38^{\circ}03'26.7''\text{E}$, *N. Holstein et al. 90* (B, DSM, M).




Figure 34. Distribution map of *C. mildbraedii* (based on 51 collections). The individual in Kenya is introduced. For Tanzania the borders of the regions are given.

14. *Coccinia mildbraedii* Gilg, Wissenschaft. Ergebn. Deutsch. Zentral-Afrika-Exped. 1907–1908 Herzog Adolf Friedrich zu Mecklenburg, Bot. 2(4): 343. 1914.

Type: Rwanda. [Western Province]: Kissenye [Gisenyi]. Bugoy [Bugoyi] forest, mixed bamboo forest, c. 2500 m, fl, fr, 30 Oct 1907, *J. Mildbraed* 1425 (Holotype: B, destroyed).

Type: Burundi. Muramvya: [Mt] Teza, 3°13'S, 29°33'E, female, fl, fr, *M. Reekmans* 7399 (Neotype, designated here: K!; isoneotypes: BR!, EA!, MO!, WAG! [WAG0225430], WAG! [WAG0225433]).

Coccinia ulugurensis Harms in Mildbraed, Notizbl. Bot. Gart. Berlin-Dahlem 11: 1091. 1934.

Type: Tanzania. [Morogoro]: Uluguru Mts, northwestern side, c. 1350 m, over shrubs at forest margin, male, fl, 14 Mar 1933, *H.J.E. Schlieben* 3643 (Holotype: B! [B

10 0154929, digital image: B, JPS], isotypes: B! [B 10 0154930, digital image: B, JPS], BM! [BM000815208], BR! [BR0000008886163, K neg. 5264, digital image: JPS], BR! [BR0000008887498, digital image: BR, JPS], G! [G00301594], HBG! [HBG506425, digital image: JPS], LISC! [LISC 002496, digital image: IICT, JPS], M! [M0105771, digital image: JPS], MA! [MA386121, digital image: JPS], MO!, P! [P00346275, digital image: JPS, P], S [S-G-1519, digital image: JPS], Z! [Z-000004448, digital image: Z], photo of isotype from BR! [EA, K]).

Description. Perennial climber. Stems up to 20 m, when young sometimes villose with whitish, articulate trichomes, later often subglabrous to glabrous. Petioles 4–8 cm long, glabrous or with pale, articulate trichomes. Leaves 9.5–16.5 × 10–16.5 cm, shallowly to profoundly 3- or 5-lobate. Lobes triangulate, ovate to elliptical. Leaf margin entire and denticulate to serrate. Upper leaf surface glabrous or with hyaline to white pustules. Lower leaf surface glabrous or sometimes villose with whitish, articulate trichomes, sometimes with white pustules on the main veins. Probracts up to 3.5 mm. Tendrils simple or bifid. Male flowers in racemes, rarely accompanied by one solitary flower, or one single flower only. Common peduncle 3–4.5 cm, pedicels up to 7 mm, bracts up to 1 mm, caducous. Pedicels of solitary flowers up to 2.5 cm, each glabrous. Perianth tube glabrous, calyx lobes up to 2.5 mm, triangulate to lineal, in buds adpressed to corolla, later spreading. Corolla 1.2–2.9 cm long, orange buff, lobes 0.3–1 cm. Filament column and anther head not seen. Pollen sacs cream yellow. Female flowers solitary, pedicel 3–8 cm, glabrous. Hypanthium glabrous, calyx lobes and corolla like in male flowers. Ovary glabrous, ribbed. Style not seen, stigma bulging, yellow. Fruit up to 20 cm long and 5 cm in diameter, unripe green with white mottling and longitudinal green lines, ripening via yellow, orange into deep red. Seeds 6–7 × 5 × 1.5 mm (L/W/H), symmetrically obovate, face flatly lenticular.

Phenology. Flowering time: January–April, June, August, September, November, December, likely throughout the year.

Distribution. Fig. 34. NW Burundi, D. R. Congo (North Kivu, South Kivu), Rwanda (Western Province), Tanzania (Iringa: Kipengere Range, Uzungwa Mts; Kigoma: Mahali Mts; Mbeya: Kipengere Range; Morogoro: Uluguru Mts; Ukaguru Mts, Uzungwa Mts), Uganda (Western Region). Elevation 1200–2600 m. Afromontane cloud forests and mist forests, upland rainforests, bamboo forest, in *Macaranga kili-mandsharica* shrubs, rarely in *Pennisetum* savannas. On basalt and graphitic schist soils, lateritic clay. Introduced in Kenya.

Vernacular names. Kihunde: mutanga (*Deru* 485), Kindanda: mwore (*Deru* 485), Kinande: mombowa (*P. Gille* 218), Kinyarwanda: umuvunguvungu (*G. Bouxin* 820), umufungofungo (*G. Troupin* 11163), umwonkalere (*Deru* 485), Kisafwa: itangalulu (*C.J. Kayombo* 1003).

Remarks. *Coccinia ulugurensis* cannot be definitely distinguished from *C. mildbraedii*. The leaves are 3-lobate with rather triangulate lobes towards central Tanzania (*C. ulugurensis*), whereas in the western areas the leaves may be deeper lobate with lanceolate lobes (*C. mildbraedii*). Collections of the *C. ulugurensis* form

also occur in the Ukaguru Mts (*M. Thulin & B. Mhoro* 2933) from which a close-by located population has been recollected for sequencing (*N. Holstein et al.* 76) because collections from the Uluguru Mts were not available. However, forms similar to *C. ulugurensis* also occur in the Western Rift area. Vice-versa, 5-lobate leaves also occur in central Tanzania. Jeffrey (1967) used to distinguish the species also by the occurrence of “crisped hairs”, but these do not seem to be specific. Such trichomes also occur in other species, such as *C. adoensis*, *C. grandiflora* and *C. mackenii*, and the fine pubescence as described for *C. ulugurensis* regularly occurs in young shoots and often disappears later on. Collections that represent the two species cluster together (Fig. 17) and either both species share major parts of their distribution ranges and only differ morphologically in nuances of lobation depth, or they belong to a single species, of which here the latter case is assumed here.

Specimens examined. (Selection, in total: 76) D.R. Congo. North Kivu: Lubero territory, Bingi, *A. Léonard* 5415 (BR, EA, WAG [WAG0225422], WAG [WAG0225423], WAG [WAG0225424]). South Kivu: Kabare territory, Marais Musisi, 28°42'E, 2°16'S, *P. Bamps* 2844 (BR, EA, WAG [WAG0225420]). Kenya. Central Province: Limuru, tea estate, introduced from Tanzania, *J.B. Gillett* 20185 (EA, MO). Rwanda. Western Province: Shangugu territoire, Mont Bigugu, *A.R. Christiaensen* 1616 (EA, WAG [WAG0225419]). Southern Province: Rutovu, km 64 on Astrida [Butare]–Shangugu [Cyangugu] route, *M. Reynders* 394 (BR). Tanzania. Iringa: Dabaga Highlands, Ihangana Forest Reserve, near Kibangu, 18 mls [29 km] S of Dabaga, *R. Polhill & S. Paulo* 1476 (B, BR, EA, K, P [P05620648], PRE). Kigoma: Mpanda district, Mahali Mts, Sisaga, c. 6°S 30°E, *T.G. Jefford & J.G.B. Newbould* 1924 (COI, EA). Mbeya: Mbeya rural district, Umalila Forest Reserve, c. 7 km W of Ruanda II on road to Izumbwe (2 km SSE of main peak of Mbogo Mt.), 9°11'S, 33°18'E, *R.E. Gereau et al.* 5060 (K, MO). Morogoro: Kilosa district, Ukaguru Mts, between Mandege and Masenge, 6°22'S, 36°58'E, *M. Thulin & B.E. Mhoro* 2792 (DSM, EA, K, MO). Uganda. Western Region: Kigezi district, Virunga chain, northern foot of Mzhavura Mt., Nkanda, *H.U. Stauffer* 931 (BR, M, Z).

15. *Coccinia ogadensis* Thulin, *Kew Bull.* 64: 485. 2009.

Type: Ethiopia. Somali Region: Harerge, 5 km S of Qarsonney, female, fr, 15 May 2006, *M. Thulin et al.* 11183 (Holotype: ETH; isotypes: K! [K000543219, digital image: K], UPS!).

Type: [Ethiopia]. [Somali Region]: Somaliland, Harradigi [Harradigit], Mar 1885, *F.L. James & J.G. Thrupp s.n.* (Paratype: K!).

Type: [Ethiopia]. [Somali Region]: Somaliland, Harradiqi [Harradigit] or Boobi, Mar or Apr 1885, *F.L. James & J.G. Thrupp s.n.* (Paratype: K!).

Type: [Ethiopia]. [Somali Region]: Agar Ven [Agar Uen], 6°30'N, 45°20'E, 2500 ft [c. 760 m], red sandy soil, bushland, 25 Oct 1953, *P. Ellis* 163 (Paratypes: FT!, K (2)!).




Figure 35. Distribution map of *C. ogadensis* (blue dots; based on 8 collections) and *C. samburuensis* (pale yellow dots; based on 4 collections). For Ethiopia the borders of the regions and for Kenya the borders of the provinces are displayed.

Type: [Ethiopia]. [Somali Region]: W of Shillavo (Scillave) [Shilabo], 6°25'N, 44°42'E, 1300 ft [c. 400 m], sandy soil, bushland, Nov 1955, *P. Ellis* 383 (Paratype: K!).

Type: [Ethiopia]. [Somali Region]: E of Gorrahei [Korahe], 700 m, 1 Nov 1967, *P.R.O. Bally* 12989 (Paratypes: G!, K!).

Type: [Ethiopia]. [Somali Region]: Scillave [Shilabo]–Wardere road, 6°13'N, 44°45'E, 1130 ft [c. 344 m], red sandy soil, open bushland, male, fl, 2 Apr 1956, *J. Simmons* S63 (Paratypes: EA!, K!).

Type: [Ethiopia]. [Somali Region]: 11 km NE Scillave [Shilabo], 6°10'N, 44°52'E, 1300 ft [c. 400 m], red sandy soil, open bushland, 13 Apr 1956, *J. Simmons* S179 (Paratypes: EA!, K!).

Type: Somalia. [Mudug]: 47 miles [75 km] from Galkayo [Gaalkacyo] on Garoe [Garroowe] road, c. 1000 ft [c. 300 m], red sandy loam and limestone ridges, 15 Oct 1959, *C.F. Hemming* 1713 (Paratypes EA!, K!).

Description. Perennial? climber or trailer. Stems up to 2 m or longer. Stems glabrous, except for nodes with short trichomes, sometimes white pustulate. Petioles 4–15 mm long, glabrous or nearly so. Leaves deeply (3- or) 5-pedately lobate. Central lobe 2–8.5 cm long, 1–8 mm wide, lateral lobes shorter. Lobes entire or dentate to lobulate, linear to oblanceolate. Leaf margins often revolute, apex obtuse with

final (brownish? colored) tooth to acute. Upper leaf surface glabrous, pale to white pustulate; pustules up to 5 mm in diam. Lower leaf surface glabrous, at base with pale aureolate glands between nerves. Probracts < 1 mm with short, whitish trichomes. Tendrils simple. Male flowers solitary, clustered, or in few-flowered racemes. Common peduncle up to 2 cm, glabrous. Pedicels up to 4–20 mm long, subglabrous to glabrous. Perianth tube glabrous, calyx lobes 1–6 mm long, in buds erect, later reflexed, glabrous or nearly so, linear to narrowly triangulate. Corolla 1.7–2.5 cm, white with green veins or yellow, lobes 0.7–1.3 cm. Petals inside with multicellular trichomes, outside with short, oligocellular trichomes. Color of filament column, anther head, and pollen sacs not seen. Female flowers not seen, but very likely solitary, pedicels, hypanthium/perianth tube, calyx lobes and petals not largely differing from male flowers. Fruits spindle-shaped to shortly cylindrical, 4.5–5.5 cm long, c. 1.5 cm in diameter, sometimes with short apical tip (“beaked”). Unripe green with elongate with spots, turning red with whitish, elongate spots. Seeds $4\text{--}5.5 \times 2\text{--}2.5 \times 1\text{--}1.5$ mm (L/W/H), slightly asymmetrically obovate, face flatly lenticular.

Phenology. Flowering time: Imperfectly known, flowering in April and in October and November during rainy seasons.

Distribution. Fig. 35. Eastern Ethiopia (C and E Somali Region), Central Somalia (Mugud). Elevation 300 to 800 m. Red sand and sandy loam, limestone soils. Open *Acacia-Commiphora* bushland, *Cordeauxia edulis* bushland, semi-desert.

Use. Fruits are reported to be edible, juicy, and thirst-quenching (*P.R.O. Bally* 12989).

Vernacular names. Somali: dudu (*P.R.O. Bally* 12989), ilgeel (Thulin 2009), Somali?: lehailu (*J. Simmons* S179), Somali?: salo dudub (*J. Simmons* S63).

Remarks. This species is similar to collections of *C. grandis* with deeply lobate leaves (described as *Coccinia palmatisecta*). However, the lobules in *C. grandis* are much more distinct when the lobulation is that deep. Apart from this, fruit and seed shape of *C. ogadensis* resemble that of *C. adoensis*.

Ellis notes on the collections no. 163 and 383 a smell of rotten meat. However, it is unclear, whether this is coming from the flowers or from crushed leaves. Several cucurbit species have a putrid smell when crushed, such as *Kedrostis foetidissima* or *Momordica foetida*, but this has never been reported for a *Coccinia* species.

Specimens examined. (in total: 10). Ethiopia. Somali Region: Ogaden, *J. Simmons* 64 (EA).

16. *Coccinia pwaniensis* Holstein, *Kew Bull.* 65(3): 435. 2010 [publ. 1 Jan 2011].

Type: Kenya. [Coast Province]: Kwale District, Buda Mafisini forest, 8 miles [12.9 km] WSW of Gazi, 80 m, male, fl, 22 Aug 1953, *R.B. Drummond & J.H. Hemsley* 3953 (Holotype: K! [3 sheets, K000309479, the other two sheets without barcode], isotype: EA!).

Type: Kenya. Coast Province: Kilifi District, Mangea Hill, 39°42'E, 03°16'S, 450 m, dry bushland with *Cynometra* sp., *Brachylaena* sp., *Manilkara* sp., *Brachystegia*

sp., *Julbernardia* sp., *Diospyros* sp., *Xylopia* sp., *Inhambanella* sp., 28 Dec 1988, W.R.Q. Luke 1601 (Paratype: EA!).

Type: Kenya. [Coast Province]: Kwale District, Cha Simba forest, 300 m, fr, 1 Feb 1953, R.B. Drummond & J.H. Hemsley 1078 (Paratype: K!).

Type: Kenya. [Coast Province]: Kwale District, Shimba Hills, Giriama Point area, 1250 ft [c. 381 m], forest edge, 17 Mar 1968, F.C. Magogo & P. Glover 315 (Paratypes: EA!, K!).

Type: Kenya. [Coast Province]: Kwale District, Shimba Hills, Pengo Hill area, 1500 ft [c. 457 m], forest, 27 Mar 1968, F.C. Magogo & P. Glover 493 (Paratypes: EA!, K!).

Type: Kenya. Shimba Hills, SE-part of Longomagandi Forest, 350 m, lowland rainforest, 13 Nov 1988, R. Schmidt 1203 (Paratype: EA!).

Type: Kenya. Kwale District, no detailed location given, 15 Jun 1957, Saunders 11241 (Paratype: EA!).

Type: Tanzania. Pwani: Bagamoyo District, Zaraninge Forest in Kiono Plateau, 38°36'E, 6°09'S, 1000 ft [c. 305 m], dry evergreen coastal forest, on sand, 14 Mar 1990, *Frontier-Tanzania Coastal Forest Research Programme* 1041 (Paratype: K!).

Type: Tanzania. [Pwani]: Kirasawe District: Pugu Hills Forest Reserve on Dar es Salaam–Kisarawe road. Roadside in forest, 100–270 m, 12 May 1970, K.H. Macaulley CVL 102 (Paratypes: DSM!, EA!).

Type: Tanzania. [Pwani]: Pugu Hills, 19 Mar 1939, J.H. Vaughan 2774 (Paratype: EA!).

Type: Tanzania. [Pwani]: Pugu Hills Forest Reserve, road W from road-tunnel, 100 m, in bushes by car-track through forest, 23 Jul 1972, R.C. Wingfield 2056 (Paratypes: DSM!, EA!).

Coccinia sp. B in C. Jeffrey, F. T. E. A.: 69. 1967. R.B. Drummond & J.H. Hemsley 1078 (K!); R.B. Drummond & J.H. Hemsley 3953 (K!, EA!); Saunders 11241 (EA!); J.H. Vaughan 2774 (EA!).

Description. Perennial climber or creeper. Stems up to 3 m long, glabrous. Petiole 0.6–4.1 cm, adaxial side glabrous or with short, stiff trichomes, abaxial side with stiff, patent trichomes that can be quite reduced, then appearing wart-like or subglabrous. Leaves 2–10.4 × 2.7–11.4 cm, shallowly to profoundly 3-(or 5-)lobate, lobes broadly triangulate to elliptic, margin minutely dentate, tips acute. Upper leaf surface minutely hyaline pustulate, nerves sometimes with tiny trichomes, lower leaf surface glabrous, rarely with blackish glands at base, nerves towards the base with stiff, patent trichomes that can be quite reduced, then appearing wart-like or subglabrous. Probracts 2–3 mm long. Tendrils simple. Male flowers in racemes, sometimes accompanied by 1–2 solitary flowers. Peduncle 3.2–7.7 cm, glabrous, pedicels of flowers in racemes 0.2–1 cm, bracts 1–1.5 mm, pedicels of solitary flowers up to 3.8 cm. Perianth tube glabrous, calyx lobes 2.5–3.5 mm long, subulate and spreading, corolla pale yellow to pale orange-yellow, 1.7–2.6 cm, lobes 1–2 cm. Color of filament stalk, anther head, and pollen sacs not seen. Female flower not seen, perianth likely like in male flowers. Style and stigmas not seen. Fruit solitary, petioles at maturity 20–33 mm long, fruit shape oblong-fusiform, 6.2–8 × 1.8–2.3 cm, rarely (?) with an up to 5.5 cm long sterile apical

tip, immature green with pale longitudinal mottling, at maturity becoming orange-red to scarlet-red with pale mottling. Seeds $6.5\text{--}7 \times 4\text{--}4.5 \times c. 1.5$ mm (L/W/H), more or less symmetrically obovate, face lenticular.

Phenology. Flowering time: January–March, June–August.

Distribution. Fig. 23. Kenya (Coast Province), Tanzania (Pwani, but likely also in Dar es Salaam region and Tanga). Elevation 80–460 m. On sandy soil. Open and disturbed places of East African coastal forests and woodlands (*Brachystegia* sp., *Julbernardia* sp., *Diospyros* sp.).

Vernacular names. Kidigo: mnokonyoka (*F.C. Magogo & P. Glover* 493), mtambaa (*F.C. Magogo & P. Glover* 315), Kijibana: muri ya nyoka (*L.J. Lap* 258).

Remarks. Morphologically this species (the only one missing DNA sequences) is close to *C. senensis*. The indumentum is reduced in prominence and in extent to the petiole and leaves in *C. pwaniensis*, and the leaves are rather 3-lobate and long petiolate, in contrast to often 5-lobate and shortly petiolate leaves in *C. senensis*. The racemes in *C. pwaniensis* have considerably more flowers than in *C. senensis*. However, both species share the subulate calyx lobes, and fruit and seed shape suggest that both species are closely related with *C. adoensis*. As *C. pwaniensis* and *C. senensis* do not co-occur, they might be sister species from allopatric speciation, with *C. pwaniensis* occurring in a refugial distribution in the northern coastal forests of East Africa.

Specimens examined. (in total: 13) Kenya. Coast Province: Kilifi district, Kaya Jibana, entering southern forest patch of Kaya Jibana following the path from shop/hoteli at Mwarakaya–Ribe road, $3^{\circ}50'0''\text{S}$, $39^{\circ}40'30''\text{E}$, *L.J. Lap* 258 (WAG [WAG0195516], WAG [WAG0195517]); Kwale district, Shimba Hills, Longomagandi Forest, *R. Schmidt* 527 (EA, UBT).

17. *Coccinia quinqueloba* (Thunb.) Cogn. in A.DC. & C.DC., *Monogr. Phan.* 3: 533. 1881.

Bryonia quinqueloba Thunb., Prodr. Pl. Cap. 1: 13. 1794. *Cephalandra quinqueloba* (Thunb.) Schrad. ex Eckl. & Zeyh., Enum. pl. afric. austral. 2: 280. 1836. *Momordica quinqueloba* (Thunb.) E.Mey. ex Drège, Zwei pflanzengeogr. Dokum.: 126, 132, 133, 137, 202. 1843.

Type: [South Africa]. [Eastern Cape]: sylva Krakakamma, male, fl, Dec, *C.P. Thunberg* 22836 (Lectotype, designated by Meeuse (1962: 99) and here: UPS-THUNB! [K neg. 2978]).

Type: ibid., male, fl, Dec, *C.P. Thunberg* 22837 (Syntype: UPS-THUNB! [K neg. 2977]).

Type: Cap. b. spei [Cape of Good Hope colony], *C.P. Thunberg s.n.* (Syntype: S! [S08-12379]).

Description. Perennial creeper or climber. Stems up to 9 m, glabrous (rarely with remote trichomes). Leaves usually sessile, petiole 3–8(–17) mm, glabrous (rarely with remote trichomes). Leaves, $3\text{--}9.5 \times 4\text{--}10$ cm, 3- or 5-lobate, auriculate. Lobes oblong,

elliptical to obovate. Leaf margin remotely dentate, apices towards lobe often serrate. Lobe apices obtuse with a point. Upper leaf surface pale pustulate, lower leaf surface glabrous, rarely a few blackish glands near base, nerves rarely white-speckled. Probracts < 1 mm or missing. Tendrils simple. Male flowers solitary or in racemes. Common peduncle 0.5–2 cm, petiole in racemous flowers up to 1.8 cm, bracts > 1 mm or missing, solitary flowers with petiole 1.8–4 cm, all glabrous. Perianth tube glabrous, calyx lobes 1.5–3 mm, narrow triangulate, erect to spreading. Corolla 1.2–2.2 cm long, pale yellow, corolla lobes 0.8–1.2 cm. Color of filament stalk, anther head, and pollen sacs not seen. Female flowers one solitary. Petiole 1–2.5 cm, glabrous. Hypanthium glabrous, calyx lobes and corolla like in male flowers. Style and stigmas not seen. Fruits 3.5–9 × 3–4 cm, elliptical to oblong, sometimes short elongated tip, unripe green with longitudinal, white mottling, ripe (orange-)red. Seeds 6–7.5 × 3–3.5 × 1–1.2 mm (L/W/H), slightly asymmetrically obovate, face (flatly) lenticular.

Phenology. Flowering time: January, February, April, July, September, November, December.

Distribution. Fig. 30. Southern and western Eastern Cape, South Africa. Elevation sea level to 1000 m. Sandy soils, also on dolomite soil. Coastal bushland, forest, dry bush, on bushes along rivers, along roadsides.

Remarks. See also under *C. mackenii*.

Taxonomic remarks. *Cephalandra quinqueloba* is the type species of the genus *Cephalandra*. Meeuse (1962) designated the lectotype of *Bryonia quinqueloba* to UPS but did not choose a specimen, which is done here.

Specimens examined. (Selection, in total: 77) South Africa. Eastern Cape: East London, Dec 1916, *H.G. Breyer s.n.* TRV23225 (PRE); Amatle Mts, Hogsback Pass, 32°36'50"S, 26°55'25"E, *P.B. Phillipson 1079* (MO, PRE); Glen Avon, Feb 1923, *Mrs. J.E. Brown s.n.* (PRE ["PRE43005"], Z); Grahamstown, Old Quarry, *R.D.A. Bayliss 8470* (G (2), M, MO, Z); 28 mls [45 km] from Grahamstown on Port Elizabeth road, *R. Story 2346* (B [B 10 0019800], L, M (2), MO, PRE, S [S08-12378]); near Port Alfred, *J.L. Sidey 1095* (PRE, S [S08-12464]).

18. *Coccinia racemiflora* Keraudren, *Adansonia*, ser. 2, 8(1): 41. 1968.

Type: Gabon. [Moyen-Ogooué]: Abanga, C. E. F. A. [Compagnie d'Exploitations Forestière Africaine] lot, male, fl, Jun 1963, *N. Hallé 2425* (Holotype: P! [P00346267, digital image: JPS], P! [P00748828, digital image: JPS, P]).

Type: Gabon. Ibid., female, fl, *N. Hallé 2305* (Paratype: P! [P00748829, digital image: JPS, P], P! [P00748830, digital image: JPS]).

Description. Perennial climber or prostate creeper. Stems up to 5 m, glabrous. Petioles 0.5–2.5 cm, on adaxial side often with line of thin smutty-beige trichomes or glabrous, abaxial side glabrous. Leaves 6.5–11 × 5–9.5 cm, hastate to 3-lobate with central lobe dominating, auriculate (auricles may reach the stem). Lobes triangulate. Leaf margin




Figure 36. Distribution map of *C. racemiflora* (blue dots; based on 8 collections, including 1 supposed hybrid).

entire to somewhat angulate, remotely dentate. Teeth darkening when dried. Upper leaf surface with waxy cover glabrous with few-celled clear pustules. Lower leaf surface glabrous with dispersed blackish glands. Probracts up to 2 mm. Tendrils bifid. Male flowers ebracteate in lax, glabrous racemes. Common peduncle up to 2.5 cm, pedicels 0.3–1 cm. Perianth tube glabrous, calyx lobes 0.5 mm, shortly lineal, spreading. Corolla c. 1.2 cm, yellowish to orange, lobes 2–4 mm. Color of filament column, anther head, and pollen sacs not seen. Female flowers ebracteate in lax, glabrous racemes, like in males. Hypanthium glabrous, calyx lobes, and corolla like in males. Ovary glabrous. Style and stigmas not seen. Unripe fruits glabrous, glaucous, globose.

Ripe fruits unknown, size c. 1.5 cm in diam.? Seeds $5 \times 3 \times 1.5$ mm (L/W/H), rather symmetrically obovate, flatly lenticular.

Phenology. Flowering time: January–March, imperfectly known.

Distribution. Fig. 36. Gabon, S Cameroon. Tropical lowland rainforest.

Taxonomic remarks. One *Hallé* 2425 specimen bears a type label but it is not clear whether it was attached by Kéraudren herself. Since both specimens appear to be part of the same individual, both are best to be treated to be holotypes mounted on two sheets.

Remarks. A collection from the Gamba area in S Gabon (*M.A. van Bergen* 490 (WAG [WAG0151338])) is morphologically close to *C. racemiflora* but shares the calyx lobes with *C. barteri* and thus may represent a hybrid. Using plastid markers, this collection (*C. barteri* 6) clusters within *C. barteri*, while in the nuclear *LEAFY*-like tree, it clusters with one representative of *C. racemiflora*, but not with both (Figs 17, 18; Holstein and Renner 2011b). Another collection (*F.J. Breteler et al.* 8835 (MO, WAG) from Saint Germain area, C Gabon, has calyx lobes typical for *C. racemiflora*, but a more condensed raceme as in the *M.A. van Bergen* 490 collection. The hypanthia of the female flowers are urceolate, whereas in male flowers they are cup-shaped. Urceolate hypanthia are also found in some *C. barteri* forms, and a phylogenetic significance, e.g., by introgressive hybridization, cannot be ruled out.

Specimens examined. (Selection; in total: 10) Cameroon. South Region: 3 km N of km 20 Kribi-Lolodorf, high forest exploitation, 3°01'N, 10°03'E, *J.J. Bos* 6590 (WAG [WAG0225514], WAG [WAG0225515]). Gabon. Estuaire: 12 km SW of Kinguéle Falls, *N. Hallé & J.F. Villiers* 5357 (K, P [P05620813]). Ngounié: 35 km on road from Lebamba to Yéno, 1°58'S, 11°25'E, *J.J.F.E. de Wilde & M. Sosef* 10456 (WAG [WAG0044628]). Ogooué-Maritime: Rabi North, 1°51.6'S, 9°51'E, *I. van Nek* 536 (WAG [WAG0044627]).

19. *Coccinia rehmannii* Cogn. in Schinz, Bull. Herb. Boiss. 3: 418. 1895.

Type: South Africa. Transvaal: boshveld ad Klippan [according to Meuse (1962) in Limpopo: Greater Sekhukhune District Municipality, Doornpoort; 24°37'S, 29°26'E], 1875–1880. *A. Rehmann* 5156 (Syntype: Z! [Z-000004445, digital image: Z]; syntype: BR! [BR0000005111602, digital image: BR, JPS]).

Type: South Africa. Ibid., *A. Rehmann* 5157 (Syntype: Z! [Z-000060807, digital image: Z]).

Type: South Africa. Ibid., *A. Rehmann* 5168 p.p. (Lectotype, designated by Meeuse (1962: 102): Z! [Z-000060808, digital image: Z]; isolectotype: BR! [BR0000005111930, digital image: BR, JPS], K! [K000313196, digital image: JPS, K]).

Type: South Africa. At Eland river, *A. Rehmann* 4944 [*sic*, must be *A. Rehmann* 4954, see Taxonomic remarks] (Syntype: Z! [Z-000060806, digital image: Z]).

Coccinia rehmannii Cogn. var. *littoralis* A.Meeuse, Bothalia 8: 104. 1962. pro parte ex *R. de Carvalho s.n.* (Paratypes: COI (2)!).

- Type: South Africa. [Eastern Cape]: [Amatole District Municipality], Komgha, Kei Mouth, *H.G. Flanagan 457* (Holotype: PRE [PRE0190559-0, digital image: JPS], isotypes: BOL?, NBG?).
- Type: South Africa. [Eastern Cape]: Cape Morgan, *H.G. Flanagan 457* (Paratype: GRA [GRA0002852-0, digital image: JPS], BOL?, NBG?).
- Type: South Africa. [Eastern Cape]: East London, Nahoon, *M.W. Nanni 151* (Paratype: PRE!).
- Type: South Africa. [Eastern Cape]: Coffee Bay, *W. Tyson 24* (Paratypes: B!, COI!, GRA, MO!, NY!, PRE!, S! [S08-12380]).
- Type: South Africa. [KwaZulu-Natal]: 10 mls NW of Mtubatuba, *L.E.W. Codd 9620* (Paratypes: COI!, M! [M-0198513], PRE).
- Type: South Africa. [Kwa-Zulu Natal]: Umhlanga Rocks, *A. Dohse & B. de Winter 223* (Paratypes: NH, PRE!).
- Type: South Africa. [Kwa-Zulu Natal]: Manaba Store, *J. Gerstner 3407* (Paratype: NH).
- Type: South Africa. [Kwa-Zulu Natal]: Dhlebe, *J. Gerstner 4261* (Paratypes: NH, PRE).
- Type: South Africa. [Kwa-Zulu Natal]: near Durban, *T.J. Jenkins TRV7092* (Paratype: PRE).
- Type: South Africa. [Kwa-Zulu Natal]: Mtunzini, *S.M. Johnson 612* (Paratype: NBG).
- Type: South Africa. [Kwa-Zulu Natal]: Stella Bush, *W.E. Marriott herb. no. 24341* (Paratype: ?).
- Type: South Africa. [Kwa-Zulu Natal]: *ibid.*, *W.E. Marriott herb. no. 27143* (Paratype: NH).
- Type: South Africa. [Kwa-Zulu Natal]: Shelly Beach, *A.O.D. Mogg 11941* (Paratype: ?).
- Type: South Africa. [Kwa-Zulu Natal]: *ibid.*, *A.O.D. Mogg 12070* (Paratypes: M! [M-0198511], M! [M-0198512], PRE!).
- Type: South Africa. [Kwa-Zulu Natal]: without detailed location, *T.B. Oatley C 15* (Paratype: PRE).
- Type: South Africa. [Kwa-Zulu Natal]: Berea, *Small herb. no. 34714* (Paratype: NH).
- Type: South Africa. [Kwa-Zulu Natal]: Ubombo coastal veld, *P.A. Tosh 28* (Paratype: NU).
- Type: South Africa. [Kwa-Zulu Natal]: Ndumu Game Reserve, *Ward 3169* (Paratype: ?).
- Type: South Africa. [Kwa-Zulu Natal]: *ibid.*, *Ward 3170* (Paratype: ?).
- Type: South Africa. [Kwa-Zulu Natal]: Umvoti, Thorns near Greytown, *J.M. Wood 5318* (Paratype: NH).
- Type: South Africa. [Kwa-Zulu Natal]: Durban, Berea, *J.M. Wood 6350* (Paratypes: BOL, L!, NBG, NH, PRE).
- Type: South Africa. [Kwa-Zulu Natal]: Doonside, *J. Wylie herb. no. 23299* (Paratype: NH).
- Type: Mozambique. Maputo: Lourenço Marques [Maputo], *J.M. Borle 253* (Paratypes: M! [M-0198510], MO!, P! [P05620807, digital image: P]).
- Type: Mozambique. Maputo: *ibid.*, *J.M. Borle 427* (Paratype: ?).
- Type: Mozambique. Maputo: *ibid.*, *J.M. Borle 442* (Paratype: PRE!).
- Type: Mozambique. Maputo: Inhaca Island, *H.G. Breyer TRV20506* (Paratype: PRE).
- Type: Mozambique. Maputo: Inhachingo, *A.W. Exell et al. 630* (Paratype: SRGH).
- Type: Mozambique. Maputo: Massinga, *A.W. Exell et al. 645* (Paratype: SRGH).

Type: Mozambique. Maputo: Lourenço Marques [Maputo], *A.J.W. Hornby* 4599 (Paratype: PRE!).

Type: Mozambique. Maputo: Delagoa Bay [Maputo Bay], *H.A. Junod* 20 (Paratypes: BR!, G (2)!, Z! [Z-000073406, digital image: Z]).

Type: Mozambique. Maputo: Inhaca Island, 6 Jul 1958, *A.O.D. Mogg* s.n. (Paratype: PRE!).

Type: Mozambique. Maputo: ibid., 14 Dec 1955, *A.R.A. Noel* s.n. (Paratype: PRE!).

Type: Mozambique. Maputo: Lourenço Marques [Maputo], *R. Schlechter* 11555 (Paratypes: BOL, COI!, G (3)!, GRA, HBG! [HBG518897], PR! [PR 801378], WAG! [WAG0234182], Z! [Z-000073405, digital image: Z]).

Type: Mozambique. Maputo: Katembe [Catembe], *R. Schlechter* 11614 (Paratypes: G (2)!, GRA, PRE!, Z! [Z-000073407, digital image: Z]).

Coccinia ovifera Dinter & Gilg in Dinter, Veg. Veldkost Südw.-Afrik.: 16. 1912.

Type: [Namibia]. Karas: Sandverhaar, *M.K. Dinter* 1214 (Syntype: ?).

Type: [Namibia]. Otjozondjupa: Otjiwarongo, female, fl, fr, Jan 1912, *M.K. Dinter* s.n. (Syntype: SAM [SAM0072115-0, digital image: JPS]).

Type: [Namibia]. Waldau, female, fr, 3 Feb 1917, *M.K. Dinter* 432 (Lectotype, designated here: SAM [SAM0066515-0, digital image: JPS]).

Further possible syntypes (cited in Dinter 1919/20) if collected before end of 1912: Oshikoto: Gaub, *M.K. Dinter* 2412 (?); Tsumeb, *M.K. Dinter* s.n. (?). Unknown: Hereroland, Palmenwald, *M.K. Dinter* s.n. (?); Hereroland, Wilhelmsberg, *M.K. Dinter* s.n. (?).

Description. Perennial climber or creeper. Stems up to 4 m, glabrous or with broad-based trichomes, when old often densely white pustulate (esp. in drier areas). Petiole 0.2–4.2 cm, glabrous or with erect, broad-based or often up to 1.5 mm long, articulate trichomes or only wart-like, when old sometimes dense white pustulate (esp. in drier areas). Leaves 0.9–9.7 × 1.4–16.6 cm, shallowly to deeply 3- or 5-lobate, auriculate, sometimes lobulate, rarely cordate. Lobes and lobules usually extending, rarely pointing towards tip, narrowly to broadly triangulate to lanceolate. Leaf margin rather remotely denticulate. Apex acute to obtuse, apiculate. Upper leaf surface more or less densely white pustulate, pustules sometimes with a thick, small trichome, on nerves often with thick, small trichomes. Lower leaf surface glabrous, sometimes with small, blackish glands between nerves, nerves usually with erect trichomes, sometimes wart-like. Probract usually absent, if present then up to 3.5 mm. Tendrils simple. Male flowers 1–3 solitary, if fasciculate or in few-flowered racemes then accompanied by 1–2 flowers. Common peduncle 0.7–4.5(–8.5) cm, glabrous or with long, articulate trichomes. Pedicel of flowers in inflorescences 0.6–2.8 cm, bracts up to 2.5 mm or missing. Pedicel of solitary flowers (0.2–)0.5–5(–9) cm, glabrous or especially at apex with long, articulate trichomes. Perianth tube usually with long (> 0.5 mm) trichomes, rarely almost glabrous. Calyx lobes 0.2–7 mm, narrowly lanceolate or lineal, when young erect, later also spreading to reflexed. Corolla 0.8–2.5 cm long, buff to more or less pale yellow, sometimes with green venation. Lobes 0.3–1.1 cm. Filament column pale buff, anthers buff, pollen sacs yellow (Fig. 10a). Female flower solitary. Pedicel 0.4–1.5 cm, glabrous or with long, articulate

trichomes. Hypanthium usually with long (> 0.5 mm) trichomes, rarely almost glabrous, calyx lobes and corolla like in male flowers. Ovary rarely glabrous, often more or less densely covered with articulate trichomes. Style columnar, green. Stigmas 2-lobed, yellow (Fig. 11b). Immature fruit, rarely also at maturity, with whitish, longitudinal mottling that develops a dark green corona during ripening. Ripe fruit globose to elliptical 1–6.2 × 1–2.8 cm, glabrous or with few articulate trichomes, bright orange to red. Seeds 4.5–7 × 2–3.5 × 1–1.2 mm (L/W/H), asymmetrically obovate to somewhat falcate, face flat.

Phenology. Flowering time: January–April, June, October–December.

Distribution. Fig. 37. South Africa (except Western Cape and SW Eastern Cape), Namibia (except hyperarid regions), Swaziland, southern Mozambique (Gaza, Inhambane, Maputo), Botswana, Zimbabwe (Manicaland, Masvingo, Matabeleland South), southern Angola (Namibe, Huila, Cunene, Cuandocubango). Elevation sea level to 1850 m. Limestone, dolomitic, quartzitic, granitic, and ultrabasic soils. Possibly some tolerance to Ni and Cu. On loam, clay, white and red sand, sandstone, and gravel, but prefers sandy (well drained) soils (Meeuse 1962). Full sun to shade. Coastal dunes, riverbanks, *Acacia sclerocarya*–*A. caffra* woodland, *Combretum apiculatum* bushland, *Grewia flava* bushland, mopane, Kalahari thornveld, grassland, semi-desert, dunes. Grazing is tolerated. Light frost seems to be tolerated (Meeuse 1962).

Use. Tuber edible after baking (Dinter 1912, *M.E. Keith* 50, *B. de Winter & O.A. Leistner* 5598). Leaves used as spinach by Shangane people in Gaza province of Mozambique (*K.L. Tinley* 3000). Fruits edible (*R.H.W. Seydel* 938, *R. Story* 5029, *B. de Winter & O.A. Leistner* 5598).

Vernacular names. Muchope [Xichope?]: fuculumué (*L.A. Grandvaux Barbosa & F. de Lemos* 8502), Otjiherero: otjimaga (*M.K. Dinter s.n. Jan* 1912), Ronga dialect [of Xitsonga language]: inyamgwazi (*A.O.D. Mogg* 31308), shiracarana (*L. Macuácuá* 73 and 75), Tsonga [Xitsonga]: inyamwazi (*A.O.D. Mogg* 31538), nyampape (*C. Liengme* 491), Zulu [isiZulu]: uselwa-iwenyoka (Sewram et al. 2006).

Remarks. The (sub-)glabrous “*littoralis*” form can be easily confused with the polymorphic *C. adoensis* (Hochst. ex A.Rich.) Cogn., which differs in shorter calyx lobes and lenticular seeds, and with *C. senensis*, which also has lenticular seeds and usually long-peduncled male racemes.

Meeuse’s variety *littoralis* is hard to define as the paratypes are variable, and characters for delimitation are unclear. For example, although the variety should lack white speckles on the stems, there are some individuals with white speckles along with long pedicels or conspicuous racemes in KwaZulu-Natal and southern Mozambique as in the variety *littoralis*. The holotype of var. *littoralis* is, in the present author’s opinion, rather intermediate between the holotype of var. *rehmannii* and the subglabrous forms, e.g., from Inhaca Island. However, the tendency that Meeuse describes is apparent. Other characters in the collections of his variety, viz. relatively long calyx lobes and petals, also occur in the high mountains of Namibia but also in the whole periphery of the *C. rehmannii* distribution range. Strangely, Meeuse does not mention the most striking difference between *C. rehmannii* collections from the inland/dry areas and coastal/peripheral collections being the globose fruit in inland/dry area individuals




Figure 37. Distribution map of *C. rehmannii* (blue dots; based on 251 collections). For South Africa the borders of the provinces are given.

or long elliptical fruit in coastal/peripheral individuals (shown and mentioned in this treatment as "*C. rehmannii* aff. var. *littoralis*"; Fig. 10a) respectively. Collections with this fruit morph also occur in Angola and Zimbabwe but do not follow Meeuse's other characters of the var. *littoralis*. On the other hand, collections with long calyx lobes and long petiolate flowers can have globose fruits. Like the area of the southeastern coast of South Africa, areas in the north of southern Africa receive more and longer rainfall per year than the inland, so there is a possible correlation between precipitation and fruit morphology. Elliptical fruits also occur in the closely related *C. trilobata* from mountainous and thus more humid habitats but not in *C. microphylla* from the dry bushlands of NE Africa whose fruit is globose. However, the characterization by Meeuse that *C. rehmannii* is an aggregate species seems to be true. It might be interesting to link morphological characters with haplotypes and to test the fitness of these morphs in the different habitats. In any case, the morphological differentiation seems to be stable in cultivated individuals, and artificial crosses between different forms (inland vs. subglabrous from the Southeast) result in the onset of a normal fruit (resulting seeds were not used for cultivation).

The placement of *C. rehmannii* var. *littoralis* forms with other *C. rehmannii* forms in plastid and nuclear phylogenies (Figs 17, 18) also rejects the hypothesis of a hybrid origin (e.g., with *C. adoensis* or *C. senensis*), but rather suggests parallel evolution due to ecological factors.

Taxonomic remarks. The protologue contains a literal mistake for the syntype from Eland River and must be corrected from 4944 to 4954. On the one hand, *A. Rehmann 4944* (GRA, K) is a Malpighiaceae. On the other hand, there is *A. Rehmann 4954* (a *C. rehmannii*) in Z from Eland river and with a remark by Cogniaux ‘sp. nov.’ Hence, *A. Rehmann 4954* is the syntype of *Coccinia rehmannii*, not *A. Rehmann 4944*.

The GRA specimen of *H.G. Flanagan 457*, which is supposed to be the isotype (cited by Meeuse) of *C. rehmannii* var. *littoralis*, is in fact merely a paratype. Meeuse stated clearly the location as “Komgha: Kei Mouth” and chose the PRE specimen from there as the holotype, but the GRA specimen is from the nearby located Cape Morgan. Apparently, Flanagan used the same collection number for different gatherings. The GRA specimen thus cannot be regarded as a duplicate despite the same number. As the GRA specimen does not have a label by Meeuse, he just cited the specimen without seeing it.

The similarity of *C. rehmannii* var. *littoralis* to *C. senensis* led to a misplaced paratype. One of the two specimens by R. de Carvalho is a syntype of *C. jatrophiiifolia* var. *australis* Cogn. and the two R. de Carvalho specimens from COI are paratypes of *C. subglabra*, which are both synonyms of *C. senensis*.

Coccinia ovifera is a validly published name, although the description is a little cryptic, hence the species name is not a *nomen nudum*. Dinter writes that he has found, viz. collected, the species around Grootfontein, in Hereroland (not in the narrow sense of the 1968 homeland) and in Sandverhaar (Namaland). Therefore, the requirements for validity are met (37.3 Note 2). The latter site is cited by him explicitly in a later publication (Dinter 1919/20). The collections designated as syntypes above match the description as given in the protologue. As the present author did not see the specimens designated as “possible syntypes”, they are tentative and might have been destroyed in Berlin.

Specimens examined. (Selection, in total: 321) Angola. Huíla: Cambos, near Chiange, *A. Menezes 3629* (LISC [LISC 031347], P). Namibe: andados ca. 55 km de Moçamedes [Namibe] para Dois Irmãos [Caraculo], *E.J. Mendes 3969* (COI, LISC). Botswana. Ghanzi: 200 mls [320 km] NW of Molepolole, *R. Story 5029* (COI (2), EA (2), PRE (2), Z). Kgalagadi: c. 50 mls [80 km] NNW of Tsabong, *O.A. Leistner 3120* (LISU, M, PRE). Ngamiland: 80 km W of Tsau on Cae Cae road, *D.G. Long & D.A.H. Rae 416* (E). Mozambique. Gaza: Vila de João Belo [Xai-Xai], entre Chicumbane e a Barra do Limpopo: próximo da povoação commercial da Barra, *F. de Lemos & A. Balsinhas 131* (BM, COI) and *133* (BM, COI (2)). Inhambane: Pomene, in hotel area, *P.C.M. Jansen et al. 7533* (G n.v. [G00305107], MO, WAG [WAG0234198], WAG [WAG0234199]). Maputo: between Costa do sol and Marracuene, Mutanhane, *A. Balsinhas 230* (BM, COI, PRE). Namibia. Erongo: Farm Anschluss, 150 km E of Swakopmund on Khomas road to Windhoek, *B. de Winter & D. Hardy 8001* (M, PRE, WAG [WAG0234173]). Hardap: c. 20 mls [32 km] from Kalkrand on road to Rehoboth, *B. de Winter 3538* (COI (2), L, M, PRE).

Karas: Dassiefontein Farm, 2–3 km E of highway, in foothills of Groot Karasberge, c. 64 km NNE of Grünau, *G. Davidse & A. Loxton* 6240 (M, MO, S). Omaheke: [Farm] Breitenberg, border of Kalahari, *R.H.W. Seydel* 2513 (COI, M, WAG [WAG0234172]). Otjozondjupa: 160 mls [257.5 km] E of Grootfontein, Gautscha Pan, *R. Story* 6219 (PRE); *ibid.*, *R. Story* 6238 (M (2), PRE (2), S [S08-12417]). South Africa. Eastern Cape: Port St. Johns distr., First Beach, *M.J. Wells* 3434 (MO). Gauteng: Pretoria, Brummeria: BRI, *A. Balsinhas* 3474 (MO, PRE, WAG [WAG0234189]). KwaZulu-Natal: Lower Tugela Valley, below Maqumbi, *D. Edwards* 3053 (B, M, HEID, PRE); 5 mls [8 km] on Nkonkoni–Pongola road, *M.J. Wells* 2162 (B, EA, M, Z [Z-000060820]). Limpopo: Immerpan, near post office on roadside, *A.D.J. Meeuse* 9452 (B, COI, L, PRE, Z [Z-000060813]). Mpumalanga: Impala, siding, *E. Retief* 1260 (PRE); *ibid.*, *E. Retief* 1261 (MO, PRE); Steelpoort, Burgersfort, 2 km E of town, 24°40'S, 30°22'E, *H.J. Venter & A. Venter* 10260 (S [S08-12382], WAG [WAG0234178]). Northern Cape: 25 ml. [40.2 km] W of Kimberley, *H.J.E. Schlieben & H.R. Tölken* 11017 (G (2), HEID, M, PRE, S). Northwest: Rooisloot, 6 Apr 1935, *A.O.D. Mogg s.n.* (B, COI, L, PRE); Farm Welgevonden, 6 Apr 1935, *A.O.D. Mogg s.n.* (B, L, PRE [PRE43077], Z [Z-000060812]). Swaziland. Lubombo: Thsaneni [Tjaneni], *I.F. La Croix* 4909 (MO, WAG [WAG0234170]). Zimbabwe. Manicaland: Chipinga district, Sabi Valley Experimental Station, *C. Soane* 162 (COI (3)). Matabeleland South: Beit Bridge [Beitbridge], *A.W. Exell et al.* 425 (LISC); *ibid.*, *L.C. Leach* 10700 (COI, MO).

20. *Coccinia samburuensis* Holstein, Kew Bull. 65(3): 438. 2010 [published on 1 Jan 2011].

Type: Kenya. [Rift Valley Province]: Samburu East District, on Wamba-Isiolo road, 0.7 km S of turnoff to Maralal, c. 1300 m, female, fl, fr, 4 Jul 1974, *R.B. Faden & A.J. Faden* 74/948 (Holotype: MO!, isotype: WAG! [WAG0234153]).

Type: Kenya. Rift Valley Province: Samburu District, Mt Nyiru, southern slopes, near a river, 2°03'N, 36°51'E, 1600 m, 1 Apr 1995, *B. Bytebier et al.* 355 (Paratypes: EA (2)!).

Type: Kenya. Operoi, 1°12'N, 36°49'E, 1350 m, rocky outcrop in *Acacia* woodland, 23 Dec 2004, *W.R.Q. Luke & P.A. Luke* 10787 (Paratypes: EA!, K!).

Type: Kenya. Near Maralal, Lowaweregoi [Lowua Werekoi Mt] 4000 ft [c. 1220 m], rocks in bushland, 5 Dec 1958, *J.G.B. Newbould* 3233 (Paratype: K!).

Description. Perennial climber. Stems up to 5 m, glabrous, except for minute few-cellular trichomes visible under 5–10× magnification. Petioles glabrous, at base white speckles may occur. Leaves 6–14 cm × 10–17 cm wide, (5- or)7-lobate. Leaf lobes elliptical, margin serrate (to lobulate), teeth (lobule tips) with yellowish glands. Lobe apex subacute, apiculate. Upper leaf surface glabrous, more or less clear to white pustulate. Lower leaf surface glabrous, nerves white-speckled. Probracts up to 4 mm. Tendrils simple. Male flowers 1–2 solitary. Pedicel up to 5 cm long, glabrous. Perianth tube glabrous, calyx lobes 6.5

mm long, linear, erect. Corolla 3.7–4 cm long, brownish yellow, lobes 2.2–2.5 cm. Female flowers solitary. Pedicel 4–5 mm, glabrous. Hypanthium tube glabrous, calyx lobes and corolla like in males. Ovary narrowly cylindrical, glabrous. Fruits c. 14×1.5 –2 cm, long cylindrical, unripe green with lighter spots, color of ripe fruit unknown but likely red. Seeds 6.5 – 7×3.5 – $4.5 \times (\geq 1)$ mm (L/W/H), symmetrically obovate, face flatly lenticular.

Phenology. Flowering time: Imperfectly known. Flowering in April, July, and December, but likely to flower as long water is available (rainy seasons).

Distribution. Fig. 35. Only known from Samburu area in Kenya (hence the epithet). Only known from seepage line in rocky (granite) outcrops in *Acacia-Commiphora* deciduous bushland.

21. *Coccinia schliebenii* Harms, Notizbl. Bot. Gart. Berlin-Dahlem 11: 685. 1932.

Type: Tanzania. Morogoro: Ulanga district, Mahenge ward, Mtimaliassi near Mahenge station. 900–1000 m, male, fl, 14 Jan 1932, *H.J.E. Schlieben* 1620 (Holotype: B! [B 10 0154928, digital image: B, JPS], isotypes: BM! [BM001010003], BR! [BR0000008886828, digital image: BR, JPS], BR! [BR0000008887580, digital image: BR, JPS], G! [G00301595], M! [M0105774, digital image: JPS], MA [MA386129, digital image: JPS], P! [P00346274, digital image: JPS, P], S [S-G-1518, digital image: S], Z! [Z-000004447]).

Coccinia calophylla Harms in Mildbraed, Notizbl. Bot. Gart. Berlin-Dahlem 12: 522. 1935.

Type: Tanzania. Lindi: Muera plateau, Bakari, fl, 26 Oct 1934, *H.J.E. Schlieben* 5551 (Holotype: B! [B 10 0154924, digital image: B, JPS], isotypes: BM! [BM001010001], BM! [BM001010002], BR! [BR0000008887153, digital image: BR, JPS], BR! [BR0000008887511, digital image: BR, JPS], G! [G00301767], G! [G00301768], HBG! [HBG506428, digital image: JPS], M! [M0105775, digital image: JPS], MA [MA386127, digital image: JPS], P! [P00346272, digital image: JPS, P], P! [P00346273, digital image: JPS, P], S [S08-11865, digital image: JPS, S], Z! [Z-000073408, digital image: Z], Z! [Z-000073409, digital image: Z]).

Description. Perennial climber. Stems up to 12 m, densely covered with short, stiff, smutty-brownish trichomes. Petioles 1.5–11 cm, indumentum as on stem. Leaves 5–18 \times 4.5–18 cm, slightly to deeply palmately 5-lobate. Lobes broadly triangulate to long elliptical, margin dentate, tips acute or obtuse. Upper leaf surface usually densely covered with short, thin trichomes. Lower leaf surface densely (esp. on nerves) covered with short, stiff, dirty-brownish-beige trichomes. Probracts up to 4.5 mm. Tendrils simple or bifid. Common peduncle 1.1–6.5 cm, with indumentum like on stem to puberulous, pedicels of flowers in racemes with up to 4 mm, indumentum as on peduncle, bracts 3–4 mm. Pedicels of solitary flowers 1.2–5 cm, indumentum as on peduncle. Perianth tube with indumentum like on stem to puberulous. Calyx lobes lineal to narrowly lanceolate 10–15 mm. Corolla 4–6.2 cm long, yellow, apricot, pale orange, sometimes marked with purple, lobes 2–3.2 cm. Filament color not seen, anther head not seen, pollen sacs dark




Figure 38. Distribution map of *C. schliebenii* (pale yellow dots; based on 27 collections). For Ethiopia and Tanzania the borders of the regions are given.

yellow to orange. Female flowers solitary, pedicels 2.5–4.5 cm long, densely covered with short trichomes. Hypanthium with indumentum like on stem to puberulous, calyx lobes, and corolla like in male flowers. Ovary with smutty-brownish trichomes. Style 3–6 mm, color not seen. Stigmas 2-lobed, orange-yellow. Fruit 7–9 × c. 2.5 cm long, oblong to shortly cylindrical, ripening from green with 10 more deeply colored ribs via yellow to red. Seeds 5.5–6 × 2.5–3 × 1 mm (L/W/H), symmetrically obovate, face lenticular.

Phenology. Flowering time: January–March, May–July, December.

Distribution. Fig. 38. Ethiopia (Benishangul-Gomaz?, Gambela, Oromia, SNNPR), Mozambique (Cabo Delgado), South Sudan (Eastern Equatoria), Tanzania

(Iringa, Lindi, Morogoro, Mtwara?, Ruvuma). Elevation 300–1900 m. Black cotton soil, reddish soil, volcanic underground. *Markhamia-Dombeya* woodland, *Chlorophora-Albizia* woodland, *Dalbergia-Pterocarpus-Combretum-Acacia* woodland, woodland with *Acanthus sennii*, *Baphia abyssinica* forests, lower afro-montane forests, termite hills, gallery forests.

Use. Fruits edible (W.J. Kindeketa et al. 2793).

Vernacular names. Didinga: moroich (J.G. Myers 10918), Kipogoro: mdalla (W.J. Kindeketa 2747), Mokonde: ncauedi (M.F. Correia 92).

Specimens examined. (Selection, in total: 31) Ethiopia. Gambela: 20 km E of Punnido, along the new road to Gog, 7°34'N, 34°24'E, I. Friis et al. 7317 (C, K). Oromia: c. 25 km E of Lekemti [Nekemte], W.J.J.O. de Wilde & B.E.E. de Wilde-Duyffes 7184 (K, MO, WAG [WAG0225409], WAG [WAG0225410], WAG [WAG0225411]). SNNPR: Ghibie [Gibe] or upper Omo gorge, 182 km SW of Addis Abeba on road to Jimma, north bank, J.W. Ash 898 (EA (2), K). Mozambique. Cabo Delgado: Maccondes, 2 km from Mueda to Negomano, near Santo António mission, M.F. Correia 92 (LISC). South Sudan. Eastern Equatoria: Didinga Mts, Mt Lotuke, Char, J.G. Myers 10918 (K). Tanzania. Iringa: Mufindi district, Lulanda village, N and NW of Ihili forest patch, 8°35'59"S, 35°37'12"E, M.A. Mwangoka & C.J. Kayombo 63 (MA n.v., MO, P [P05620800]). Lindi: Rondo Plateau, E. Milne-Redhead & P. Taylor 7630 (EA, K). Morogoro: Ulanga district, Kitonga subvillage, 8°46'46"S, 36°42'34"E, G.S. Laizer et al. 1449 (BM, MO). Ruvuma: [near Gumbiro], by R. Mtandazi [river], E. Milne-Redhead & P. Taylor 8538 (B [B 10 0312902], EA, K, LISC, P [P05620801]) and 8539 (EA, P [P05620802]).

22. *Coccinia senensis* (Klotzsch) Cogn. in A.DC. & C.DC., *Monogr. Phan.* 3: 535. 1881.

Cephalandra senensis Klotzsch in W.C.H. Peters, *Naturw. Reise Mossambique*: 151. 1862. Type: Mozambique. [Zambézia Province]: Rios de Sena [province], without detailed locality, in grassland, W.C.H. Peters s.n. (Holotype: B, destroyed).

Type: Tanzania, Lindi Region: 40 km W of Lindi, Lake Lutamba, hill, woodland, climbing over bushes, c. 240 m, male, fl, 6 Sep 1934, H.J.E. Schlieben 5259 (Neotype, designated in Holstein and Renner (2010: 441): M! [M-0165202]; isoneotypes: B! [B 10 0379053, digital image: B], BM!, G! [G00301596], HBG! [HBG518898], MO!, P!, S! [S08-12156, digital image: S], Z!).

Coccinia jatrophiiifolia (A.Rich.) Cogn. var. *australis* Cogn. [sphalm.: *Coccinia jatrophæfolia* var. *australis* Cogn.] Bol. Soc. Brot. ser. 1, 7: 228. 1889.

Type: Mozambique. [Nampula]: Mossuril et Cabaceira (Zambézia), male, fl, 1884, R. de Carvalho 15 (Lectotype, designated here: BR!).

Type: Mozambique. Ibid., male, fl, 1884–1885, R. de Carvalho s.n. (isolectotype: COI!).

Coccinia fernandesiana C. Jeffrey, *Kew Bull.* 30(3): 478. 1975.

Type: Mozambique. Niassa: Erati, between Namapa and Ocúa, near river Lúrio bridge, female, fl, fr, 9 Mar 1960, F. de Lemos & L. Macuácua 29 (Holotype: COI,




Figure 39. Inflorescence of a male *Coccinia senensis*, note the long triangulate (may be narrower in other collections, then subulate) calyx lobes in contrast to the calyx lobes of *C. adoensis* var. *adoensis* in Fig. 21; picture taken from the neotype (*H.J.E. Schlieben 5259* (M)). Black bar equals 1 cm.

isotypes: BM! [BM001010006], BM! [BM001010007], K!, LISC! [LISC 002485, digital image: IICT, JPS], LMA, PRE! [PRE0592949-0, digital image: JPS], PRE! [PRE0592950-0, digital image: JPS], SRGH! [SRGH0106711-1, digital image: JPS], SRGH! [SRGH0106711-2, digital image: JPS], SRGH! [SRGH0106711-3, digital image: JPS]).

Type: Tanzania. Mtwara: Masai Distr. [*sic*, must be Masasi Distr.], W of R. Bangala, 390 m, in woodland on gravelly soil, 17 Dec 1955, *E. Milne-Redhead & P. Taylor 7703* (Paratypes: EA!, K (2)!, LISC!, P!).

Type: Tanzania. Lindi: Mlinguru, 275 m, shrub woodland, 18 Dec 1934, *H.J.E. Schlieben 5745* (Paratypes: B! [B 10 0379052, digital image: B], B!, BR!, EA!, HBG! [HBG518899], K (2)!, P!, LISC!, MO!, PRE!, SRGH).

Type: L. cl., *F. de Lemos & L. Macuácuá 30* (Paratypes: BM! [BM001010005], COI! [2 sheets], K!, LISC! [LISC 002482, digital image: IICT, JPS], LMA, P!, SRGH).

Type: Mozambique. Niassa: Erati, between Namapa and Nacarea, *F.A. Mendonça 1128* (Paratypes: LISC! [LISC 002483, digital image: IICT, JPS]).

Type: Ibid., *F.A. Mendonça 1129* (Paratypes: LISC! [LISC 002484, digital image: IICT, JPS]).

Type: Mozambique. Zambézia: Milange, 95 km towards Quelimane, *A.R. Torre & M.F. Correia 14060* (Paratypes: K! [K000313233, digital image: JPS, K], LISC! [LISC 002481, digital image: IICT, JPS]).

Type: Tanzania. Lindi: Nachingwea, *Pterocarpus-Combretum* woodland, *B. Anderson 815* (Paratypes: EA!, K!, NHT!).

Type: Tanzania. Lindi: Mbemkuru [also called Mbwenburu, Mto Bwamkuro], in deciduous thicket by roadside, 135 m, *E. Milne-Redhead & P. Taylor 7473* (Paratypes:

BR!, EA!, K (2)!, P!), *E. Milne-Redhead* & *P. Taylor* 7473A (Paratypes: K (2)!), *E. Milne-Redhead* & *P. Taylor* 7473B (Paratype: K), *E. Milne-Redhead* & *P. Taylor* 7473C (Paratype: K), *E. Milne-Redhead* & *P. Taylor* 7473D (Paratype: K).

Coccinia subglabra C.Jeffrey, Kew Bull. 30(3): 479. 1975.

Type: Mozambique. Nampula: Nacala, 11 km from Itoculo towards Nacala, 130 m, male, fl, 4 Dec 1963, *A.R. Torre* & *J. Paiva* 9417 (Holotype: LISC [LISC 002486, digital image: IICT, JPS]; isotypes: COI, K! [K000313458, digital image: JPS, K], LMA).

Type: Mozambique. Ibid., *A.R. Torre* & *J. Paiva* 9417A (Paratypes: COI, K! [K000313457, digital image: JPS, K], LISC, LMA).

Type: Mozambique. Niassa: Ruvuma River, *J. Kirk* s.n. (Paratype: K!).

Type: Mozambique. Nampula: Mossuril e Cabaceira, *R. de Carvalho* s.n. (Paratypes: COI (2)!).

Type: Mozambique. Zambézia: 23 km on road [from vila] Maganja da Costa [= Olinga] towards Namacurra, *A.R. Torre* & *M.F. Correia* 14176 (Paratypes: EA!, LISC [LISC 002487, digital image: IICT, JPS], MO!).

Description. Perennial climber or creeper. Stems up to 3 m, glabrous or with erect, stiff, articulate, pale trichomes, glabrescent, when older sometimes with white pustules. Leaves subsessile or distinctly but not long petiolate. Petiole 0.4–4 cm long, abaxial side more or less densely covered with erect, stiff, articulate, pale trichomes, sometimes glabrous. Leaves 4–14 × 5–16 cm, cordate or shallowly to deeply 3- or 5-lobate, sometimes auriculate (Fig. 6). Lobes triangulate, lanceolate to lineal, sometimes with broader end. Finely to coarsely serrate-dentate, especially towards the lobe tips, sometimes lobulate. Lobe apices acute to subacute, with an apical tooth. Upper leaf surface with erect, stiff, articulate, pale trichomes or with whitish pustules. Lower leaf surface glabrous or covered with erect, stiff, articulate, pale trichomes, nerves with same indumentum or with white pustules. Between the main nerves at lamina base often with blackish glands. Probracts up to 4 mm. Tendrils simple, rarely bifid. Male flowers (Fig. 39) in few-flowered racemes, sometimes accompanied with 1–2 solitary flowers. Common peduncle 0.3–6.2 cm long, often more or less densely covered with erect, stiff, articulate, pale trichomes. Petiole of flowers in racemes 0.2–1.2 cm. Bracts up to 2.5 mm, often missing. Petioles of solitary flowers 3.2–6.2 cm long. Petioles in each case with indumentum of peduncle. Perianth tube glabrous or with erect, stiff, articulate, pale trichomes. Calyx lobes 2.5–6.5 mm long, subulate to narrowly triangulate-lanceolate. Corolla 1.1–3 cm, yellow, orange to salmon, lobes 0.9–1.1 cm. Color of filament column, anther head, and pollen sacs not seen. Pedicel of female flower 1.2–2.3 cm. Hypanthium, calyx lobes, and corolla like in male flowers. Ovary glabrous or more or less covered with erect, articulate trichomes. Style and stigma not seen. Fruit 3.5–4.4 × 1–1.5 cm, long ovoid to shortly cylindrical, often with apical sterile tip. Unripe pale green with dark green longitudinal lines, ripe red. Seeds 5.5–7 × 3–5 × 1.2–1.3 mm (L/W/H), symmetrically obovate, face lenticular.

Phenology. Flowering time: January–April, September, December.

Distribution. Fig. 23. Central Tanzania (Iringa, Lindi, Morogoro, Ruvuma), Malawi (Southern Region), Mozambique (Cabo Delgado, Nampula, Tete, Zambezia). Elevation 0–700 m. Sandy soil. *Coccinia senensis* seems to be a typical element of the Zambezian center of endemism (White 1983b). Deciduous woodlands, termitaria, riverine thickets, coastal forests.

Vernacular names. Kihehe: mtumbulansoka (W. Carmichael 171), Macua [Makhuwa]: muuco-uco (F. de Lemos & L. Macuácuá 29)

Remarks. The species is recognizable by the combination of few-flowered racemes, long subulate calyx lobes, and the often subsessile leaves. The trichome type (often appearing articulate when dried) is the same as in *C. rehmannii*, where (sub-)glabrous collections also occur (see also the Taxonomic remarks). Except for the degree of trichome density, a subglabrous collection (E.M.C. Groenendijk et al. 1031) from 11 km from the collecting site of the *C. subglabra* holotype was neither morphologically nor genetically (Holstein and Renner 2011b) distinguishable from *C. fernandesiana*, and *C. senensis* (sensu Flora Zambesiaca (Jeffrey 1978) and sensu Holstein and Renner (2010)). Fruit shape and length as well as the length of the female pedicel are variable, so *C. subglabra* is synonymized. Without calyx lobes, *C. senensis* is only hardly, if at all, distinguishable from glabrous *C. adoensis* collections or those with long articulate trichomes, which are described as *C. adoensis* var. *jeffreyana* in this treatment. The fruit and seed shape also match the variable *C. adoensis*. Usually, *C. adoensis* var. *adoensis* has short trichomes and calyx lobes are ≤ 2 mm, but where both species meet (Malawi, NW Mozambique, S Tanzania), exceptions can be found (listed and further discussed as *C. adoensis* var. *jeffreyana*). Whether *C. adoensis* and *C. senensis* are truly separate species and the role of these intermediates needs to be tested by artificial hybridization, field observations, and/or a phylogeographic analysis.

Taxonomic remarks. Although the holotype of *C. senensis* burned during the destruction of the Berlin herbarium in 1943, and the name appears to have been lost, the protologue mentions several characters that allow *C. senensis* to be synonymized with Jeffrey's *C. fernandesiana*. The *C. senensis* protologue points out “articulate” trichomes and an overall appearance like *C. quinqueloba*, which matches perfectly with many collections of *C. fernandesiana*. Interestingly, many of these collections have been identified as “*Coccinia quinqueloba*” or “*Coccinia palmata*” by various collectors and scientists. The similarity, including the calyx lobes, is easily visible in many collections, but both species are restricted to southern Africa.

Cogniaux described var. *australis* of *C. jatrophiifolia* (synonym to *C. adoensis*) recognizing the similarity to the polymorphic *C. adoensis*. However, he differentiated between the R. de Carvalho specimens with long lineal lobes (BR, COI) and specimens with lanceolate lobes (BR, COI), which he determined as *C. senensis*. When Jeffrey described *C. subglabra*, he cited the two COI specimens (as deduced from his ID labels), but he did not refer to Cogniaux' variety, which must have been overlooked. The one COI specimen is therefore paratype of *C. subglabra* and syntype of *C. jatrophiifolia* var. *australis*. The two COI specimens are also misplaced paratypes of Meeuse's *C. rehmannii* var. *littoralis*. The similarity of the COI specimens of Meeuse's variety to *C. senensis*

is striking, but the long peduncles and the conspicuous black sublaminar glands refer rather to *C. senensis* than to *C. rehmannii*.

Specimens examined. (Selection, in total: 49) Malawi. Northern Region: Rumphi district, Nyika Plateau, 20 mls N of M1, *J. Pawek 13339B* (MO). Southern Region: Bvumbwe, *I.F. La Croix 2653* (MO); Lengwe National Park, near Mukanyu ravine, *A. Hall Marker 1051* (K). Mozambique. Cabo Delgado: Mueda Plateau, 11°22'S, 39°20'E, *W.R.Q. Luke et al. 10084* (EA, K). Nampula: Monapo district, Monapo, forest reserve of Mr. Wolf, *E.M.C. Groenendijk et al. 1031* (WAG [WAG0104327]). Tete: Cabora bassa [Cahora bassa], police post no. 3, 5 km from barrage, *A.R. Torre et al. 18788* (MO). Tanzania. Lindi: Selous Game Reserve, Kingupira, 8°28'S, 38°33'E, *K. Vollesen 1908* (EA); *ibid.*, *K. Vollesen MRC 4316* (DSM, EA, K, WAG [WAG0234144]); *ibid.*, *R.C. Wingfield et al. 3466* (DSM). Morogoro: Kilosa district, Ilonga Research Institute, 9.5 km NNE of Kilosa on road to Dumila, 6°46'31.3"S 37°2'23"E, *N. Holstein et al. 66* (DSM, M).

23a. *Coccinia sessilifolia* (Sond.) Cogn. var. *sessilifolia*

Cephalandra sessilifolia Sond. in Harv. & Sond., Fl. Cap. 2: 493. 1862. *Coccinia sessilifolia* (Sond.) Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 534. 1881.

Type: South Africa. Vaal river, *J. Burke 289* (Syntypes: K! [K000313207, digital image: K], PRE, SAM).

Type: South Africa. Slengerfontein in Nieuwe Hantom [area where the provinces Western Cape, Eastern Cape and Free State meet], on rocks, 4500–5000 ft, female, fr, 1839, *J.F. Drège 3375* (Lectotype, designated here: P! [P00346268, digital image: P]).

Type: South Africa. Nieuwe Hantom, on rocks, 4500–5000 ft, 1839, *J.F. Drège s.n.* (Syntypes: BR! [BR0000005111596, digital image: BR], G! [G00301769], K! [K000313206, digital image: K], L!, P! [P00346270, digital image: P], S! [S08-12468, digital image: JPS, S], W! [W 0026938: digital image: WU]).

Type: South Africa. Transvaal, *C.L.P. Zeyher 580* (Syntypes: BM! p.p., E! [E00303259], K! p.p. [K000313205, digital image: K], P! p.p. [P00346271, digital image: P]).

Coccinia sessilifolia var. *major* Cogn. in Schinz, Verh. Bot. Ver. Provinz Brandenburg 30: 152. 1889.

Type: Namibia. Hereroland, male, fl, 1885, *A. Lüderitz 133* (Lectotype, designated here: Z!).

Type: Namibia. Walvis bay to Odyitambi, Dec 1885–Feb 1886, *A. Lüderitz 1a* (Syntype: Z!).

Coccinia schinzii Cogn., Bull. Herb. Boiss. 3: 419. 1895.

Type: South Africa. Transvaal: Klippan [according to Meuse (1962) in Limpopo: Greater Sikhukhune District Municipality, Doornpoort; 24°37'S, 29°26'E], bushveld, 1875–1880, *A. Rehmann 5162* (Lectotype, designated by Meuse (1962:




Figure 40. Distribution map of *C. sessilifolia* var. *sessilifolia* (pale yellow dots; based on 133 collections) and *C. sessilifolia* var. *variifolia* (blue dots; based on 7 collections). For South Africa the borders of the provinces are given.

98): Z! [Z-000004446, digital image: Z], isoelectotype: BR! [BR0000005112265, digital image: BR, JPS]).

Description. Perennial climber or creeper. Stems up to 5 m long, with slight waxy cover, glaucous (Figs 2b, 7b), glabrous (first shoots may have short, white trichomes). Leaves sessile to amplexicaul (first leaves after appearance of stem can be distinctly petiolate, rarely also when mature (up to 1.5 cm; Figs 2b, 4b, 7b)), glaucous, $1.5\text{--}12.5 \times 2.2\text{--}13.5$ cm, (cordate to) deeply palmately 5-lobate. Lobes linear, lanceolate to elliptic. Leaf margin remotely denticulate, with or without lobules. Lobe apex obtuse to acute, apiculate. Upper leaf surface glabrous, clear to white pustulate. Lower leaf surface glabrous, sometimes with dark glands near base of lamina. Probracts up to 1.7 mm or missing. Tendrils simple, very rarely bifid. Male flowers solitary or clustered in few-(rarely many-)flowered racemes (Figs 2b, 7b). Pedicels of solitary flowers 1–4 cm, glabrous. Peduncle 1–6 cm long, glabrous. Pedicels of flowers in racemes 0.3–2.5 cm, glabrous. Bracts glabrous, up to 1.8 mm,

or missing. Perianth tube glabrous, calyx lobes 1.5–3.5 mm long, lanceolate to (narrow) triangulate, erect to reflexed. Corolla 1.5–3 cm long, whitish cream to pale yellow, rarely dull orange-brown with conspicuous green venation, lobes 0.9–2 cm. Filament column not seen, anther head not seen, pollen sacs yellow. Female flowers one solitary. Pedicel 1–3 cm, glabrous. Hypanthium glabrous, calyx and corolla like in males. Ovary glabrous. Style columnar, greenish yellow. Stigmas bulging, greenish yellow. Fruit 8–12 × 3–4 cm, ellipsoid to oblong, when immature green with white longitudinal spots to stripes with waxy bloom, ripe red (Fig. 13b). Seeds 6–8 × 3–3.5 × 1–1.5 mm (L/W/H), symmetrically to slightly asymmetrically obovate, face flat (Fig. 14d, e).

Phenology. Flowering time: January–May, October–December.

Distribution. Fig. 40. South Africa, Botswana, Namibia, except high mountains, hyper-arid regions, and Cape floristic region. Elevation 300–1500 m. Stony soil, sand, sandy loam, clay loam. Granite or calcareous substrate. Semi-desert, grassland (e.g., *Rhynchelytrum* sp.), bushland, open woodland. Full sun to shade. Moderate disturbance tolerated.

Use. Unripe fruits are baked in ashes and eaten (Dinter 1912), ripe fruits edible (*N. Holstein* 119, *R.H.W. Seydel* 3829, *R.H.W. Seydel* 4100).

Vernacular names. Afrikaans: bobbejaan komkommer (*C.A. Smith* 3981), Otjiherero: ekungu (singular), omakungu (plural) (Dinter 1912; 1919/20), Tswana: mogábalá (*J. Snyman & C. Noailles* 229), !Kun [Kung]: kitwa (*R. Story* 5167).

Taxonomic remarks. The *C.L.P. Zeyher* 580 specimens (syntype) in BM, K, and P are mixed with a *Trochomeria* sp.

Specimens examined. (Selection, in total: 184) Botswana. Central District: Mahalapye, 2 mls [3.8 km] SW of Kalamare, *H.J. van Rensburg* B4019 (PRE). Kgatleng: 15 km SE of Artesia (Mosomane), *D.T. Cole* 1542 (PRE). North-West District: Aha Hills, *H. Wild & R.B. Drummond* 6953 (COI). South-East District: Lobatsi [Lobatse], *F.A. Rogers* 6281 (G, Z). Namibia. Erongo: Karibib, Okongawa, *R.H.W. Seydel* 3026 (B (2), COI, FR (2), G (3), H, HEID, M, WAG [WAG0234195], WAG [WAG0234197]). Khomas: [Farm] Aris, mountain in the west, *R.H.W. Seydel* 4100 (B (2), M, MO) and 4100a (B (2)). Oshana: Amboland, Uukuanjama [Oukwanyama], Omupanda, *A. Wulffhorst* 18 (Z). Otjozondjupa: 32 mls [51.2 km] N of Nurugas on road to Karakuwisa, *B. de Winter* 3710 (M, PRE). South Africa. Eastern Cape: in valley near Graaff-Reinet, *H. Bolus* 364 (S [S08-12381]). Free State: Kroonstad townland, NE of confluence of Blomspruit and Vals river, *J.C. Scheepers* 1720 (EA, LISU, PRE, S [S08-12461]). Gauteng: Pretoria, Brummeria, Botanical Garden, *A. Balsinhas* 3406 (MO, PRE, WAG [WAG0234191]); *ibid.*, *A. Balsinhas* 3476 (MO, PRE, WAG [WAG0234190]). Limpopo: Penge mine, *E. Retief* 1354 (MO, PRE, WAG [WAG0234188]); c. 30 mls [48 km] W of Louis Trichardt, western part of Zoutpansberg, near Mara, Buysdorp, *H.J.E. Schlieben* 7453 (B, G, HBG, M). Mpumalanga: Blyderivierspoort Nature Reserve, Sybrand van Niekerk resort at camp area, *E. Retief* 1340 (PRE, MO, WAG [WAG0234187]), *ibid.*, *E. Retief* 1341 (PRE). Northern Cape: Colesburg, Achtertang, 16 Apr 1934, *J.O. Swinford* s.n. (PRE [PRE42988]). North West: near Klerksdorp,

H.J.E. Schlieben 10695 (HEID, PRE, S [S08-12463]); 60 mls [96 km] NW of Vryburg, Farm Palmyra, *R.J. Rodin 3605* (MO, P [P05620787], PRE).

23b. *Coccinia sessilifolia* var. *variifolia* (A.Meeuse) Holstein, stat. nov.

urn:lsid:ipni.org:names:77148916-1

Coccinia variifolia A.Meeuse, Bothalia 8: 100. 1962.

Type: South Africa. [Limpopo]: Waterberg, Vaalwater, about 2.25 km from Vaalwater on road to Hermanusdoorns, male, fl, 6 Jan 1959, *A.D.J. Meeuse & R.G. Strey 10413* (Holotype: PRE [2 sheets: PRE0188239-1 and PRE0188239-2, digital image: JPS], isotypes: BOL, L, SRGH).

Type: South Africa. Ibid., *A.D.J. Meeuse & R.G. Strey 10413bis* (Paratype: PRE!).

Type: South Africa. Limpopo: Palala river, *M.G. Breyer[-Brandwijk] TRV25226* (Paratype: ?).

Type: South Africa. Limpopo: Rietspruit near Nylstroom [Modimolle], *G.P.F. van Dam TRV23372* (Paratype: PRE!).

Type: South Africa. Limpopo: Nabomspruit, Mosdene, *E.E. Galpin s.n.* (Paratype: ?).

Type: South Africa. Limpopo: 11 km from Warmbaths [Bela Bela] on Nylstroom road, *R. Story 1525* (Paratype: ?).

Type: South Africa. Limpopo: Warmbaths, c. 3600 ft [1100 m], grassland/bush veld, *H. Bolus 11893* (Paratype: BR!).

Type: South Africa. Ibid., *R. Leendertz TRV7579* (Paratype: ?).

Type: South Africa. [Limpopo]: Waterberg, 5.5 mls [8.85 km] NNE of Warmbaths, c. 1220 m, sour bushveld, *J.P.H. Acocks 13903* (Paratype: S! [S08-12475]).

Description. Perennial climber. Stems up to 1.2 m, likely also longer, glabrous. Petiole 0.7–1.6 cm, glabrous. Leaves 5.2–6 × 6–7.5 cm, deeply to shallowly 5-lobate, lobes outwards lobulate. Leaf margin remotely dentate, apex obtuse with final tooth. Upper leaf surface glabrous, with clear to whitish pustules. Lower leaf surface glabrous, with glands at base between nerves. Probracts up to 2 mm. Tendrils simple. Male flowers in racemes, accompanied by a solitary flower. Common peduncle 1–1.4 cm, pedicel in racemes 3–6 mm, each glabrous. Bracts up to 1.5 mm, narrowly ovate. Pedicel of solitary flowers 0.9–2 cm, glabrous. Perianth tube glabrous. Calyx lobes 2.5–4 mm, subulate to narrowly triangulate, erect. Corolla c. 2 cm, pale buff, lobes not measured. Filament column, anther head, and pollen sacs not seen. Female flowers solitary or clustered in reduced 2-flowered racemes. Common peduncle 1 mm, pedicel in racemes 0.9–1 mm, pedicel of solitary flower not seen, each glabrous. Hypanthium most likely glabrous and perianth as in male flowers. Ovary glabrous. Style and stigma not seen. Fruit and seeds not seen.

Phenology. Flowering time: January–March, November, December. Likely as in *C. sessilifolia* var. *sessilifolia*.

Distribution. Fig. 40. Only known from Limpopo Province in South Africa. Elevation 800–1200 m. On sandstone, well-drained stony sand. Low closed woodland.

Remarks. The new status of *C. variifolia* was chosen due to the minor differences to *C. sessilifolia* s.str. Subsessile leaves spontaneously occur in *C. sessilifolia* (H. Bolus 364, F.A. Rogers 19262) and young individuals usually (always?) have petiolate leaves (N. Holstein 131, Fig. 4b). However, these collections of mature plants with distinctly petiolate leaves have only been observed in Limpopo Province in South Africa. Sessile *C. sessilifolia* leaves can be quite variable, profoundly to deeply lobate, sometimes also lobulate. Compared to the rather uniform *C. quinqueloba* (Meeuse 1962), the leaves thus appear to be extraordinarily variable. Meeuse's *C. variifolia* shares the sublaminar glands (cp. Fig. 7b) and the calyx lobes of *C. sessilifolia*, and it is geographically nested within this species (hence no climatic differentiation). Acocks (J.P.H. Acocks 13903) also reports a “stark glaucous” appearance, just as in *C. sessilifolia*. As petiolate leaves also occur in young *C. sessilifolia* plants, and subsessile leaves also occur in mature plants, it is more likely that the distinctly petiolate *C. sessilifolia* individuals represent a local fixation of this character. As *C. sessilifolia* var. *sessilifolia* is derived from petiolate plants, this variety might even represent a remnant population of these.

Specimens examined. (in total: 8) South Africa. Limpopo: Waterberg Distr., F.A. Rogers 24932 (Z [Z-000073427]), N. Rooyen 1667 (PRE), R.H. Westfall 2136 (PRE).

24. *Coccinia subsessiliflora* Cogn., Bull. Jard. Bot. État Brux. 4(1): 225. 1914.

Type: D. R. Congo. [Équateur Province]: around Likimi, male, fl, 15 Oct 1910, L.C.E. Malchair 433 (Lectotype, designated here: BR! [BR0000008887481, digital image: BR, JPS], syntype: BR! [BR0000008886835, digital image: BR, JPS]). *Coccinia* sp. D in C. Jeffrey, F. T. E. A.: 70. 1967. Uganda. [Western Region]: Kigezi District [(Kanungu District/Kisoro District)], Kayonza Forest Reserve [Bwindi Forest Reserve / Impenetrable Central Forest Reserve], S. Paulo 644 (EA!, K!, MO!); [Central Region]: Mengo district, Mabira forest, M.V. Loveridge 87 (?); [Central Region]: Mabira Forest, near Kiwala, R.A. Dummer 3195 (?).

Description. Perennial creeper or climber. Stems up to 4 m, glabrous. Petiole 2–12 cm, glabrous, sometimes with white pustules. Leaves 5.5–15 × 6.5–17.5 cm, almost to the base palmately 5-lobate. Lobes lanceolate, sometimes lobulate; tip acute, acuminate. Margin serrate-lobulate, denticulate. Upper leaf surface glabrous with clear to white pustules, rarely with few fine (up to 1.5 mm long) trichomes. Lower leaf surface glabrous, rarely with dispersed small blackish glands, rarely with tiny trichomes; sometimes nerves with white pustules. Probracts up to 1.5 mm or missing. Tendrils simple. Male flowers in glabrous, dense, compact racemes. Peduncles up to 6 mm. Pedicels up to 4 mm. Bracts 2–2.5 mm. Perianth tube glabrous. Calyx lobes 1–2 mm, subulate, triangulate to lineal, erect to reflexed. Corolla 1.2–1.3 cm, orange, pale yellow-orange, yellow, lobes c. 3 mm. Filament column, anther head, and pollen sacs not seen. Female flowers solitary or in few flowered racemes. Common peduncle up to 1 cm, glabrous. Pedicel of flowers in racemes up to 4 mm. Bracts up to 2 mm or missing.




Figure 41. Distribution map of *C. subsessiliflora* (blue dots; based on 18 collections). For D.R. Congo the borders of the provinces (until 1988) are given.

Pedicle of solitary flowers up to 1.1 cm, glabrous. Hypanthium glabrous, calyx lobes and corolla like in males. Ovary glabrous. Style columnar, pale yellow. Stigma 2-lobed, yellow. Fruit 2–2.4(–7) × 1.7 cm, globose to long ovoid, unripe green with glaucous waxy cover, ripe color not known, most likely red. Seeds $\geq 4.5 \times 2\text{--}2.5 \times 1\text{--}1.5$ mm (L/W/H), asymmetrically obovate, face flat.

Phenology. Flowering time: January, April, July, August, October, December.

Distribution. Fig. 41. Congo basin (Central African Republic, D. R. Congo). Forested mountains of NW Burundi, D. R. Congo (North Kivu, South Kivu), W Rwanda, Uganda (Western Province), South Sudan (Eastern Equatoria/Sharq al-'Istiwa'iyah: Lotti Forest). Elevation 300–1950 m. Soil preferences unknown. Tropical rainforests. *Macaranga* [= *Gilbertiodendron*?] forest, swamp forest, disturbed ground in open forest.

Vernacular names. Kihunde: mutangatanga (*R. Gutzwiller* 965), Lissongo [Mapti]: kanganga (*C. Tisserant (Équipe)* 1103), Turumbu: ndombo (*J. Louis* 2709).

Taxonomic remarks. There are two *Malchair* 433 specimens. As they do not contain any indication of having been separated from a single specimen, they are treated as syntypes. The two specimens do not differ in quality of the material, so the specimen with the original label was chosen to be the lectotype.

Remarks. Collections from the eastern parts of the distribution (esp. E of the Western Rift) have longer fruits but it appears to be a variable character.

Rarely (*J. Louis* 5672, *J. Louis* 13030), the lower leaf lamina and the adaxial petiole side have short trichomes and the upper lamina has some long trichomes. These features are unusual, but the other characters match the species.

Although *C. subsessiliflora* is nested within *C. barteri* in the molecular tree from plastid markers (Fig. 17), it can be regarded as a proper morphospecies. The deeply lobate leaves are a distinct character, and the species is distributed only in the Congo Basin and the eastern rainforests. The present author supposes that *C. subsessiliflora* might have evolved peripatrically in an arid period of the Pliocene/early Pleistocene period, eventually near the Kivu Mts, and *C. barteri* populated these areas later on.

Specimens examined. (Selection; in total: 21) Burundi. Bubanza: Bubanza, *J. Lewalle* 6504 (BR, EA); *ibid.*, *M. Reekmans* 1477 (BR (2)). Central African Republic. Lobaye: Boukoko, *C. Tisserant (Équipe)* 1103 (BM, G, P [P05620797], P [P05620798]); *ibid.*, 2276 (BM, G, P [P05620796], P [P05620799], P [P05621163]). D.R. Congo. Équateur: [Nord-Ubangi district] Businga territoire, between Karawa and Businga, *J. Lebrun* 1928 (BR, P [P05620794], WAG [WAG0225403]). Maniema: between Kindu and Katakokombe, *J. Lebrun* 6011 (P [P05620791], WAG [WAG0225402]). North Kivu: Beni territory, Kiandolili river, Gongobotsi Camp of Albert National Park guards, *H. Fredericq in Herb. G.F. de Witte* 8288 (BR, M, PRE, WAG [WAG0225407]). Orientale: Haut-Uélé district, Faradje territoire, Kurukwata (Aba), *P. Gerard* 3564 (BR, EA, WAG [WAG0225405]); Ituri district, Mambasa territoire, Réserve de Faune à Okapi, Epulu, 1°25'N, 28°35'E, *C.E.N. Ewango* 2290 (M, MO); Yangambi, *J. Louis* 13030 (BR [BR0000008912916], BR [BR0000008913272], P [P05620795], WAG [WAG0225404]). South Sudan. Eastern Equatoria: Torit district, Lotti Forest, *J.K. Jackson* 3026 (K). Uganda. Western Region: [Masindi district], Bunyoro, Bujenje county, Budongo Forest, *A.B. Katende* K2801 (MO); *ibid.* *T.J. Synott* 1322 (EA).

25. *Coccinia trilobata* (Cogn.) C. Jeffrey, *Kew Bull.* 15: 349. 1962.

Peponia parviflora var. *trilobata* Cogn., Bot. Jahrb. Syst. 21: 210. 1895. *Peponia trilobata* (Cogn.) Engl., Pflanzenw. Ost-Afr. C: 399. 1895, nom. illeg. [*Peponia* is a diatom genus]. *Peponium trilobatum* (Cogn.) Engl., Engl. & Prantl, Pflanzenfam., Nachtr.: 318. 1897.

Type: Tanzania. Kilimanjaro: Mkuu [c. 3°10'S, 37°36'E], 1500 m, in hedges, fl, fr, Mar 1894, *G. Volkens* 1956 (Holotype: B, destroyed; lectotype acc. to sheet, but not published, so designated here: BR! [BR0000008887160, digital image: BR, JPS]; isolectotype: BR!).

Coccinia kilimandjarica Cogn. ex Harms in Fries, Notizbl. Bot. Gart. Berlin-Dahlem 8: 489. 1923.

Type: Tanzania. Kilimanjaro: Kibohöhe [farm at c. 3°15'50"S, 37°12'0"E], 1100–1200 m, fl, *R. Endlich* 52a (Holotype: B, destroyed; lectotype, designated here: M! [M0105772, “1122”, digital image: JPS], isolectotype: H!).

Coccinia kilimandjarica var. *subintegrifolia* Cogn. ex Harms in Fries, Notizbl. Bot. Gart. Berlin-Dahlem 8: 490. 1923.

Type: Tanzania. Kilimanjaro: Kibohöhe, 1100–1200 m, fl, *R. Endlich* 52 (Holotype: B, destroyed; lectotype, designated here: M! [M0105773, “1121”, digital image: JPS], isolectotype: H!).

Description. Perennial climber. Stems up to 3 m, with soft, whitish trichomes, at least along nerves. Petiole 1.5–16.5 cm, with whitish trichomes (Fig. 8b). Leaves 2.6–14.5 × 3.2–18 cm, cordate, 5-angulate to 5-lobate, sometimes lobulate. Margin denticulate. Apex at least of central lobe acute, often acuminate. Upper leaf surface with hyaline to white pustules and usually with white trichomes. Lower leaf surface more or less covered with soft trichomes, denser on nerves. Probracts up to 1.5 mm or missing. Tendrils simple. Male flowers solitary or in short few-flowered racemes. Common peduncle up to 1.7 cm long, pedicel in raceme up to 3.2 cm, each glabrous or with short trichomes. Bracts up to 1 mm. Pedicel in solitary flowers 0.7–4.7 cm, indumentum as in racemes. Perianth tube with articulate trichomes. Calyx lobes 2–5 mm long, lineal, erect to reflexed. Corolla 0.7–2.2 cm, orange-yellow to reddish-orange, deeper colored on the inner side of the lobes, outside with green venation, lobes 0.7–1.3 cm. Filament column pale yellowish green, anther head a bit darker than filament column. Pollen sacs yellow. Female flowers 1(–2) solitary, pedicel 0.5–4 cm, glabrous or sparsely covered with short trichomes. Hypanthium with articulate trichomes, calyx lobes and corolla like in males. Ovary with trichomes, becoming glabrous towards fruit ripening. Style and stigmas not seen. Fruits ovoid to oblong, 4–4.7(–9) × 2 cm, unripe green with white longitudinal stripes that develop a dark green corona during ripening, ripe fruits orange-red. Seeds 6.5–7 × 2.5–3.5 × 1.5 mm (L/W/H), more or less asymmetrically obovate, face flat (Fig. 14c).

Phenology. Flowering time: January, May–July, October–December.

Distribution. Fig. 33. Tanzania (Arusha, Kilimanjaro, Tanga), Kenya (Central, Coast, Eastern, Nairobi, southern Rift Valley Province). Elevation 1100–2100 m. Red soil, black soil. Open forest, savannas, evergreen bushland, shrubland, grassland.

Use. Leaves eaten as vegetable (*Coilly?* 24, *F. Msajiri* 19).

Vernacular names. Dholuo: angwe (*G.R. Williams* 307), Kinandi: notondwe (*G.R. Williams* 307), Meru: katakeru (*Coilly?* 24), Kikuyu: kigerema (*P. Njogu* EA13835), Kipare: itotwe (*W.J. Kindeketa* 648).

Remarks. The fruits are reported to be poisonous (*G.R. Williams* 307).

There are some collections that have a mixed (not intermediate) phenotype with *C. microphylla*: the calyx lobes are unusually long (up to 7 mm), which is a strong argument for *C. trilobata*, but the indumentum matches *C. microphylla*. These morphs do not occur in single location but are found in the Ndoto Mts (*O. Kerfoot* 2644), in Koboko (*P. Kirika et al.* 002/020/2011), and around Voi (*M. Hucks* 579, *B. Verdcourt* 3888, *R. Polhill & S. Paulo* 962). Whether these are hybrids (F2 or later) or just a variation is not known. These collections look also quite like *C. megarrhiza*, which occurs in northern Kenya and Ethiopia, however, the indumentum does not match either. A clarification where these collections belong to would require sequence data and a better understanding of the plastid and nuclear haplotypes in the three species, which is not available so far.

The collections in the Usambara Mts are often quite glabrous or the trichomes are minute and thus easy to mix up with *C. microphylla*.

Specimens examined. (Selection, in total: 63) Kenya. Central Province: South Nyeri district, S of road (D450), c. 4 km E of Nairobi–Nanyuki road, 3 km N of Kiganjo, *S.S. Hooper & C.C. Townsend* 1697 (K [K000353353]). Eastern Province:

Nkunga Crater Lake, *P.A. Luke et al.* 7256 (EA). Nairobi: Nairobi river Valley, Chirromo, 1°16'30"S, 36°48'E, *R.B. Faden & A.J. Faden* 74/822 (BR, DSM, EA, MO, WAG [WAG0234201]). Rift Valley Province: Naivasha District [Nakuru district], Ol Longonot Estate, *O. Kerfoot* 3543 (EA, S [S08-12472]). Tanzania. Arusha: Small Momela Lake, *H.M. Richards* 20036 (EA (2), K [K000353413]). Kilimanjaro: valley slopes near Alt Moschi [Old Moshi], *A. Peter* 56453 (B). Tanga: western Usambara Mts, Mombo–Soni road, *R.B. Drummond & J.H. Hemsley* 3007 (B, EA, K, LISC, S [S08-12484]).

Insufficiently known taxa

***Coccinia* sp. A C.Jeffrey, Fl. Zambes.: 450. 1978.**

Zambia. Northern Province: Chilongowelo, Tasker's Deviation waterfall, 4900 ft, female, fl, 27 Feb 1952, *H.M. Richards* 883 (K!).

Distribution. Only known from single collection.

Remarks. Jeffrey (1978) suggested that this collection is allied with *C. barteri*. The inflorescence matches that of *C. racemiflora*, which differs in an urceolate perianth tube/hypanthium, more coriaceous leaves, and a glabrous surface. Simple tendrils also occur in *C. barteri*, so this is not a good distinctive character. As the plastid haplotypes of *C. racemiflora* are nested in *C. barteri*, and the distribution of *C. sp. A* is within the *C. barteri* range, Jeffrey's hypothesis is sound. However, the corolla lobes are quite long, and it seems that the corolla is open campanulate. Therefore, there is some similarity with *C. mildbraedii* and even with *C. grandiflora*.

***Coccinia* sp. B C.Jeffrey, Fl. Zambes.: 450. 1978.**

Zambia. Southern Province: Mazabuka, on Nanga Estate near Kafue pilot polder [c. 15°45'S, 27°54'E], female, fl, 7 Mar 1963, *H.J. van Rensburg* 1620 (K!).

Distribution. Only known from single collection.

Remarks. Like *C. sp. A*, Jeffrey (1978) suggested a closer relationship to *C. barteri*. The tubular corolla and the coriaceous leaves support that. The collection was found in riverine bush in *Acacia* woodland and *Hyparrhenia/Setaria* grassland in a flood plain area, which is unusual for *C. barteri* as it rather occurs in (rain) forests. White (1983a) calls the phytochorion of that region “edaphic grassland with semi-aquatic vegetation”. On the one hand, the local soil conditions are not known, and water might be available throughout the year, on the other hand this individual might also represent a local adaptation towards increased drought tolerance.

Dubious names

***Coccinia aostae* Buscal. & Muschl., Bot. Jahrb. Syst. 59: 499. 1913.**

Type: [Eastern Africa]. At Mbusi river [authors state that this river flows into the Indian Ocean in Mozambique, the collector went upstream towards Zambezi river and Victoria falls; most likely the Buzi River is meant], tree steppe, fl, 14 Dec 1909, *H. von Aosta* [H.L.F.H. d'Orléans] 105 (Syntype: B destroyed; duplicate ?).

This species is supposed to be from Mozambique. However, the describing author, Muschler, provoked a scandal with this work as Georg Schweinfurth (1915) and his former supervisor Adolf Engler (Engler et al. 1915; Ryding 2001) accused him of fraud. Gilg, who contributed corrections in the Cucurbitaceae, suggested that *Coccinia aostae* had been described using the *G.A. Schweinfurth* 578 specimen from Eritrea, which bore the ms. name *Coccinia lalambae* Schweinf. A drawing of this species exists in BR! (K neg. 4887), which likely represents a *C. adoensis*. However, the name for *G.A. Schweinfurth* 578 remains unpublished and the *von Aosta* 105 specimen is destroyed, and the name remains dubious. According to the describing authors, duplicates of the von Aosta specimens have been distributed, and White (1962) found some in Florence (FI or FT). However, a loan from FT did not contain any *Coccinia* collections by von Aosta.

***Coccinia buettneriana* Cogn., Bull. Acad. Roy. Sci. Belgique ser. 3, 14: 351. 1887.**

Type: Gabon. No detailed information given, Sep 1884, *R. Büttner* 18 (Holotype: B destroyed).

Cogniaux and Harms synonymize (1924) *C. buettneriana* under *Momordica gabonii* Cogn. (1881) as it was collected in close vicinity of *Büttner* 17 (*Momordica gabonii*), which is, according to Cogniaux himself almost not distinguishable from *C. buettneriana*.

***Coccinia calantha* Gilg, Bot. Jahrb. Syst. 34: 358. 1904.**

Type: Tanzania. [Tanga]: Usambara Mts, Duga, near Nikunde village, 100 m, in bush and on fencing, fl, Jul, *C.H.E. W. Holst* 3190 (Holotype: B destroyed; duplicates ?).

As the holotype is destroyed and the description does not give enough sufficient characters to relate *C. calantha* to other species, the name remains dubious. Zimmermann (1922b) presents a drawing of an anther, but the thecae are too narrow for a *Coccinia* but would match *Eureiandra* species. On the other hand, *Eureiandra* has free petals, whereas *C. calantha* ought to be sympetalous.

***Coccinia helenae* Buscal. & Muschl., Bot. Jahrb. Syst. 59: 498. 1913.**

Type: [Eastern Africa]. At Mbusi river [authors state that this river flows in the Indian Ocean in Mozambique, the collector went upstream towards Zambezi river and Victoria falls; likely the Buzi river is meant], steppe, fl, fr, 3 Dec 1910, *H. von Aosta* [H.L.F.H. d'Orléans] 87 (Holotype: B, destroyed; duplicates ?).

As for *C. aostae*, *C. helenae* seems to be mistaken. Gilg (Engler et al. 1915) suggested that *Coccinia helenae* had been described using the *G.A. Schweinfurth* 932 collection from Blue Nile. A drawing of this species exists in BR (K neg. 4846). The drawing, if it represents a *Coccinia* (the two subsessile female flowers on one node are suspicious), does not match any species of the present author's knowledge from Blue Nile area entirely. It might be *C. abyssinica* if it was collected in the Ethiopian highlands or *C. adoensis* but the fruit would be unusually ovoid. However, if it is from the area as given by Muschler, it might be *C. rehmannii*. As the *von Aosta* 87 specimen is destroyed the name remains dubious. According to the describing authors duplicates of the *von Aosta* specimens have been distributed, and White (1962) found some von Aosta specimens in Florence (FI or FT). However, a loan from FT did not contain any *Coccinia* specimens by von Aosta.

***Coccinia longipetiolata* Chiov., Fl. somala 2: 223. 1932.**

Type: Somalia. [Jubbada Dexhe/Jubbada Hoose border]: between Afmadù [Afmadow] and Saamoggia, 1926, *P. Gorini* 149 (Syntype: FT [K neg. 4851, digital image: JPS]).
Type: Somalia. [Jubbada Dexhe/Jubbada Hoose border]: between Afmadù [Afmadow] and Saamoggia, 1926, *P. Gorini* 150 (Syntype: FT [K neg. 4850, digital image: JPS]).

Remarks. The specimens are quite poor. No leaf is spread out, and generative characters are missing. However, 7-lobate leaves, according to description, do only occur in *C. samburuensis*, which differs in coriaceous leaves and a serrate margin with glandular teeth. Hence, this species name is not synonymous with any *Coccinia* species. The tendrils in *C. longipetiolata* are almost equally bifid, which is not found in *Coccinia*, especially not in species not from rainforests. The drawing accompanying the protologue shows stipules, but this can only be seen in a single node of *P. Gorini* 149, while the other nodes are more typical of Cucurbitaceae. It shows, however, more likely a bud and a probract of similar sizes that give the impression of stipules. In all, the specimens are likely to belong to the Cucurbitaceae.

There are neither characters supporting a relationship with *Coccinia*, nor characters contradicting it, except for the tendrils. Jeffrey (1967) suggests a relationship to his *Coccinia* sp. E sensu F.T.E.A. (*Jarman* 66), but as this specimen could not be examined by the present author, it cannot be discussed. Therefore, that species is treated as dubious. Eventually, sequencing could give disclosure about the relationships.

Invalid names

***Coccinia abdallai* Zimm., nom. nud.**

The name is mentioned on *P.W.A. Zimmermann G6594* (EA!) and in *Die Cucurbitaceen* 2: 8 (1922b) but not described. This is a *Coccinia trilobata*.

***Bryonia acerifolia* D.Dietr., Syn. pl. 5: 367. 1852, nom. nud.**

Dietrich mentions this species in his synopsis as a name by Willdenow. However, no such name by Willdenow is known. As Cogniaux (1881) synonymizes *B. acerifolia* with *B. alceaefolia* Willd., and Dietrich uses the exact same words for *B. acerifolia* as for *B. alceaefolia*, this name is likely just mistaken.

***Bryonia barbata* Buch.-Ham. ex Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 530. 1881, nom. nud.**

If the plate 625 in the East India Company's Museum is additioned by a printed label with description, then this name might be valid, but Cogniaux (1881) mentioned the name as unpublished. He synonymized it with *Coccinia cordifolia* (see also there), so it is not clear whether *B. barbata* is a *C. grandis* or a *Cucumis maderaspatanus*. However, the epithet suggests the existence of rigid trichomes, which does not match with *C. grandis*.

***Coccinia cordifolia* var. *triangularis* A.Chev., nom. nud.**

This name appeared on labels in P specimens of *A.J.B. Chevalier* 8886, 9527, and 10934, and it is apparently unpublished.

***Coccinia crassifolia* H.K.Walter, Naturwissenschaft und Landwirtschaft 9: 33. 1926, nom. nud.**

This is a typographical error of *Caccinia crassifolia* Kuntze, a Boraginaceae.

***Coccinia dubia* Palacký, Lotos 10(4): 70. 1860, nom. nud.**

Palacký cites a *Coccinia dubia*, which was supposed to be described by von Bunge in his "Reliquiae lehmanniae" (von Bunge 1854). However, Palacký mistyped the genus, which is in fact called *Caccinia*, a Boraginaceae.

***Coccinia glandis* nom. nud.**

This is a typographical mistake for *C. grandis* that has been published several times (Tewtrakul et al. 2006, Jiwajinda et al. 2002). This epithet should hence not be used in *Coccinia*.

***Physedra gracilis* A.Chev., Explor. Bot. Afrique occ. franç. 1: 292. 1920, nom. nud.**

Ivory Coast: Bassin de la Moyenne-Sassandra, at Guidéko, *A.J.B. Chevalier 16416* (P! [P05591866, digital image: P]). Ivory Coast: Bassin de la Moyenne-Sassandra, at Guidéko, *A.J.B. Chevalier 19013* (P! [P05591864], P! [P05591865]).

This is a nom. nud. (ICN 32.3) because the note that the plants have yellow flowers cannot be regarded as intended to describe a new species. The specimens are *Coccinia keayana*.

***Cephalandra indica* var. *triangularis* A.Chev., nom. nud.**

This name appeared on a label by Chevalier from October 1908 on *A.J.B. Chevalier 9527*, but apparently was not published.

***Cucumis inedulis* Forssk., Fl. aegypt.-arab.: CXXII. 1755, nom. nud.**

For details, see the Taxonomic remarks of *Coccinia grandis*, to which the name would belong.

***Cephalandra ivorensis* A.Chev., Explor. Bot. Afrique occ. franç. 1: 295. 1920, nom. nud.**

Collections connected to this nomen nudum have been synonymized with *Physedra eglandulosa* (Hook.f.) Hutch. & Dalziel (now *Ruthalicia eglandulosa* (Hook.f.) C.Jeffrey).

***Bryonia lagenaria* E.Mey. ex Drège, Zwei pflanzengeogr. Dokum. 54, 169. 1843, nom. nud.**

This name appears on some Drège collections and is only listed in the work of Meyer. However, the collections are type specimens *Coccinia sessilifolia*.

***Coccinia medica* M.T.H.Khan in: Gottschalk-Batschkus and Green, Handbuch der Ethnotherapien: 377, 384, 517, 537. 2002, nom. nud.**

Khan used this name but without taxonomic context. Most likely, he meant *Coccinia indica*, an illeg. name for *C. grandis*.

***Coccinia monteroi* Hort., Catalogue des graines du Jardin botanique de Bordeaux. 1866.**

The protologue was not available to the present author, but Cogniaux (1881) listed this name as nomen tantum. It is, however, not mentioned in the notes sections of the Bordeaux Garden catalogues from 1866 or 1867 where Naudin described some new species, so it might well be just a nomen nudum.

***Coccinia moshiensis* Zimm., nom. nud.**

Mentioned on *P.W.A. Zimmermann G6599* (EA!). This is a *Coccinia trilobata*.

***Coccinia natalensis* Burt-Davy, A manual of the flowering plants and ferns of the Transvaal with Swaziland, South Africa 1: 237. 1926, nom. nud.**

Cephalandra natalensis Oliv., unknown.

Coccinia natalensis (Oliv.) Cogn., unknown.

The names of Cogniaux und Oliver are mentioned in Burt-Davy and Pott-Leendertz in *Ann. Transvaal Mus.* 3(3): 121. 1912. However, no citation is given. Neither *Cephalandra natalensis* is mentioned in Daniel Oliver's *Flora of Tropical Africa*, nor any Cogniaux publication with this name is known. Burt-Davy writes in 1926 that the name 'appears to have been an unpublished MS. name'.

***Coccinia lalambensis* Schweinf. ex Penzig, Atti Congr. Bot. Int. Genova (1892): 342. 1893, nom. inval.**

Eritrea. [Anseba Province]: Monte Lalambensis near Keren, c. 2000 m, 20 Mar, *G.A. Schweinfurth* 568 (B, destroyed). [Northern Red Sea Province]: Habab, *J.M. Hildebrandt plant. Habab* 1802 (LE?).

No description given, therefore this name is not validly published.

***Cucurbita laevigata* Bl.?, nom. nud.?**

This name is written on a specimen in L herbarium (L0587542). The specimen was part of the collection of C. G. C. Reinwardt but lacks collector, collecting site, and date. One ink-written label solely states "1766.E.5138." and the species name. Another label, written with a pencil, says "*Cucurbita laevigata*" "mihi" and "Callelet W[...]". The last word is unreadable to the present author. Another specimen (L0587515) bears a similar label with "1766.E.5138.", however without a pencil-written label. Since both specimens are *Coccinia grandis*, *Cucurbita laevigata* would be a synonym, if it had been validly published. A Waitz collection (L0587563) bears the names "*Cucurbita laevigata* Bl." and "Callelet Bl.", so the former name is maybe a Blume manuscript name and the latter one is indigenous.

***Bryonia quinquefolia* Noronha, Verh. Bat. Genootsch. 5: 155(8). 1790, nom. nud.** Miquel (1855) synonymizes this nomen nudum under *C. wightiana* M.Roem., which is *C. grandis* (see Taxonomic remarks there). However, he also synonymizes *Bryonopsis pedata* Hassk., which cites *Bryonia quinquefolia*. The description of *Bryonopsis pedata* mentions lacinate, almost pinnatifid leaves and male flowers in oblong clustered racemes. This does not match *C. grandis* at all but eventually *Diplocyclos palmatus*, which Roemer excluded from *C. indica* as *Coccinia palmata* M.Roem. The identity of *Bryonia quinquefolia* might be solved, if one finds a Noroña specimen stating “*Bryonia quinqueloba*” or “Oyot-kekèp”, the latter name being the Javanese term for this species. However, the epithet rather links to the deeply lobate leaves of *D. palmatus*.

***Bryonopsis pedata* Hassk., Cat. hort. bot. bogor.: 189. 1844, nom. nud.**

Hasskarl cites Noroña’s *Bryonia quinquefolia* and a vernacular name “aroy kalanyar beurriet”. The given description of lacinate, almost pinnatifid leaves and male flowers in oblong clustered racemes does not match *C. grandis*. According to Filet (1859), the vernacular name is used in Sundanese and refers to *Bryonia* [*sic*] *pedata* Hassk., two *Trichosanthes* species and *Luffa cordifolia* Bl. None of these names have been referred to *Coccinia*, so it seems unlikely that *Bryonopsis pedata* does.

***Coccinia peterii* Zimm., nom. nud.**

This is an unpublished ms. name on *R. Soleman* 6046 (EA!). The specimen, however, is a *C. grandis*.

***Bryonia ruderalis* Zipp. ex Span., Linnaea 15: 206. 1841, nom. illeg. & nom. nud.**

In L herbarium, there is a specimen determined as “*Bryonia ruderalis* Zp.” from Timor (L0587573), which is a Zippelius collection of *Coccinia grandis*. However, the name is a later homonym of *Bryonia ruderalis* Salisb. Additionally, it lacks a description in the publication, so it is a nomen nudum, too.

***Cucurbita schimperiana* Hochst., nom. nud.**

The name was used on printed labels of *G.H.W. Schimper* 1570 (effective publication), which lack a proper description (hence a nom. nud.). Under this distribution number, specimens from two different shipments are included. One is taken from package “P. 16 K. no. 4”, collected on 23 Apr 1841 in Djeladjeranné (label on P specimen). The data of this label were used for C. F. F. Hochstetter’s printed labels. The TUB-004724 and TUB-004725 specimens bear a Schimper label from package “P. 10 D. no. 23” from “Landschaft Modat” collected in April 1839. An unnumbered W specimen also

notes this collecting site, hence the specimen might be from the same shipment. Specimens of both collections are *Coccinia grandis*.

***Coccinia schultzei* Gilg, Namaland & Kalahari: 697. 1907, nom. nud.**

Apparently a collection by L. Schultze (*Schultze 320a*) in B herbarium, but not validly published by Gilg afterwards. However, if so, then the holotype was burned in the Berlin herbarium fire in 1943.

***Coccinia sericea* Zimm., nom. nud.**

Zimmermann marked the specimen *P.W.A. Zimmermann G6600* (EA!) to be a new species, but B. Verdcourt pointed out on the specimen that Zimmermann never published it. In any case, this specimen belongs to *C. grandis*.

***Bryonia sinuosa* Wall., Numer. List 6716. 1832, nom. nud.**

The present author did not see a specimen with this number, so it cannot be decided whether Cogniaux' (1881) partial synonymization of *Coccinia cordifolia* refers to *Coccinia grandis* or *Cucumis maderaspatanus*. Wallich himself supposed that this collection is a mix of (*Bryonia*) *Coccinia grandis* and *Melothria indica*.

***Cephalandra sylvatica* A.Chev., Explor. Bot. Afrique occ. franç. 1: 295. 1920, nom. nud.**

Collections connected to this nomen nudum (ICN 32.3) have been synonymized with *Physedra eglandulosa* (Hook.f.) Hutch. & Dalziel (now *Ruthalicia eglandulosa* (Hook.f.) C.Jeffrey).

***Cucurbita triangulata* Hochst. ex Cogn. in A.DC. & C.DC. Monogr. Phan. 3: 532. 1881, nom. nud.**

Cogniaux cites a Schimper specimen (*Iter Abyss. Sect. 3 no. 1202*) that was supposed to be labeled by C. F. F. Hochstetter. There are several sheets with this distribution number in Paris, but only one bears this name. The location is given by "In Semen" [Semien Mts]. The other Paris specimens with this number are from Baria Dikeno (collected on 6 Aug 1853). The collection is a *Coccinia grandis*.

***Coccinia wightii* Miq., Fl. Ned. Ind. 1(1): 1112. 1855, nom. nud.**

Name variation of *Coccinia wightiana* M.Roem. in the index of the book.

Excluded taxa

***Physedra bequaertii* De Wild., Pl. Bequart. 1: 569. 1922.**

Type: D.R. Congo. Along the Semliki [river], female, fl, 16 Jun 1914, *J. Bequaert* 4791 (Syntypes: BR [BR0000008886330, digital image: BR, JPS], BR [BR0000008887122, digital image: BR, JPS], BR [BR0000008887245, digital image: BR, JPS]).

Type: D.R. Congo. Along Ruthuru river, female, fl, 17 Nov 1914, *J. Bequaert* 6315 [*sic*, should be 6215] (Syntypes: BR [BR0000008886477, digital image: BR, JPS], BR [BR0000008886521, digital image: BR, JPS]).

Remarks. As *Physedra* is a synonym of *Coccinia* but the specimens are belonging to the genus *Bambekea*, the name *P. bequaertii* has to be excluded. Jeffrey (1962: 364) published the new combination *Bambekea bequaertii* (De Wild.) C. Jeffrey, however, it is not clear whether this is a species separate from *Bambekea racemosa* Cogn.

***Coccinia cordifolia* (L.) Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 529. 1881. pro parte.**

Bryonia foliis cordatis oblongis quinquangularibus dentatis scabris L., Fl. zeyl.: 168. 1747.
Bryonia cordifolia L., Sp. pl. 2: 1012. 1763.

Type: Sri Lanka. No detailed location given, *P. Hermann* 354 (Type: Herm. Flora zeylanica 2:22, BM! [BM000521582, digital image: BM]).

Remarks. Cogniaux cites *Bryonia cordifolia* sensu Linnaeus' Species plantarum 2nd edition, where Linnaeus synonymizes Rumphius' *Vitis alba indica*, which is *Coccinia grandis*. If *B. cordifolia* would have been originally described in 1763, Cogniaux' choice would have been valid, but *B. cordifolia* was described in 1753 (Species plantarum 1st edition), where *Vitis alba indica* is not mentioned, but only a collection from Hermann herbarium, which is *Cucumis maderaspatanus* L.

***Coccinia decipiens* (Hook.f. in Oliv.) Cogn. in A.DC. & C.DC., Monogr. Phan. 3: 539. 1881.**

Cephalandra decipiens Hook.f. in Oliv., F. T. A. 2: 552. 1871.

Type: Angola. [Cuanza Norte]: Pungo Adongo, grassland, *FM.J. Welwitsch* 816 (Holotype: BM! [BM000799218], isotypes: COI [COI00005507, digital image: JPS], K [K000313451, digital image: JPS, K], LISU [LISU00214555, digital image: JPS]).

Remarks. The sessile beaked fruits match to the genus *Diplocyclos*, which has been correctly observed by Jeffrey (1962) as *Diplocyclos decipiens* (Hook.f.) C. Jeffrey.

***Coccinia ecirrhosa* Cogn., Bull. Herb. Boissier 4(12): 822. 1896.**

Type: Somalia. Abdallah, 1891, *C. Keller* 106 [sphalm. 116 in l.c.] (Type: BR! [BR0000008885999, digital image: BR, JPS], Z! [Z-000004442, digital image: Z, K neg. 4832]).

Remarks. The type does not contain much material, but the lower surface of a leaf shows a pinnatifid venation pattern, which is unknown in *Coccinia*. Jeffrey (1967) transferred it correctly (Kocyan et al. 2007) to another genus, namely *Cephalopentandra*.

***Coccinia gabonensis* Keraudren, Adansonia 8: 40. 1968.**

Type: Gabon. Ogooué-Ivindo: Bélinga, 950–1000 m, male and female, fl, Nov 1964, *N. Hallé* 3018 (Holotype: P [P00348266, digital image: JPS, P], isotypes: P! [P00348264, digital image: JPS, P], P! [P00348265, digital image: JPS, P], K! [K000313237, digital image: JPS, K]).

Remarks. This species does certainly not belong to *Coccinia*. It is monoecious, has free petals, and rather large bracts, in contrast to the dioecious, sympetalous *Coccinia* species with much smaller bracts. It rather belongs to *Momordica*.

***Coccinia macrantha* nom. nud.**

Physedra macrantha Gilg, Bot. Jahrb. Syst. 34: 356. 1904.

Type: Liberia. Gran Bassa: Fishtown, in bush, sand, fl, 10 m, 27 Aug 1898, *M. Dinklage* 2019 [Cucurbitaceae no. 1846] (Holotype: B! [B 10 0154925, digital image: B, JPS]).

Remarks. The combination *Coccinia macrantha* was only used in B herbarium for storage but apparently never published. *Physedra macrantha* Gilg has been synonymized, correctly, with *Physedra eglandulosa* Hutch. & Dalziel (1928), which is now in the genus *Ruthalicia*.

***Coccinia obbadiensis* (Chiov.) Cufod., Supplem.: Enum. Pl. Aethiop. Spermatophyta 35(2): 1050. 1965.**

Cephalopentandra obbadiensis Chiov., Fl. somala: 187, tab. 20, fig. 1. 1929.

Type: Somalia. [Mudug]: Obbia [Hobyó] Sultanate, between Dolobscío and Magghiòle, 27 Apr, *G. Stefanini* & *N. Puccioni* 458 (Holotype: FT [FT003569, K neg. 4845, digital image: JPS], isotype: K! [K000313183, digital image: JPS, K]).

Remarks. The author notes five stamens, which are a good reason for not including this species in *Coccinia*. The leaves of the specimen on the picture look like these of *Coccinia quercifolia*, which is also excluded from *Coccinia* and separated by Jeffrey (1967).

***Coccinia quercifolia* Hutch. et E.A.Bruce, Bull. Misc. Inform. Kew 2: 99. 1941.**

Type: Somalia. [Nugaal]: Boundary Pillar 93, 45°9'E, 8°37'N, 990 m, fl, 6 Oct, *J.B. Gillett 4194* (Holotype: K! [K000313174, digital image: JPS, K], K! [K000313174, digital image: JPS, K]).

Type: Ethiopia. [Somali Region]: Harradigit [c. 7°45'N, 45°30'E], Apr, *F.L. James & J.G. Thrupp no.?* (Paratype: K?).

Remarks. The leaves have a pinnatifid venation, just like *Cephalopentandra* (*Coccinia*) *ecirrhosa*, but are deeply lobed. Since the leaf form may vary in Cucurbitaceae, it is likely that these specimens belong together, so that the odd leaf venation excludes this type from *Coccinia*. Jeffrey already synonymized this name with *C. ecirrhosa*.

***Coccinia palmata* (L.) M.Roem., Syn. pepon.: 93. 1846.**

Bryonia palmata L., Sp. pl. 2: 1012. 1753. *Bryonia zeylanica, folio quinquepartito* Burm., Thes. zeylan.: 49. 1737. *Bryonia foliis palmatis levibus quinquepartitis: laciniis lanceolatis repando-serratis* L., Fl. zeyl.: 146. 1747.

Type: [Sri Lanka]. No location given. *P. Hermann 25* (Type lost?).

Type: [Sri Lanka]. No location given. *P. Hermann 41* (Type lost?).

Type: [Sri Lanka]. No location given. *P. Hermann 62* (Type lost?).

Type: [Sri Lanka]. No location given. *P. Hermann 353* [Musæum zeylonicum 2:58] (Lectotype, designated by Jeffrey (1962: 352): BM [BM-000621700]).

Remarks. The name *Coccinia palmata* has been applied illegitimately for *C. mack-enii* for a long time due to an overlooked combination. When Wight and Arnott published the name *Coccinia indica*, they cited *Bryonia grandis* L. and also tentatively included the citation of *Bryonia palmata* L. More likely, however, they meant a specimen in Herbarium Madras that was identified as *B. palmata* L. One year after Voigt's correction to *Coccinia grandis* (L.) Voigt, Roemer (1846) also recognized the seemingly missing combination and that Linnaeus' *B. palmata* and *B. grandis* indeed referred to different species. Roemer treated them, amongst other species, as *C. grandis* (L.) M.Roem. (nom. illeg.) and *C. palmata* (L.) M.Roem. Roemer cited the second edition of Linnaeus' *Species plantarum* (1763), which has the identical description for this species as the first edition, so he explicitly meant *Bryonia palmata* to be a part of *Coccinia*. The *Bryonia palmata* typelectotype belongs to what is now widely called *Diplocyclos*, due to the globose striped fruits

easily recognizable as being not part of *Coccinia*. Apart from that, another species from South Africa was described with the name *Cephalandra palmata* E.Mey. ex Sond. (Harvey and Sonder 1862). Cogniaux (1881) accepted this species in *Coccinia* but overlooked *Coccinia palmata* (L.) M.Roem. He thus created an illegitimate *Coccinia palmata* (E.Mey. ex Sond.) Cogn., which has since been used for this species. Holstein and Renner (2010) called attention to this erroneous usage by resurrecting the correct name, *Coccinia mackenii* Naudin ex C.Huber, while *C. palmata* is a synonym of *Diplocyclos palmatus* in spite of its wide usage for *C. mackenii* since then.

***Coccinia petersii* Gilg, Bot. Jahrb. Syst. 34: 356. 1904.**

Type: Mozambique. [Zambézia]: Rios de Sena and Boror, without detailed locality, male and female, on dry ground, 1846, *W.H.C. Peters s.n.* (Holotype: B destroyed, isotype: K).

Remarks. Jeffrey (1967) synonymized this species with *Eureiandra fasciculata* (Cogn.) C.Jeffrey.

***Coccinia polyantha* Gilg, Bot. Jahrb. Syst. 34: 356. 1904.**

Type: Tanzania. Lindi: Kilwa district, near Mariwe, upon low shrub, in light, slightly wet ground pori, fl, Dec, *W. Busse 512* (Syntype: B destroyed, isosyntype: EA [EA000002139, digital image: JPS]).

Type: Tanzania. [South central Tanzania], Kissaki steppe at Rufiji river, 250 m, on sandy laterite soil, fr, Nov, *Götze 80* (Syntype: B destroyed).

Remarks. The seeds are described as globose to subglobose, but *Coccinia* seeds are rather flat. It is therefore unlikely that this species belongs to *Coccinia*. Jeffrey (1967) synonymized this species with *Eureiandra fasciculata* (Cogn.) C.Jeffrey.

***Cucumis rheedii* Kostel., Allg. med.-pharm. Fl. 2: 738. 1833.**

Schunambuvalli Rheede, Hort. malab. 8: 21, t. 11. 1688.

Type: drawing in l.c.

Remarks. Cogniaux and Harms (1924) synonymized this species with *Coccinia cordifolia* (L.) Cogn., a name that Cogniaux used for *C. grandis* (L.) Voigt. However, the drawing shows 3–5-fid tendrils and small? subglobose fruits on an ebracteate panicle. This does not correspond to *Coccinia*.

***Coccinia stefaninii* Chiov., Result. Sc. Miss. Stefan.-Paoli Somal. Ital. 1: 84. 1916.**

Type: Somalia. Somali Republic, Jubaland, Dintorni di El Uré, *G. Paoli 1069* (Syntype: FT! [2 sheets, FT003512, digital image: JPS]).

Type: Somalia. Fra Jagdoudou e Duddumai, *G. Paoli 1179* (Syntype: FT! [FT003513, digital image: JPS]).

Remarks. This name has been transferred to the genus *Dactyliandra* by Jeffrey (1985).

***Coccinia stolzii* Harms, Bot. Jahrb. Syst. 57: 241. 1923.**

Type: Nyasaland [Tanzania]. Kyimbila district, Mbaku, 600 m, fl, fr, Jun 1913, *A.F. Stolz 2028* (Holotype: ?, isotype: BM!, G! [G00301602], K [K000313456, digital image: JPS, K], L!, P! [P05590096, digital image: P], PR!, PRE [PRE0592846-0, digital image: JPS], S [S08-12469, digital image: JPS], U! [U0074536], W!, Z! [Z-000004440, digital image: Z]).

Remarks. Monoecious plant with several small subglobose fruits per node apply clearly to *Diplocyclos* and must therefore be synonymized as it has been done by Jeffrey (1967) to *Diplocyclos decipiens*.

***Coccinia suburceolata* Cogn., Symb. Antill. (Urban) 1: 450. 1900.**

Type: Haiti. Near Port-au-Prince, in Tête bois de pin, 1800 m, male, fl, Nov, *L. Picarda 1498* (Holotype: BR! [BR0000009939141, digital image: JPS], isotypes: B, L, NY, S?).

Remarks. This plant was found on Haiti and is therefore geographically far away from the natural distribution range of the genus *Coccinia*. According to the protologue, the tendrils are often trifid and the fruit is spherical and apple-sized, which does not fit to the morphospace of any *Coccinia* species. Urban (1921) put the specimen and thus species, amongst others, into a new genus: *Penelopeia*, which has been confirmed by Kocyan et al. (2007).

Acknowledgements

I thank the curators of B, BAR, BM, BR, BRI, C, CANB, CBG, COI, DSM, EA, FR, FT, G, GAT, GOET, H, HBG, HEID, JE, K, L, LISC, LISU, M, MO, MSB, NHT, P, PERTH, PR, PRC, S, U, UBT, UPS, W, WAG, Z, ZT, and Charlie Jarvis for making the collections available, sending loans or scans, or helpful comments about the collections. I am grateful to Jean-Luc Gatard for sending seeds and plants, without which

most of the morphological findings and the crossing experiments would not have been possible. I also thank Frank M. Mbago for his help and excellent eye on the field trip in Tanzania.

Thanks are given to Hanno Schäfer for communicating various observations and specimens from DNA, NY, US, Christian Bräuchler for communicating and discussion of Schimper specimens from TUB, and Eberhard Fischer for communicating material from Rwanda. Shixiao Luo is thanked for help with translations from Chinese language, Werner Greuter for clarifying doubts with the validity of *Cucurbita exanthematica*, and Brigitta and Willem de Wilde for discussion of the *Momordica bicolor* specimens and sharing their field experience with *C. grandis*. I thank Eva Facher for help with staining of the petiole sections and Mila Vovsky for preparing the chromosomes. Thanks are given particularly to Marc Gottschling for many fruitful discussions and advice on the ms., esp. in taxonomic questions and Susanne S. Renner for supervision and English revision. I also thank Sandy Knapp and two anonymous reviewers for many helpful comments on the manuscript.

This work has been financed by Deutsche Forschungsgemeinschaft (RE603/6-1 and 6-2 given to S.S. Renner). Travel to the herbaria, BR, P, and W have been funded by travel grants by Munich Graduate School for Evolution, Ecology and Systematics of the LMU Munich. This research received support (visit to L and U collections) from the SYNTHESYS Project (<http://www.synthesys.info/>), which is financed by European Community Research Infrastructure Action under the FP7 “Capacities” Program. I also like to acknowledge the Tanzania Commission for Science and Technology (COSTECH) for providing research clearance to undertake fieldwork in Tanzania.

References

- Abera G, Guteta H (2007) Response of anchote (*Coccinia abyssinica*) to organic and inorganic fertilizers rates and plant population density in western Oromia, Ethiopia. *East African Journal of Sciences* 1: 120–126. doi: 10.4314/eajsci.v1i2.40350
- Adanson M (1757) *Histoire naturelle du Sénégal*. C.-J.-B. Bauche, Paris, France, 159 pp.
- Addis G, Baskaran R, Raju M, Ushadevi A, Asfaw Z, Woldu Z, Baskaran V (2009) Effect of blanching and drying process on carotenoids composition of underutilized Ethiopian (*Coccinia grandis* L. Voigt) and Indian (*Trigonella foenum-graecum* L.) green leafy vegetables. *Journal of Food Processing and Preservation* 33: 744–762. doi: 10.1111/j.1745-4549.2008.00308.x
- Aga HF, Badada KU (1997) Nutritional and antinutritional characteristics of anchote (*Coccinia abyssinica*). *Ethiopian Journal of Health Development* 11: 163–168.
- Agarwal PK, Roy RP (1975) Colchicine induced hexaploid in *Coccinia indica* W. & A. *The Nucleus* 18: 122–124.
- Agarwal PK, Roy RP (1983) X-ray irradiation studies in *Coccinia indica* W. et A. *Cytologia* 48: 691–697. doi: 10.1508/cytologia.48.691
- Agarwal PK, Roy RP (1984) Karyotype of *Coccinia indica*. *Indian Journal of Genetics & Plant Breeding* 44: 117–120.

- Agarwal VM, Rastogi N (2008) Deterrent effect of a guild of extrafloral nectary-visiting ant species on *Raphidopalpa foveicollis*, a major insect pest of sponge gourd, *Luffa cylindrica*. *Entomologia Experimentalis et Applicata* 128: 303–311. doi: 10.1111/j.1570-7458.2008.00727.x
- André E (1900) *Coccinia dinteri*. *Revue horticole*, 268–269.
- Aryal KP (2007) Uncultivated plants: an option for livelihood support of the people in mid-hills of Nepal. Uppsala Universitet, Uppsala, Sweden.
- Asfaw Z (1997) Conservation and use of traditional vegetables in Ethiopia In: Guarino L (Ed.) *Traditional african vegetables: Proceedings of the IPGRI International Workshop on genetic resources of traditional vegetables in Africa Conservation and use*. ICRAF-HQ, Nairobi, Kenya, 57–65.
- Asfaw Z, Tadesse M (2001) Prospects for sustainable use and development of wild food plants in Ethiopia. *Economic Botany* 55: 47–62. doi: 10.1007/BF02864545
- Avetta G (1894) Sui cistoliti delle foglie de alcune *Coccinia*. *Annuario del Reale Istituto botanico di Roma* 5: 181–185.
- Awasthi P (1962) Palynological investigations of Cucurbitaceae part II. *Proceedings of the National Institute of Science in India - B* 28: 485–496.
- Azad Khan AK, Akhtar S, Mahtab H (1980) Treatment of diabetes mellitus with *Coccinia indica*. *British Medical Journal* 280: 1044. doi: 10.1136/bmj.280.6220.1044
- Baker HG, Cox PA (1984) Further thoughts on dioecism and islands. *Annals of the Missouri Botanical Garden* 71: 244–253. doi: 10.2307/2399068
- Balinsky BI (1962) Patterns of animal distribution on the African continent. *Annals of the Cape Provincial Museums* 2: 299–310.
- Bamba JP, Miller RH, Reddy GVP, Muniappan RN (2009) Studies on the biology, host specificity, and feeding behavior of *Acythopeus cocciniae* O'Brien and Pakaluk (Coleoptera: Curculionidae) on *Coccinia grandis* (L.) Voigt (Cucurbitaceae) and *Zehneria guamensis* (Merrill) Fosberg (Cucurbitaceae). *Micronesica* 41: 71–82.
- Barua AB, Goswami BC (1979) Carotenoids of *Cephalandra indica* (*Coccinia indica*). *Current Science* 48: 630–632.
- Bekele A, Feyissa T, Tesfaye K (2014) Genetic diversity of anchote (*Coccinia abyssinica* (Lam.) Cogn.) from Ethiopia as revealed by ISSR markers. *Genetic Resources and Crop Evolution* 61: 707–719. doi: 10.1007/s10722-014-0090-9
- Bentham G, Hooker JD (1867) *Genera Plantarum*. Reeve & Co.; Williams & Norgate, London, UK, XV, 319 pp.
- Berndt R (2007) A global survey of *Puccinia*-rust on Cucurbitaceae. *Mycological Progress* 6: 151–178. doi: 10.1007/s11557-007-0535-z
- Bhaduri P, Bose P (1947) Cyto-genetical investigations in some common cucurbits, with special reference to fragmentation of chromosomes as a physical basis of speciation. *Journal of Genetics* 48: 237–256. doi: 10.1007/BF02989384
- Bhagavan Reddy P, Reddy SM (1987) Physiology of host-pathogen interaction in the fruit rot of *Coccinia* (*Coccinia indica*) caused by *Colletotrichum gloeosporioides*. *Indian Journal of Mycology and Plant Pathology* 17: 289–294.
- Bhagavan Reddy P, Reddy SN (1988) Role of oxidases and cell degrading enzymes in the fruit rot of coccinia caused by *Rhizoctonia solani*. *Indian Phytopathology* 41: 138–141.

- Bhakuni DS, Srivastava SN, Sharma VN, Kaul KN (1962) Chemical examination of the fruits of *Coccinia indica*. Journal of Scientific & Industrial Research Section B 21B: 237–238.
- Bharathi LK (2007) Ivy gourd. In: Peter KV (Ed.) Underutilized and underexploited horticultural crops. New India Publishing, New Delhi, India, 275–282.
- Bhatia SK, Mahto Y (1968) Note on breeding of fruit flies *Dacus ciliatus* Loew and *D. cucurbitae* Coquillett in stem galls of *Coccinia indica* W. & A. Indian Journal of Entomology 30: 244–245.
- Bhatt D, Kumar A (2001) Foraging ecology of red-vented bulbul *Pycnonotus cafer* in Haridwar, India. Forktail 17: 109–110.
- Bhave MR, Gupta VS, Ranjekar PK (1986) Arrangement and size distribution of repeat and single copy DNA sequences in four species of Cucurbitaceae. Plant Systematics and Ecology 152: 133–151. doi: 10.1007/BF00989423
- Bhave MR, Lagu M, Ranjekar PK (1984) Molecular analysis of Cucurbitaceae genomes: I. Comparison of DNA reassociation kinetics in six plant species. Plant Science Letters 33: 127–136. doi: 10.1016/0304-4211(84)90002-6
- Bhuskute SM, Makde KH, Deshpande PK (1986) Staminal organization in *Coccinia grandis* (L.) Voigt. Annals of Botany 57: 415–418.
- Bleher B, Potgieter CJ, Johnson DN, Böhning-Gaese K (2003) The importance of figs for frugivores in a South African coastal forest. Journal of Tropical Ecology 19: 375–386. doi: 10.1017/S0266467403003420
- Bosch CH (2004) *Coccinia sessilifolia* (Sond.) Cogn. In: Grubben GJH, Denton OA (Eds) Vegetables/Légumes PROTA (Plant Resources of Tropical Africa / Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands, 230–231.
- Bradstock RA, Auld TD (1995) Soil temperatures during experimental bushfires in relation to fire intensity: consequences for legume germination and fire management in south-eastern Australia. Journal of Applied Ecology 32: 76–84. doi: 10.2307/2404417
- Bulbul IJ, Nahar L, Haque M (2011) Antibacterial, cytotoxic and antioxidant activity of chloroform, n-hexane and ethyl acetate extract of plant *Coccinia cordifolia*. Agriculture and Biology Journal of North America 2: 713–719. doi: 10.5251/abjna.2011.2.4.713.719
- Chakravarty HL (1948) Extrafloral glands of Cucurbitaceae. Nature 162: 576–577. doi: 10.1038/162576b0
- Chakravarty HL (1954) Morphology of the staminate flowers in the Cucurbitaceae with special reference to the evolution of the stamens. Bulletin of the Botanical Society of Bengal 8: 186–215.
- Chakravorti AK (1947) The development of female gametophyte and seed of *Coccinia indica* W. & A. Journal of the Indian Botanical Society 26: 95–104.
- Chakravorti AK (1948) Cytology of *Coccinia indica* W. & A. with reference to the behaviour of its sex-chromosomes. Proceedings of the Indian Academy of Sciences - Section B 27: 74–86.
- Chatage VS, Bhale UN (2010) Fungi associated with fruit rot diseases of ivy gourd (*Coccinia indica* Wight & Arn.) from Maharashtra, India. Bioinfolet 7: 242–243.
- Chittchang U, Jittinandana S, Sungpuag P, Chavasit V, Wasantwisut E (1999) Recommending vitamin A-rich foods in southern Thailand. Food and Nutrition Bulletin 20: 238–243.
- Chopra RN (1955) Some observations on endosperm development in the Cucurbitaceae. Phytomorphology 5: 219–230.

- Chopra RN, Bose JP (1925) *Cephalandra indica* (telakucha) in diabetes. Indian Journal of Medical Research 13: 11–16.
- Chun ME (2001) Biology and host specificity of *Melittia oedipus* (Lepidoptera: Sesiidae), a biological control agent of *Coccinia grandis* (Cucurbitaceae). Proceedings of the Hawaiian Entomological Society 35: 85–93.
- Chun ME (2002) Biological control of ivy gourd, *Coccinia grandis* (Cucurbitaceae), in Hawai'i. In: Smith CW, Denslow J, Hight S (Eds) Workshop on biological control of native ecosystems in Hawaii. Pacific Cooperative Studies Unit, University of Hawaii at Manoa, Department of Botany, 8–12.
- Cogniaux A (1878) Remarques sur les Cucurbitacées brésiliennes, et particulièrement sur leur dispersion géographique. Bulletin de la Société Royale de Botanique de Belgique 17: 273–303.
- Cogniaux A (1881) Cucurbitacées. In: de Candolle AP, de Candolle C (Eds) Monographiae Phanerogamarum. G. Masson, Paris, France, 340–954. doi: 10.2307/3666537
- Cogniaux A (1916) Cucurbitaceae congolanae novae: herbarii bruxellensis. Bulletin du Jardin Botanique de l'État à Bruxelles 5: 109–116.
- Cogniaux A, Harms HAT (1924) Cucurbitaceae-Cucurbiteae-Cucumerinae. In: Engler A (Ed.) Das Pflanzenreich. Verlag von Wilhelm Engelmann, Leipzig, Germany, 246 pp.
- Das VM, Sultana S (1979) Five new species of the genus *Pratylenchus* from vegetable crops of Hyderabad (Andhra Pradesh). Indian Journal of Nematology 9: 5–14.
- Datta SK (1988) Studies in *Cephalandra indica* Naud. Cytologia 53: 731–737. doi: 10.1508/cytologia.53.731
- Davis RI, Ruabete TK (2010) Records of plant pathogenic viruses and virus-like agents from 22 Pacific island countries and territories: a review and an update. Australasian Plant Pathology 39: 265–291. doi: 10.1071/AP10047
- de Boer HJ, Kool A, Broberg A, Mziray WR, Hedberg I, Levenfors JJ (2005) Anti-fungal and anti-bacterial activity of some herbal remedies from Tanzania. Journal of Ethnopharmacology 96: 461–469. doi: 10.1016/j.jep.2004.09.035
- de Wilde WJJO, Duyfjes BEE (2008) Cucurbitaceae. In: Santisuk T, Larsen K, Nielsen I, Chayamarit K (Eds) Flora of Thailand. Forest Herbarium, Royal Forest Department, Bangkok, Thailand, 136 pp.
- de Wilde WJJO, Duyfjes BEE, Phonsena P, van der Ham RWJM (2011) Miscellaneous South East Asian cucurbit news IV. Thai Forest Bulletin 39: 1–22.
- de Winter B (1971) Floristic relationships between the northern and southern arid areas in Africa. Mitteilungen der Botanischen Staatssammlung München 10: 424–437.
- Deshpande PK, Bhuskute SM, Makde KH (1986) Microsporogenesis and male gametophyte in some Cucurbitaceae. Phytomorphology 36: 145–150.
- Dewanjee S, Kundu M, Maiti A, Majumdar R, Majumdar A, Mandal SC (2007a) *In vitro* evaluation of antimicrobial activity of crude extract from plants *Diospyros peregrina*, *Coccinia grandis* and *Swietenia macrophylla*. Tropical Journal of Pharmaceutical Research 6: 773–778. doi: 10.4314/tjpr.v6i3.14658
- Dewanjee S, Maiti A, Kundu M, Mandal SC (2007b) Evaluation of anthelmintic activity of crude extracts of *Diospyros peregrina*, *Coccinia grandis* and *Schima wallichii*. Dhaka University Journal of Pharmaceutical Science 6: 121–123.

- Dharmamaraju E (1968) A note on the occurrence of fruit fly maggots of *Dacus cucurbitae* Coq. in Cecidomyiid galls of *Coccinia indica* in Andhra Pradesh. Indian Journal of Entomology 30: 87.
- Dinter MK (1912) Die vegetabilische Veldkost Deutsch-Südwest-Afrikas. Selbstverlag, Okahandja, Namibia, 47 pp.
- Dinter MK (1919/20) Index der aus Deutsch-Südwest-Afrika bis zum Jahre 1917 bekannt gewordenen Pflanzenarten IV. Feddes Repertorium 26: 167–174. doi: 10.1002/fedr.19190160507
- Dunnill PM, Fowden L (1965) The amino acids of seeds of the Cucurbitaceae. Phytochemistry 4: 933–944. doi: 10.1016/S0031-9422(00)86271-8
- Dutt MN (Ed.) (1891–1894) The Ramayana. translated into English prose from the original Sanskrit of Valmiki. Manmatha Nath Dutt, Calcutta, India, 1933 pp.
- Edwards S (1815) *Bryonia quinqueloba*. The botanical register 1: 82.
- Elangovan V, Marimuthu G, Kunz TH (2001) Temporal patterns of resource use by the short-nosed fruit bat, *Cynopterus sphinx* (Megachiroptera: Pteropodidae). Journal of Mammalogy 82: 161–165. doi: 10.1644/1545-1542(2001)082<0161:TPORUB>2.0.CO;2
- Engle LM, Altoveros NC, Geal GG, Phanmaliving K, Xaygnalis P, Opena RT (1998) Domestication of *Coccinia grandis*: an underutilized vegetable species in Lao PDR. Philippine Journal of Crop Science 23 (Supplement no. 1): 73.
- Engler A, Diels FLE, Gilg EF, Graebner P, Harms HAT, Lindau G, Loesener T, Schlechter R, Volkens GLA (1915) Berichtigungen zu den von R. Muschler in Engl. Bot. Jahrb. XLIII. (1909), XLVI. (1911), XLIX. (1913) und L. Suppl. (1914) veröffentlichten Diagnosen afrikanischer Pflanzen. Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 53: 366–375.
- Enslin PR, Joubert FJ, Rehm S (1956) Bitter principles of the Cucurbitaceae. III.—elaterase, an active enzyme for the hydrolysis of bitter principle glycosides. Journal of the Science of Food and Agriculture 7: 646–655. doi: 10.1002/jsfa.2740071004
- Eshrat MH (2003) Effect of *Coccinia indica* (L.) and *Abroma augusta* (L.) on glycemia, lipid profile and on indicators of end-organ damage in streptozotocin induced diabetic rats. Indian Journal of Clinical Biochemistry 18: 54–63. doi: 10.1007/bf02867368
- Farrukh U, Shareef H, Mahmud S, Ali SA, Rizwani GH (2008) Antibacterial activities of *Coccinia grandis* L. Pakistan Journal of Botany 40: 1259–1262.
- Filet GJ (1859) De inlandsche plantennamen bijeenverzameld en in alphabetische orde gerangschikt. Natuurkundig tijdschrift voor Nederlandsch Indië 19: 1–280.
- Forsskål P (1775) Flora Ægyptiaco-Arabica sive descriptiones plantarum quas per Ægyptum Inferiorem et Arabiam felicem detexit. Möller, Hauniæ [Copenhagen], Denmark, CXXVI, 220 pp.
- Friis I (1983) Notes on the botanical collections and publications of Pehr Forsskal. Kew Bulletin 38: 457–467. doi: 10.2307/4107837
- Friis I (1984) An analysis of the protologues of new genera with new species in Forsskål's Flora Ægyptiaco-Arabica. Taxon 33: 655–667. doi: 10.2307/1220782
- Gelmesa D (2010) Shifting to alternative food source: potential to overcome Ethiopias' malnutrition and poverty problems. In: Coudel E, Devautour H, Soulard C, Hubert B (Eds) ISDA 2010, Montpellier, France, 10 pp.

- Getahun A (1973) Developmental anatomy and germination of seeds of anchoté, *Coccinia abyssinica* (W. & A.) Cogn. (Cucurbitaceae). *Botaniska Notiser* 126: 437–449.
- Getahun A (1974a) Developmental anatomy of seedlings of anchoté, *Coccinia abyssinica* (W. & A.) Cogn. (Cucurbitaceae). *Botaniska Notiser* 127: 211–223.
- Getahun A (1974b) The role of wild plants in the native diet in Ethiopia. *Agro-Ecosystems* 1: 45–56. doi: 10.1016/0304-3746(74)90006-7
- Gignoux J, Clobert J, Menaut J-C (1997) Alternative fire resistance strategies in savanna trees. *Oecologia* 110: 576–583. doi: 10.1007/s004420050198
- Gillett JB (1972) W. G. Schimper's botanical collecting localities in Ethiopia. *Kew Bulletin* 27: 115–128. doi: 10.2307/4117875
- Gmelin JF (1791) *Systema naturae*. Georg Emanuel Beer, Leipzig, Germany, 777 pp.
- Gradé JT, Tabuti JRS, Van Damme P (2009) Ethnoveterinary knowledge in pastoral Karamoja, Uganda. *Journal of Ethnopharmacology* 122: 273–293. doi: 10.1016/j.jep.2009.01.005
- Graidist P, Purintrabipan J (2009) The biological activity of *Coccinia indica* on glucose transporter 1 (GLUT1) promoter. *Songklanakarin Journal of Science and Technology* 31: 247–253.
- Guha A, Sinha RK, Sinha S (2004) Average packing ratio as a parameter for analyzing the karyotypes of dioecious cucurbits. *Caryologia* 57: 117–120. doi: 10.1080/00087114.2004.10589379
- Guha J, Sen SP (1973) Antigibberellins of the Cucurbitaceae. *Nature New Biology* 244: 223–224. doi: 10.1038/newbio244223a0
- Harvey WH, Sonder OW (1862) *Flora capensis: being a systematic description of the plants of the Cape colony, Caffraria, & Port Natal*. Hodges, Smith, and Co.; A.S. Robertson, Dublin, Ireland; Cape Town, South Africa, 621 pp.
- Hawaiian Department of Agriculture (1992) Hawaii administrative rules 4: 68: Noxious weed rules.
- Holstein N, Renner SS (2010) *Coccinia* (Cucurbitaceae) gains two new species from East Africa, three new synonyms, and one new combination. *Kew Bulletin* 65: 435–441. doi: 10.1007/s12225-010-9229-9
- Holstein N, Renner SS (2011a) *Coccinia intermedia* – a new Cucurbitaceae species from West Africa. *PhytoKeys* 7: 27–36. doi: 10.3897/phytokeys.7.2032
- Holstein N, Renner SS (2011b) A dated phylogeny and collection records reveal repeated biome shifts in the African genus *Coccinia* (Cucurbitaceae). *BMC Evolutionary Biology* 11: 28. doi: 10.1186/1471-2148-11-28
- Hooker JD (1871) Cucurbitaceae. In: Oliver D (Ed.) *Flora of tropical Africa*. L. Reeve & Co., London, UK, 521–569.
- Hora A (1995) *Anchote: an endemic tuber crop*. Jimma College of Agriculture, Jimma, Ethiopia, 75 pp.
- Hossain MZ, Shibib BA, Rahman R (1992) Hypoglycemic effects of *Coccinia indica*: inhibition of key gluconeogenic enzyme, glucose-6-phosphatase. *Indian Journal of Experimental Biology* 30: 418–420.
- Huber C (1864) *Catalogue pour le printemps 1864*. Ch. Huber Frères, Hyères, France.
- Huber C (1865) *Catalogue pour le printemps 1865*. Ch. Huber Frères, Hyères, France.
- Hussain A, Wahab S, Rizvi A, Hussain MS (2011) Macroscopical, anatomical and physico-chemical studies on leaves of *Coccinia indica* Wight & Arn., growing wildly in eastern Uttar Pradesh region of India. *Indian Journal of Natural Products and Resources* 2: 74–80.

- Hutchinson J, Dalziel JM (1928) Tropical African Plants: III. Bulletin of Miscellaneous Information 1928: 221–229.
- Hutchinson L, Dalziel MD, Hepper FN (1954) Flora of West Tropical Africa. Crown Agents, London, UK, 295 pp.
- Imbumi MD (2004) *Coccinia grandis* (L.) Voigt. In: Grubben GJH, Denton OA (Eds) Vegetables/Légumes. PROTA (Plant Resources of Tropical Africa / Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands, 227–229.
- Jacot Guillarmod A (1971) Flora of Lesotho (Basutoland). J. Cramer, Lehre, Germany, 474 pp.
- Jarvis A, Reuter HI, Nelson A, Guevara E (2008) Hole-filled seamless SRTM data V4. International Centre for Tropical Agriculture (CIAT).
- Jeffrey C (1962) Notes on Cucurbitaceae, including a proposed new classification of the family. Kew Bulletin 15: 337–371. doi: 10.2307/4115586
- Jeffrey C (1967) Cucurbitaceae. In: Milne-Redhead E, Polhill RM (Eds) Flora of Tropical East Africa. Crown Agents for Oversea Governments and Administrations, London, UK, 157 pp.
- Jeffrey C (1978) Cucurbitaceae. In: Launert E (Ed.) Flora Zambesiaca. Flora Zambesiaca Managing Committee, London, UK, 414–499.
- Jeffrey C (1985) Further notes on Cucurbitaceae: VII. preliminary to the Flora of Ethiopia. Kew Bulletin 40: 209–211. doi: 10.2307/4108496
- Jeffrey C (1995) Cucurbitaceae. In: Edwards S, Tadesse M, Hedberg I (Eds) Flora of Ethiopia and Eritrea. Addis Abeba University & Uppsala University, Addis Abeba, Ethiopia & Uppsala, Sweden, 17–59.
- Jeffrey C, Fernandes RB (1986) Cucurbitaceae. In: Mendes EJ (Ed.) Flora de Moçambique. Instituto de Investigação Científica Tropical, Lisboa, Portugal, 117 pp.
- Jensen NT (2005) The medical skills of the Malabar doctors in Tranquebar, India, as recorded by surgeon T L F Folly, 1798. Medical history 49: 489–515. doi: 10.1017/S0025727300009170
- Jha U, Asad M, Asdaq SMB, Das AK, Prasad VS (2010) Fertility inducing effect of aerial parts of *Coccinia cordifolia* L. in female rats. Journal of Ethnopharmacology 127: 561–564. doi: 10.1016/j.jep.2009.10.021
- Johns T, Kokwaro JO (1991) Food plants of the Luo of Siaya District, Kenya. Economic Botany 45: 103–113. doi: 10.1007/BF02860055
- Johnston AK (1861) The royal atlas of modern geography. William Blackwood & Sons, Edinburgh & London, UK, 96 pp.
- Jongjitvimol T, Wattanachaiyingcharoen W (2006) Pollen food sources of the stingless bees *Trigona apicalis* Smith, 1857, *Trigona collina* Smith, 1857 and *Trigona fimbriata* Smith, 1857 (Apidae, Meliponinae) in Thailand. The Natural History Journal of Chulalongkorn University 6: 75–82.
- Jongkind CCH (2004) A new species of *Coccinia* (Cucurbitaceae) from West Africa. BLUMEA 49: 83–85. doi: 10.3767/000651904X486205
- Kapoor A, Mahor R, Gupta RN (1981) A new fruit rot disease of kanduri (*Coccinia cordifolia* Linn.). Current Science 50: 281.
- Kéraudren-Aymonin M (1975a) Cucurbitaceae. In: Bamps PRJ (Ed.) Flore d'Afrique centrale. Jardin botanique de Belgique, Meise, Belgium, 152 pp.

- Kéraudren-Aymonin M (1975b) Cucurbitacées. In: Aubréville A, Leroy J-F (Eds) Flore du Cambodge, du Laos et du Viêt-nam. Muséum National d'Histoire Naturelle, Paris, France, 123 pp.
- Kéraudren M (1967) Cucurbitaceae. In: Aubréville A (Ed.) Flore du Cameroun. Muséum National d'Histoire Naturelle, Paris, France, 192 pp.
- Kéraudren M (1968) Deux Cucurbitacées nouvelles du Gabon. *Adansonia* 8: 39–44.
- Khunwasi C (1998) Palynology of the Cucurbitaceae. Ph.D. thesis, Leopold-Franzens-Universität, Innsbruck, Austria.
- Kirkwood JE (1904) Comparative embryology of the Cucurbitaceae. *Bulletin of the New York Botanical Garden* 3: 313–402.
- Koch K (1865) Die neuen Pflanzen von Ch. Huber frères & Co. in Hyères. *Wochenschrift des Vereines zur Beförderung des Gartenbaues in den Königlichen Preussischen Staaten für Gärtnerei und Pflanzenkunde* 8: 51–55.
- Kocyan A, Zhang L-B, Schaefer H, Renner SS (2007) A multi-locus chloroplast phylogeny for the Cucurbitaceae and its implications for character evolution and classification. *Molecular Phylogenetics and Evolution* 44: 553–577. doi: 10.1016/j.ympev.2006.12.022
- Krishnamurthy KV (1984) Plant-insect-fungus association in some plant galls. *Proceedings of the Indian Academy of Sciences - Animal Sciences* 93: 265–273. doi: 10.1007/BF03186245
- Kumar LSS, Deodikar GB (1940) Sex chromosomes of *Coccinia indica* Wight and Arn. *Current Science* 9: 128–130.
- Kumar LSS, Vishveshwaraiah S (1952) Sex mechanism in *Coccinia indica* Wight and Arn. *Nature* 170: 330–331. doi: 10.1038/170330a0
- Kuriyan R, Rajendran R, Bantwal G, Kurpad AV (2008) Effect of supplementation of *Coccinia cordifolia* extract on newly detected diabetic patients. *Diabetes Care* 31: 216–220. doi: 10.2337/dc07-1591
- Lal S (1973) Parthenocarpic fruit development in *Coccinia indica* W. A. *Indian Journal of Horticulture* 30: 453–454.
- Leeuwenberg AJM (1985) A revision of *Carvalhoa* K. Schum. Wageningen Agricultural University Papers 85: 48–55.
- Leins P, Galle P (1971) Entwicklungsgeschichtliche Untersuchungen an Cucurbitaceen-Blüten. *Österreichische Botanische Zeitschrift* 119: 531–548. doi: 10.1007/BF01377505
- Li N, Wu Y, Li L, Zhong Q (2001) A study of seed germination in *Coccinia grandis* and *Amaranthus spinosus*. *Journal of Hainan Normal University (Natural Science)* 14: 14–16.
- Linnaeus C (1747) *Flora Zeylanica*. Sumtu & Literis Laurentii Salvii [Lars Salvius], Holmiae [Stockholm], Sweden, 240 pp.
- Linnaeus C (1753) *Species plantarum: exhibentes plantas rite cognitatas, ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas*. Impensis Laurentii Salvii [Lars Salvius], Holmiae [Stockholm], Sweden, 560 pp.
- Linnaeus C (1763) *Species plantarum, exhibentes plantas rite cognitatas ad genera relatas cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas*. Impensis Direct. Laurentii Salvii [Lars Salvius], Holmiæ [Stockholm], Sweden, 899 pp.

- Martcorena C (1963) Material para una monografia de la morfologia del polen de Cucurbitaceae. Grana Palynologica 4: 78–91. doi: 10.1080/00173136309437861
- Maurice NG, Ramteke PW (2012) Development of *Epilachna vigintioctopunctata* (Fabricius) (Coleoptera: Coccinellidae) on essential and alternative foods. International Journal of Advance Pharmaceutical and Biological Sciences 2: 199–204.
- Mayr E (1942) Systematics and the origin of species, from the viewpoint of a zoologist. Columbia University Press, New York, USA, 334 pp.
- Mazumder PM, Sasmal D, Nambi RA (2008) Antiulcerogenic and antioxidant effects of *Coccinia grandis* (Linn.) Voigt leaves on aspirin-induced gastric ulcer in rats. Indian Journal of Natural Products and Resources 7: 15–18.
- McKay JW (1930) Chromosome numbers in the Cucurbitaceae. Botanical Gazette 89: 416–417. doi: 10.1086/334075
- Meeuse ADJ (1962) The Cucurbitaceae of Southern Africa. Bothalia 8: 1–111.
- Mengesha D, Belew D, Gebreselassie W, Sori W (2012) Growth and yield performance of anchote [*Coccinia abyssinica* (Lam.) Cogn.] in response to contrasting environment. Asian Journal of Plant Sciences 11: 172–181. doi: 10.3923/ajps.2012.172.181
- Mennicken M, Maier W, Oberwinkler F (2005) A contribution to the rust flora (Uredinales) of southern Africa, with an emphasis on Namibia. Mycological Progress 4: 55–75. doi: 10.1007/s11557-006-0109-5
- Ming R, Bendahmane A, Renner SS (2011) Sex chromosomes in land plants. Annual Review of Plant Biology 62: 485–514. doi: 10.1146/annurev-arplant-042110-103914
- Miquel FAW (1855) Flora van Nederlandsch Indië. Friedrich Fleischer, Leipzig, Germany, 1130 pp. doi: 10.5962/bhl.title.93
- Miró M (1995) Cucurbitacins and their pharmacological effects. Phytotherapy Research 9: 159–168. doi: 10.1002/ptr.2650090302
- Moideen K, Sherief SH, Sengottuvelu S, Sivakumar T (2011) Hepatoprotective and antioxidant activity of *Coccinia grandis* root extract against paracetamol induced hepatic oxidative stress in Wistar albino rats. International Journal of Research in Ayurveda & Pharmacy 2: 858–863.
- Motooka P, Castro L, Nelson D, Nagai G, Ching L (2003) Weeds of Hawaii's pastures and natural areas: an identification and management guide. CTAHR Publication, Honolulu, HI, USA, 185 pp.
- Mubalama L (2000) Population and distribution of elephants (*Loxodonta africana africana*) in the central sector of the Virunga National Park, eastern DRC. Pachyderm 28: 44–55.
- Muhammad Ilyas MH (1992) Studies on the extrafloral nectaries in some members of Cucurbitaceae. Acta botanica indica 20: 116–119.
- Munasinghe MAAK, Abeysena C, Yaddhegige IS, Vidanapathirana T, Piyumal KPB (2011) Blood sugar lowering effect of *Coccinia grandis* (L.) J. Voigt: path for a new drug for diabetes mellitus. Experimental Diabetes Research: Article ID 978762. doi: 10.1155/2011/978762
- Muniappan RN, Cruz J, Bamba J (2002) Invasive plants and their control in Micronesia. Micronesica Supplement 6: 85–92.
- Muniappan RN, Reddy GVP, Raman A (2009) *Coccinia grandis* (L.) Voigt (Cucurbitaceae). In: Muniappan RN, Reddy GVP, Raman A (Eds) Biological control of tropical weeds

- using arthropods. Cambridge University Press, New York, USA, 175–182. doi: 10.1017/CBO9780511576348.010
- Murthy DV (1959) Ovipositional habits of *Dacus* sp. (Trypetidae: Diptera). *Current Science* 28: 72–73.
- Nadkarni KM, Nadkarni AK (1976) Dr. K.M. Nadkarni's Indian materia medica with Ayurvedic, Unani-tibbi, Siddha, allopathic, homeopathic, naturopathic & home remedies, appendices & indexes Popular Prakasham, Mumbai, LXXII, 1319 pp.
- Naudin CV (1859) Revue des Cucurbitacées cultivées au Muséum, en 1859. *Annales des Sciences Naturelles Botanique* 12: 79–164.
- Naudin CV (1862) Cucurbitacées cultivées au muséum d'histoire naturelle en 1862. *Annales des Sciences Naturelles Botanique* 18: 159–208.
- Naudin CV (1866) Cucurbitacées nouvelles. *Annales des Sciences Naturelles Botanique* 5: 5–43.
- Nazimuddin S, Naqvi SSH (1984) Cucurbitaceae. In: Nasir E, Ali SI (Eds) *Flora of Pakistan*. Dept. of Botany, University of Karachi, Karachi, Pakistan, 56 pp.
- Niazi J, Singh P, Bansal Y, Goel R (2009) Anti-inflammatory, analgesic and antipyretic activity of aqueous extract of fresh leaves of *Coccinia indica*. *Inflammopharmacology* 17: 239–244. doi: 10.1007/s10787-009-0010-3
- Nieuwenhuis von Üxküll-Güldenbandt M (1907) Extraflorale Zuckerausscheidungen und Ameisenschutz. *Annales du jardin botanique de Buitenzorg* 21: 195–322.
- Njoroge GN, Newton LE (1994) Edible and poisonous species of Cucurbitaceae in the central highlands of Kenya. *Journal of East African Natural History* 83: 101–115. doi: 10.2982/0012-8317(1994)83[101:eapsoc]2.0.co;2
- O'Brian CW, Pakaluk J (1998) Two new species of *Acythopeus* Pascoe (Coleoptera: Curculionidae: Baridinae) from *Coccinia grandis* (L.) Voight (Cucurbitaceae) in Kenya. *Proceedings of the Entomological Society of Washington* 100: 764–774.
- Oberthür C (1878) Études d'entomologie: Faunes entomologiques; descriptions d'insectes nouveaux ou peu connus. Oberthür & son, Rennes, France, 48 pp. doi: 10.5962/bhl.title.9398
- Okoli BE (1984) Wild and cultivated cucurbits in Nigeria. *Economic Botany* 38: 350–357. doi: 10.1007/BF02859015
- Okoli BE, Onofeghara FA (1984) Distribution and morphology of extrafloral nectaries in some Cucurbitaceae. *Botanical Journal of the Linnean Society* 89: 153–164. doi: 10.1111/j.1095-8339.1984.tb01006.x
- Orech FO, Akenga T, Ochora J, Friis H, Aagaard-Hansen J (2005) Potential toxicity of some traditional leafy vegetables consumed in Nyang'oma division, western Kenya. *African Journal of Food and Nutritional Sciences* 5: article 9.
- Owolabi AT, Rabenstein F, Ehrig F, Maiss Edgar M, Vetten HJ (2012) Strains of Moroccan watermelon mosaic virus isolated from *Lagenaria breviflorus* and *Coccinia barteri* in Calabar, Southeastern Nigeria. *International Journal of Virology* 8: 258–270. doi: 10.3923/ijv.2012.258.270
- Pangelinan M (2002) Commonwealth of the Northern Mariana Islands (CNMI). In: Shine C, Reaser JK, Gutierrez AT (Eds) *Prevention and management of invasive alien species: proceedings of a workshop on forging cooperation throughout the Austral-Pacific*. Global Invasive Species Programme, Cape Town, South Africa, 62.

- Pasha MK, Sen SP (1995) Molecular analysis of Cucurbitaceae genome: reassociation kinetic classes and its evolutionary significance. *Biochemical Systematics and Ecology* 23: 399–406. doi: 10.1016/0305-1978(95)00023-N
- Patankar S, Joshi C, Ranade S, Bhawe M, Ranjekar P (1985) Interphase nuclear structure in plants: role of nuclear DNA content and highly repeated DNA sequences in chromatin condensation. *Proceedings of the Indian Academy of Sciences - Plant Sciences* 94: 539–551. doi: 10.1007/BF03053221
- Patel JN, Patel NK (2010) Study of parasite hosts of the genus *Cuscuta* and its traditional uses in Palanpur Taluka, Gujarat, India. *Ethnobotanical Leaflets* 14: 126–135.
- Pattanayak SP, Sunita P (2009) In vivo antitussive activity of *Coccinia grandis* against irritant aerosol and sulfur dioxide-induced cough model in rodents. *Bangladesh Journal of Pharmacology* 4: 84–87. doi: 10.3329/bjp.v4i2.1537
- Perveen A, Qaiser M (2008) Pollen flora of Pakistan - LVI. Cucurbitaceae. *Pakistan Journal of Botany* 40: 9–16.
- Philip S, Ramakrishnan CK, Menon MR (1972) Leaf blight of *Coccinia indica* (Wight & Arn.) caused by *Corynespora cassiicola*. *Agricultural Research Journal of Kerala* 10: 196.
- Phillips EP (1933) A contribution to the flora of the Leribe Plateau and environs: with a discussion on the relationships of the floras of Basutoland, the Kalahari, and the South-eastern regions. *Annals of the South African Museum* 16: 1–472.
- Purcifull DE, Simone GW, Baker CA, Hiebert E (1988) Immunodiffusion tests for six viruses that infect cucurbits in Florida. *Proceedings of the Florida State Horticultural Society* 101: 400–403.
- Ramachandran K, Subramaniam B (1983) Scarlet gourd, *Coccinia grandis*, little-known tropical drug plant. *Economic Botany* 37: 380–383. doi: 10.1007/BF02904197
- Rangaswami G, Chandrasekaran S (1961) *Cercospora* species on cucurbitaceous hosts in South India. I. Identification of species. *Mycologia* 53: 371–377. doi: 10.2307/3756582
- Rangaswami G, Mahadevan A (2004) Diseases of crop plants in India. PHI Learning, New Delhi, India, 548 pp.
- Reddy GVP, Cruz ZT, Muniappan RN (2009) Life-history, host preference and establishment status of *Melittia oedipus* (Lepidoptera: Sesiidae), a biological control agent for *Coccinia grandis* (Cucurbitaceae) in the Mariana Islands. *Plant Protection Quarterly* 24: 27–31.
- Rehm S, Enslin PR, Meeuse ADJ, Wessels JH (1957) Bitter principles of the Cucurbitaceae. VII.—the distribution of bitter principles in this plant family. *Journal of the Science of Food and Agriculture* 8: 679–686. doi: 10.1002/jsfa.2740081203
- Richard A (1847) *Tentamen florae Abyssinicae*. Arthus Bertrand, Paris, France, XI, 472 pp.
- Roemer MJ (1846) *Familiarum naturalium regni vegetabilis synopses monographica*. Landes-Industrie-Comptoir, Vimarier [Weimar], Germany, 222 pp.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology*. doi: 10.1093/sysbio/sys029
- Roy RP (1974) Sex mechanism in higher plants. *Journal of the Indian Botanical Society* 53: 141–155.
- Roy RP, Roy PM (1971a) An intergeneric cross in the Cucurbitaceae (*Coccinia indica* W. and A. × *Bryonopsis laciniosa* Arn.). *Current Science* 40: 46–48.

- Roy RP, Roy PM (1971b) Mechanism of sex determination in *Coccinia indica*. *Journal of the Indian Botanical Society* 50A: 391–400.
- Roy RP, Saran S (1990) Sex expression in the Cucurbitaceae. In: Bates DM, Robinson RW, Jeffrey C (Eds) *Biology and utilization of the Cucurbitaceae*. Comstock Publishing Associates, Ithaca, NY, USA and London, UK, 251–268.
- Ruby J, Nathan PT, Balasingh J, Kunz TH (2000) Chemical composition of fruits and leaves eaten by short-nosed fruit bat, *Cynopterus sphinx*. *Journal of Chemical Ecology* 26: 2825–2841. doi: 10.1023/A:1026446011693
- Rumphius GE (1747) *Herbarium amboinense*. Apud Franciscum Changuion, Hermannum Uytwerf, Haga [Den Haag], Netherlands, 492 pp.
- Ryding O (2001) Reno Muschler and the identity of his African labiates. *Biologiske Skrifter* 54: 155–172.
- Sadikot S (2009) Does this article help or hinder traditional medicine? *International Diabetes Monitor* 21: 26–27.
- Satheesh LS, Murugan K (2011) Antimicrobial activity of protease inhibitor from leaves of *Coccinia grandis* (L.) Voigt. *Indian Journal of Experimental Biology* 49: 366–374.
- Schaefer H, Renner SS (2010) A three-genome phylogeny of *Momordica* (Cucurbitaceae) suggests seven returns from dioecy to monoecy and recent long-distance dispersal to Asia. *Molecular Phylogenetics and Evolution* 54: 553–560. doi: 10.1016/j.ympev.2009.08.006
- Schaefer H, Renner SS (2011a) Cucurbitaceae. In: Kubitzki K (Ed.) *Families and genera of flowering plants*. Springer, Berlin, Germany, 112–174.
- Schaefer H, Renner SS (2011b) Phylogenetic relationships in the order Cucurbitales and a new classification of the gourd family (Cucurbitaceae). *Taxon* 60: 122–138.
- Schrödter K (1926) Zur physiologischen Anatomie der Mittelzelle drüsiger Gebilde. *Flora, oder allgemeine botanische Zeitung* 120: 19–86.
- Schweinfurth GA (1867) *Beitrag zur Flora Aethiopiens*. Georg Reimer, Berlin, Germany, 311 pp.
- Schweinfurth GA (1893) *Abyssinische Pflanzennamen*. *Abhandlungen der königlichen Akademie der Wissenschaften zu Berlin*, 1–84.
- Schweinfurth GA (1915) Dr. Reno Muschlers Fälschungen. *Verhandlungen des Botanischen Vereins der Provinz Brandenburg* 56: 170–175.
- Selby AD (1899) Further studies of cucumber, melon and tomato diseases, with experiments. *Bulletin of the Ohio Agricultural Experiment Station* 105: 217–235.
- Sewram V, Shephard GS, van der Merwe L, Jacobs TV (2006) Mycotoxin contamination of dietary and medicinal wild plants in the Eastern Cape Province of South Africa. *Journal of Agricultural and Food Chemistry* 54: 5688–5693. doi: 10.1021/jf060483b
- Shackleton SE, Dzerefos CM, Shackleton CM, Mathabela FR (1998) Use and trading of wild edible herbs in the central Lowveld savanna region, South Africa. *Economic Botany* 52: 251–259. doi: 10.1007/BF02862142
- Shah JJ, Nair GM, Thanki YJ, Kothari IL (1983) Isolated sieve tubes in the fruit of *Coccinia grandis* (L.) Voigt (Cucurbitaceae). *Annals of Botany* 51: 251–253.
- Shaheen SZ, Bolla K, Vasu K, Charya MAS (2009) Antimicrobial activity of the fruit extracts of *Coccinia indica*. *African Journal of Biotechnology* 8: 7073–7076.

- Shibib BA, Khan LA, Rahman R (1993) Hypoglycaemic activity of *Coccinia indica* and *Momordica charantia* in diabetic rats: depression of the hepatic gluconeogenic enzymes glucose-6-phosphatase and fructose-1,6-bisphosphatase and elevation of both liver and red-cell shunt enzyme glucose-6-phosphate dehydrogenase. *Biochemical Journal* 292: 267–270.
- Siddiqi MR (2000) Tylenchida: parasites of plants and insects. CABI Publishing, Egham, UK, 848 pp. doi: 10.1079/9780851992020.0000
- Siebert F, Eckhardt HC, Siebert SJ (2010) The vegetation and floristics of the Letaba exclosures, Kruger National Park, South Africa. *Koedoe* 52: article 777. doi: 10.4102/koedoe.v52i1.777
- Sims J (1816) *Bryonia quinqueloba*. Five-fingered Cape Bryony. *Curtis's Botanical Magazine* 43: 1820.
- Solereider H (1899) Systematische Anatomie der Dicotyledonen. Ferdinand Enke, Stuttgart, Germany, 984 pp.
- Stafleu FA, Cowan RS (1988) Taxonomic literature. Bohn, Scheltema & Holkeman / Dr. W. Junk b.v., Publishers, Utrecht, Antwerpen / The Hague, Netherlands, Boston, USA, 653 pp.
- Stamatakis A (2006) RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22: 2688–2690. doi: 10.1093/bioinformatics/btl446
- Stanford CB, Nkurunungi JB (2003) Behavioral ecology of sympatric chimpanzees and gorillas in Bwindi Impenetrable National Park, Uganda: Diet. *International Journal of Primatology* 24: 901–918. doi: 10.1023/A: 1024689008159
- Starr F, Starr K, Loope LL (2010) New plant records from the Hawaiian Archipelago. *Bishop Museum Occasional Papers* 107: 61–68.
- Stearn WT (2004) Botanical Latin. Timber Press, Portland, OR, USA, 560 pp.
- Sutaria RN (1936) On the chromosome number in some cultivated and wild cucurbits of Gujarat. *Journal of the University of Bombay* 4: 21–28.
- Thanki YJ (1989) Structure and development of hairs in cucurbitaceous fruits. *Feddes Repertorium* 100: 377–379.
- Thulin M (2009) New species of *Coccinia* and *Momordica* (Cucurbitaceae) from north-eastern tropical Africa. *Kew Bulletin* 64: 485–489. doi: 10.1007/s12225-009-9124-4
- Uchida GK, Vargas RI, Beardsley JW, Liquido NJ (1990) Host suitability of wild cucurbits for melon fly, *Dacus cucurbitae* Coquillett, in Hawaii, with notes on their distribution and taxonomic status. *Proceedings of the Hawaiian Entomologic Society* 30: 37–52.
- Umamaheswari M, AsokKumar K, Somasundaram A, Sivashanmugam T, Subhadradevi V, Ravi TK (2007) Xanthine oxidase inhibitory activity of some Indian medical plants. *Journal of Ethnopharmacology* 109: 547–551. doi: 10.1016/j.jep.2006.08.020
- Umamaheswari M, Chatterjee TK (2008) *In vitro* antioxidant activities of the fractions of *Coccinia grandis* l. leaf extract. *African Journal of Traditional, Complementary and Alternative Medicines* 5: 61–73. doi: 10.4314/ajtcam.v5i1.31258
- Unni PN, Raghavan P, Philip VJ (1976) A histopathological study of anomalous growth in the stem of *Coccinia indica* W. & A. infested with *Neolasioptera cephalandrae* Mani. *Annals of Botany* 40: 493–497.
- Urban I (1921) Sertum antillanum X. *Repertorium specierum novarum regni vegetabilis* 17: 6–11.

- Vadivu R, Krithika A, Biplab C, Depeepya P, Shoeb N, Lakshmi KS (2008) Evaluation of hepatoprotective activity of the fruits of *Coccinia grandis* Linn. *International Journal of Health Research* 1: 163–168.
- van der Meer FW, Garnett HM (1987) Purification and identification of a South African isolate of Watermelon Mosaic Virus – Morocco. *Journal of Phytopathology* 120: 255–270. doi: 10.1111/j.1439-0434.1987.tb04440.x
- van der Pijl L (1982) Principles of dispersal in higher plants. Springer, Berlin, Germany, 214 pp. doi: 10.1007/978-3-642-87925-8
- Verma VS, Singh S, Padma R (1983) Mosaic disease of *Coccinia indica* Wight and Arn. *Acta Agronomica Academiae Scientiarum Hungaricae* 32: 182–183.
- Voigt JO (1845) Hortus Suburbanus Calcuttensis: A catalogue of the plants which have been cultivated in the Hon. East India company's Botanical Garden, Calcutta, and in the Serampore Botanical Garden, genera. Bishop's College Press, Calcutta, India, 747 pp.
- von Aufseß H (1973) Mikroskopische Darstellung des Verholzungsgrades durch Färbemethoden. *Holz als Roh- und Werkstoff* 31: 24–33. doi: 10.1007/bf02608218
- von Bunge A (1854) Beitrag zur Kenntniss der Flor Russlands und der Steppen Central-Asiens. Mémoires présentés a l'académie impériale des sciences de St-Petersbourg par divers savants et lus dans ses assemblées 7: 1–535.
- Wada M (1930) Über Citrullin, eine neue Aminosäure im Presssaft der Wassermelone, *Citrullus vulgaris* Schrad. *Biochemische Zeitschrift* 224: 420.
- Wasantwisut E, Viriyapanich T (2003) Ivy Gourd (*Coccinia grandis* Voigt, *Coccinia cordifolia*, *Coccinia indica*) in human nutrition and traditional applications. In: Simopoulos AP, Gopalan C (Eds) Plants in human health and nutrition policy. Karger, Basel, Switzerland, 60–66. doi: 10.1159/000069929
- Weiersbye IM, Witkowski ETF (2003) Seed fate and practical germination methods for 46 perennial species that colonize gold mine tailings and acid mine drainage-polluted soils in the grassland biome. In: Seydack AHW, Vorster T, Vermeulen WJ, van der Merwe IJ (Eds) Natural Forests and Woodlands Symposium III : Multiple Use Management - Policy Refinements and Scientific Progress. Department of water affairs and forestry, Berg-en-Dal, Kruger National Park, South Africa, 221–255.
- White F (1962) A history of plant collecting in Northern Rhodesia. In: Fernandes A (Ed.) Comptes rendus de la IV^e réunion plénière de l'association pour l'étude taxonomique de la flore d'Afrique tropicale. Junta de Investigações do Ultramar, Lisbon, Portugal, 175–203.
- White F (1983a) Unesco/AETFAT/UNSO vegetation map of Africa. UNESCO, Paris, France, 1: 5.000.000.
- White F (1983b) The vegetation of Africa. Unesco, Paris, France, 356 pp.
- WHO (2009) Global prevalence of vitamin A deficiency in populations at risk 1995–2005. WHO global database on vitamin A deficiency. World Health Organization, Geneva, Switzerland, 55 pp.
- Wight R, Arnott GAW (1834) Prodrromus Florae Peninsulae Indiae Orientalis: containing abridged descriptions of the plants found in the peninsula of British India, arranged according to the natural system. Parbury, Allen & Co., London, UK, 480 pp.

- Wilson HH (1867) The Megha-dūta (Cloud messenger) by Kālidāsa. Trübner & Co., London, UK, 179 pp.
- Wu C-Y, Raven PH, Hong D-Y (2011) Flora of China. Vol. 19 (Cucurbitaceae through Valerianaceae, with Annonaceae and Berberidaceae). Science Press/Missouri Botanical Garden Press, Beijing, China/St. Louis, MO, USA, 884 pp.
- Xu Y-K, Liu H-M, Dao X-S (2003) The nutritional contents of *Coccinia grandis* and its evaluation as a wild vegetable. Acta Botanica Yunnanica 25: 680–686.
- Xu Y-K, Liu H-M, Xiao C-F, Chan Y, Wu Z-L, Dao X-S, Cai C-T (2004) An analysis of selenium contents of wild vegetable in Xishuangbanna. Acta Botanica Yunnanica 26: 668–672.
- Zahur MS (1962) Early ontogeny of the ovule in *Coccinia indica*. Biologia 8: 179–197.
- Zaman MR (2009) Effect of pH on *in vitro* pollen germination of fourteen cultivated and wild species of cucurbit. Journal of Bio-Science 17: 129–133. doi: 10.3329/jbs.v17i0.7120
- Zimmermann A (1922a) Die Cucurbitaceen 1. Gustav Fischer, Jena, Germany, 206 pp.
- Zimmermann A (1922b) Die Cucurbitaceen 2. Gustav Fischer, Jena, Germany, 186 pp.
- Zimmermann JG (1932) Über die extrafloralen Nektarien der Angiospermen. Beihefte zum Botanischen Centralblatt 44: 99–196.

Supplementary material 1

List of examined specimens and geo-references of the localities

Authors: Norbert Holstein

Data type: specimens data.

Explanation note: Voucher information and (predominantly inferred) coordinates of geo-references (in decimal degrees). Specimens included in this study, with the geographic origin of material (only country and 1st administrative division given).

Herbarium acronyms follow Index Herbariorum; barcodes or other unique identifiers in brackets behind the acronym. Indication of digitized images refer to the homepage of the corresponding institution indicated by the herbarium acronym, except for JPS (JSTOR plant science) and CVH (= Chinese Virtual Herbarium).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Supplementary material 2

GenBank accessions

Authors: Norbert Holstein

Data type: specimens data.

Explanation note: Voucher information and GenBank accession numbers. Specimens included in this study, with the geographic origin of material (only country and 1st administrative division given) and GenBank accession numbers for all sequences. Herbarium acronyms (in parentheses behind the voucher name) follow Index.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.